

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/361965887>

# Two New Pentacyclic Triterpenoids, an Alkaloid and a Long-chain Fatty Acid from *Albizia Coriaria* (Welw ex. Oliver)

Article in *Journal of Chemistry* · July 2022

DOI: 10.17721/fujcV10I1P128-141

CITATION

1

READS

65

3 authors:



**Timothy Omara**

University of Natural Resources and Life Sciences Vienna

97 PUBLICATIONS 582 CITATIONS

[SEE PROFILE](#)



**Ambrose Kiprof**

Moi University

60 PUBLICATIONS 493 CITATIONS

[SEE PROFILE](#)



**Viola Jepchumba Kosgei**

University of Lorraine

17 PUBLICATIONS 102 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Plants Used in Antivomom Therapy in Rural Kenya: Ethnobotany and Future Perspectives [View project](#)



Design and Assembly of a Domestic Water Temperature, pH and Turbidity Monitoring System [View project](#)

## **Two New Pentacyclic Triterpenoids, an Alkaloid and a Long-chain Fatty Acid from *Albizia Coriaria* (Welw ex. Oliver)**

Timothy Omara<sup>a,b,c\*</sup>, Ambrose K. Kiprop<sup>a,b</sup>, Viola J. Kosgei<sup>a,b</sup>

<sup>a</sup> *Africa Center of Excellence II in Phytochemicals, Textile and Renewable Energy (ACE II PTRE),  
Moi University, P. O. Box 3900-30100, Eldoret, Kenya*

<sup>b</sup> *Department of Chemistry and Biochemistry, School of Sciences and Aerospace Studies, Moi  
University, P. O. Box 3900-30100, Eldoret, Kenya.*

<sup>c</sup> *Department für Chemie, Universität für Bodenkultur Wien, Vienna Gregor-Mendel-Straße 331180  
Vienna, Austria.*

[prof.timo2018@gmail.com](mailto:prof.timo2018@gmail.com), [prof.timo2018@mu.ac.ke](mailto:prof.timo2018@mu.ac.ke)

**Keywords:** *Albizia*, *triterpenes*, *pterin-6-carboxylic acid*, *oleanolic acid*, *undecanol*.

Herein, we report the occurrence of four new compounds in ethanolic extract of *Albizia coriaria* Welw ex. Oliver leaves along with other compounds previously reported in this species. The compounds were isolated and characterized using column chromatography, Fourier Transform Infrared (FTIR) and gas chromatography/mass spectrometry. FTIR spectrum of the extract showed phenolic O–H stretching ( $3362.30\text{ cm}^{-1}$ ), C=O ( $1660.08\text{ cm}^{-1}$ ), C–O stretching ( $1369.46\text{ cm}^{-1}$  and  $1319.00\text{ cm}^{-1}$ ) and C–N stretch ( $1072.44\text{ cm}^{-1}$ ) which confirmed the presence of alcohols, carboxylic acids and nitrogen-containing compounds. Oleanolic acid (**1**), oleanolic acid acetate (**2**), pterin-6-carboxylic acid (**3**), undecanol (**4**), betulinic acid (**5**), betulin (**6**) and benzyl alcohol (**7**) were tentatively identified in the extract. Compounds **1-4** are being reported for the first time in *Albizia coriaria*.

### **Introduction**

Many lead compounds and therapeutic molecules have been discovered from natural products. Plants represent a huge proportion of natural sources of drugs currently used in both human and veterinary medicine [1]. Species emanating from the genus *Albizia* are widely used in traditional medicine, owing to their inherent possession of bioactive compounds [2]. *Albizia coriaria* Welw ex. Oliver (*A. coriaria*) is

a representative species of this genus that is widely utilized in African herbal medicine [3]. Its organs and the whole plant are used for managing various ailments and conditions including dermatological diseases, gastrointestinal diseases and infections, ophidian bites, malignancies, malaria, cough (tuberculosis), cardiovascular diseases, anaemia, venereal diseases and reproductive conditions [4-30]. It is also used in the treatment of lungworms/ascaris worms in

cattle, sheep and goats, as a mosquito repellent and toothbrush [18, 25, 31-33].

Various bioactivities of the stem bark, leaves, flowers and roots of this species have been validated *in vitro*. These include molluscicidal [34], antimicrobial [7, 35-42], antiproliferative [43], anti-giardial [23], antiplasmodial [44], anti-inflammatory [45], antimycobacterial [46, 47] and antioxidant activities [36, 37, 40].

Investigation of the chemical composition of *A. coriaria* stem bark has led to the identification of lupeol, lupenone, benzyl alcohol, betulinic acid, (+)-catechin and acacic acid lactone [39], and 43 other compounds including hexanal, (2,4-dinitrophenyl)hydrazine; hydroquinone and 7-hydroxy-3'-methoxyflavone, trimethylsilyl ether [46]. In the root bark extracts, Gummiferaoside C and Coriariosides A-E were earlier characterized [43, 48]. Recently, we reported the presence of lupenone, lupeol, betulinic acid, betulin and benzyl alcohol in the leaves of this species sampled from Mbarara district of Western Uganda [49]. The current study reports for the first time, the occurrence of two pentacyclic triterpenoids (oleanolic acid and oleanolic acid acetate), an alkaloid (pterin-6-carboxylic acid), a long-chain fatty acid (undecanol), along with betulinic acid, betulin and benzyl alcohol in ethanolic extract of *A. coriaria* leaves from Kole district of Northern Uganda, East Africa.

## **Experimental part**

### *Sample collection and preparation*

*A. coriaria* leaves were sampled from Otangula village, Kole district, Uganda (2°17'35.0''N 32°46'31.2''E) on Wednesday 20<sup>th</sup> January 2021. A specimen voucher (no. 50995) was authenticated by a taxonomist at Natural Chemotherapeutics Research Institute (NCRI), Kampala, Uganda. It was deposited at Department of Botany, Makerere University herbarium, Kampala, Uganda for future reference.

Owing to the hot weather in Uganda in January 2021, the laboratory sample was washed under running tap water to remove dusts and then shade-dried for two days in Otino Waa Secondary School Biology Laboratory, Kole, Uganda. This was done to avoid sample deterioration prior to transportation for further drying. The sample was then packaged in sterilized polyethylene bags and transported to School of Sciences and Aerospace Studies, Chemistry Laboratory, Moi University, Eldoret, Kenya where they were dried under shade at room temperature for 3 weeks.

The dry leaves were powdered using a NutriBullet® 600 Series electric grinder (Capbran Holdings, USA) and then 0.5 kg of it was serially macerated in 1000 ml of ethyl acetate and ethanol at room temperature. The extracts were separately filtered through cotton wool and then Whatman No.1 filter paper. They were eventually concentrated to dryness by

rotary evaporation at 40 °C. The crude extracts were transferred to a desiccator of anhydrous sodium sulphate to remove any traces of water.

*Fourier transform infrared scanning of the crude extract*

The ethanolic extract was subjected to Attenuated Total Reflection-Fourier transform infrared (ATR-FTIR) characterization. The scan was done on a Nicolet 6700 FTIR spectrophotometer (Thermo Scientific, USA) at a spectral resolution of 4 cm<sup>-1</sup> and scan range of 500 cm<sup>-1</sup> to 4000 cm<sup>-1</sup>.

*Thin layer chromatography and column chromatography of the crude extract*

Thin layer chromatography (TLC) was used to optimize the solvent ratios for column chromatography [50]. Briefly, the ethanolic extract was reconstituted in ethanol (1: 20, w/v), spotted onto TLC plates, dried and developed in solvent systems of different hexane: ethyl acetate and ethyl acetate: ethanol ratios. The number of compounds and their retention factors (R<sub>f</sub>) were calculated after visualization under UV light (254 nm and 365 nm).

The extract was subjected to silica gel (60-120 mesh) column chromatography (7.5 × 100 cm column) using the optimized solvent ratios. Silica gel (100 g) was mixed with hexane: ethyl acetate solvent system to form a slurry before it was poured into the column. The sample was introduced into the column and then eluted with

the established solvent systems [49]. For every elution, equal fractions were collected and those with similar TLC profiles were combined and concentrated to dryness by rotary evaporation.

*Characterization of compounds in the fractions*

Seven fractions from chromatographic isolation and TLC profiling were re-dissolved in dichloromethane/methanol (1:1). They were separately filtered through 0.45 µm filters and then transferred to 2 mL vials for gas chromatography-mass chromatography (GC-MS) characterization as previously described [49].

Briefly, the characterization was performed in a gas chromatograph interfaced with mass spectrometer triple-quad system (Agilent 8890A GC and Agilent 5977 GC/MSD, Agilent Technologies) supplied with an Agilent 7693A automatic liquid sampler, a National Institute of Standards and Technology (NIST) library, an installed Mass Hunter Workstation software and an HP-5MS ultra inert column (30 m × 0.25 mm × 0.25 µm).

Compound identification was based on the gas chromatogram elution times, mass spectra matching with those of the compounds in NIST 11 spectral library and/or compared with published spectroscopic data. Compound molecular ions were established using Nitrogen rule [51]. Further, the identified molecular ions were individually analyzed for their capacity to


yield important ions in the high mass region by logical neutral losses [52].

## Results and discussion

### Fourier Transform infrared spectrum of the crude extract

The FTIR spectrum of ethanolic extract of *A. coriaria* leaves (**Figure 1**) had a stretch at  $3362.30\text{ cm}^{-1}$  which could be assigned to O-H

stretching of phenolics [53]. The peaks observed at  $2358.93\text{ cm}^{-1}$ ,  $1660.08\text{ cm}^{-1}$ ,  $1369.46\text{ cm}^{-1}$  and  $1319.00\text{ cm}^{-1}$ ,  $1370.12\text{ cm}^{-1}$  and  $1072.44\text{ cm}^{-1}$  are due to O=C=O stretching [54], conjugated or aromatic C=O bond [55], C–O stretching [56] and C–N stretching [57], respectively. This confirmed the presence of alcohols, carboxylic acids and nitrogenous compounds in the extract.



**Figure 1.** FTIR spectrum of ethanolic extract of *A. coriaria* leaves from Kole district, Northern Uganda.

### Compounds tentatively identified in ethanolic extract of *A. coriaria* leaves

Chromatographic fractions 6–19 eluted with hexane: ethyl acetate (4:6) afforded a white solid which was analyzed and found to contain oleanolic acid (**1**,  $R_f = 0.570$ ). Fractions 22–30 gave crystals which contained Oleanolic acid acetate (**2**,  $R_f = 0.533$ ). Further elution of the column yielded fractions 35–48 and 52–66 which contained Pterin-6-carboxylic acid (**3**,  $R_f = 0.490$ ) and Undecanol (**4**,  $R_f = 0.440$ ), respectively. Further chromatographic elution

using 30% ethyl acetate in ethanol gave fractions 2–17, 20–33 and 37–46 which afforded betulinic acid (**5**,  $R_f = 0.596$ ), betulin (**6**,  $R_f = 0.500$ ) and benzyl alcohol (**7**,  $R_f = 0.480$ ), respectively (**Figure 2**). Of the seven compounds identified, **1-4** are reported for the first time in *A. coriaria*.

### Compound 1 (Oleanolic acid)

Compound **1** was obtained as a white solid, soluble in ethanol. Its GC retention time was 31.875 minutes, with a molecular ion at  $m/z$  456 which suggested a molecular formula  $C_{30}H_{48}O_3$ . The mass spectrum of compound **1**

(Figure 3) had a molecular ion peak at  $m/z$  455, which corresponded to a deprotonated oleanolic acid molecule [58-60]. The spectral data also showed presence of a carboxylic acid group at  $m/z$  407, that strongly suggested a C-12 unsaturated pentacyclic triterpene containing a carboxylic acid group in either ring D or E [61]. Another peak was observed at  $m/z$  391 due to loss of  $-CH_2$  group from the  $m/z$  407 fragment (Figure 4)[62, 63]. The other peaks observed at  $m/z$  255, 269, 283 and 297 are fragments of the enhanced products from the molecular ion [64]. Thus, compound 1 was tentatively deduced to be oleanic acid/oleanolic acid.




Figure 2. Structure of compounds identified in ethanolic extract of *A. coriaria* leaves from Kole district, Uganda.





Figure 4. Fragmentation pattern of compound 1 (Oleanolic acid)

Free oleanolic acid is being reported for the first time in *A. coriaria*. However, it has been previously reported in the ethanolic extracts of *Albizia julibrissin* stem bark [65], a sister species in the *Albizia* genus. Oleanolic acid is one of the best known pentacyclic triterpenoids with widespread occurrence throughout Kingdom plantae in the form of free acid or aglycones for triterpenoid saponins [58, 66].


#### Compound 2 (Oleanolic acid acetate)

Compound 2, eluted with ethyl acetate: ethanol (3:7) had a GC retention time of 30.992 minutes. Its mass spectrum (Figure 5) had a molecular ion at  $m/z$  497, with a molecular formula  $C_{32}H_{50}O_4$ . Fragments  $m/z$  269 and  $m/z$  423, characteristic of pentacyclic triterpenes possessing carboxylic acid groups were present [67, 68]. The spectrum further suggested the presence of a carboxylic acid group,  $[M-COOH]^+$  at  $m/z$  301 [67, 68]. The ion peak observed at  $m/z$  469 could have been due to loss

of HCHO from the molecular ion (**Figure 6**). Further loss of HCHO and H<sub>2</sub>O molecules as in oleanolic acid could have given the peak observed at  $m/z$  423 [67, 68]. These spectral characteristics supported the NIST library suggestion that compound **2** is 3-*O*-acetyloleanolic acid (oleanolic acid acetate)[69].



**Figure 3.** Expanded mass spectrum of compound **1** (Oleanolic acid)



**Figure 5.** Mass spectrum of compound **2** (Oleanolic acid acetate)




Figure 6. Fragmentation pattern for major ions of compound 2 (Oleanolic acid acetate)

**Compound 3 (Pterin-6-carboxylic acid)**

Compound 3 was eluted with hexane: ethyl acetate (4:6) from the column. It had a GC

retention time of 21.579 minutes. The molecular ion of compound 3 occurred at  $m/z$  207, suggesting a molecular formula  $C_7H_5N_5O_3$  (Figure 7).





Figure 7. Mass spectrum of compound 3 (pterin-6-carboxylic acid)

The fragment observed at  $m/z$  163 is due to loss of carbon dioxide ( $CO_2$ , mass = 44) from the molecular ion [70]. Loss of  $N_2HCHO$  (mass = 58) from  $m/z$  163 (Figure 8) could have yielded the fragment at  $m/z$  105 as observed in the fragmentation of pterins [71]. The fragment at  $m/z$  149 is due to loss of glycolaldehyde gas ( $HOCH_2CHO$ , mass = 59) from the molecular

ion [71]. Loss of  $H_2O$  from  $m/z$  149 gives the fragment at  $m/z$  131. The other peaks at  $m/z$  57, 69, 105 and 122 are usually observed in the spectrum of pterin-6-carboxylic acid [72-75]. Thus, compound 3 was suggested to be 2-amino-4-hydroxy-6-pteridine carboxylic acid (pterin-6-carboxylic acid) as initially indicated by NIST 11 library matching. Compound 3, a typical



alkaloid, was previously characterized in *Albizia lebbbeck* leaf extracts [75].



**Figure 8.** Fragmentation pattern of compound 3 (pterin-6-carboxylic acid)


#### Compound 4 (Undecanol)

Compound 4 was obtained as a pale greenish liquid with mild odor, soluble in ethanol. Its GC retention time was 15.307 minutes. The mass spectrum of compound 4 (**Figure 9**) had a base peak at  $m/z$  55 and a molecular ion at  $m/z$  172, which suggested a


molecular formula  $C_{11}H_{24}O$ . The fragment observed at  $m/z$  154 is due to loss of water from the molecular ion [76, 77]. The fragment at  $m/z$  125 is due to loss of water and vinyl ( $CH_2 = CH$ ) group from the molecular ion [76, 78]. Loss of ethene ( $CH_2=CH_2$ ) molecule from  $m/z$  125 gave the fragment at  $m/z$  97 (**Figure 10**).

The other fragments observed at  $m/z$  at 83, 69, 55, 41 and 27 are due to loss of  $-CH_2$  from the preceding fragments ( $m/z$  97, 83, 69, 55 and 41, respectively)[79]. These spectral patterns confirmed that compound 4 is Undecan-1-ol (undecanol or 1-undecanol). Though it has never been reported before in *A. coriaria*, undecanol was earlier reported in *A. zygia* stem bark fixed oil analyzed by GC-MS [80].

The other compounds: betulinic acid (5), betulin (6) and benzyl alcohol (7) were identified as reported previously [49].



**Figure 9.** Mass spectrum of compound 4 (Undecanol).



**Figure 10.** Fragmentation pattern of compound 4 (Undecanol)

All in all, the compounds identified in the ethanolic extract support the traditional use of this species in the management of bacterial diseases and oxidative stress-mediated complications in Northern Uganda. For example, compounds **1** and **2** have been reported to elicit antibacterial activity against *Staphylococcus aureus* and *Pseudomonas aeruginosa* [66, 68, 81-83]. Compounds **3** and **4** had antibacterial activity [75, 84], though their antioxidant activity has not been reported. The other compounds (**5-7**) have well documented bioactivities [49]. The occurrence of bioactive compounds **1-4** in this species, further lends credence to its wide use in traditional medicine in Uganda and across Africa. It also confirms that there is intraspecific variation in the bioactive compounds present in this species.

## Conclusions

Isolation and characterization of the ethanol extract of *A. coriaria* leaves from Kole

district of Uganda resulted in the identification of two triterpenes: oleanolic acid (**1**) and oleanolic acid acetate (**2**); an alkaloid (pterin-6-carboxylic acid **3**), and an organic alcohol (undecanol **4**), along with betulinic acid (**5**), betulin (**6**) and benzyl alcohol (**7**). Compounds **1-4** have been identified for the first time in this species. The isolated compounds possess an array of pharmacological activities, lending credence to the use of the leaves in Ugandan herbal medicine.

## Acknowledgements

The authors are grateful to the World Bank and the Inter-University Council of East Africa (IUCEA) through Africa Center of Excellence II in Phytochemicals, Textile and Renewable Energy (ACE II PTRE) hosted at Moi University, Kenya for funding this study (Credit No. 5798-KE).

## References

- [1] Atanasov AG, Zotchev SB, Dirsch VM, International Natural Product Sciences Taskforce, Supuran CT. Natural products in drug discovery: advances and opportunities. *Nat Rev Drug Discov.* 2021;20(3):200–216.
- [2] He Y, Wang Q, Ye Y, Liu Z, Sun H. The ethnopharmacology, phytochemistry, pharmacology and toxicology of genus *Albizia*: A review. *J Ethnopharmacol.* 2020;257:112677.
- [3] Omara T, Kiprop AK, Kosgei VJ. *Albizia coriaria* Welw ex Oliver: a review of its ethnobotany, phytochemistry and ethnopharmacology. *Adv Tradit Med.* 2021.
- [4] Namukobe J, Kasenene JM, Kiremire BT, Byamukama R, Kamatenesi-Mugisha M, Krief S et al. Traditional plants used for medicinal purposes by local

- communities around the northern sector of Kibale National Park, Uganda. *J Ethnopharmacol.* 2011;136:236–245.
- [5] Anywar G, Kakudidi E, Byamukama R, Mukonzo J, Schubert A, Oryem-Origa H. Indigenous traditional knowledge of medicinal plants used by herbalists in treating opportunistic infections among people living with HIV/AIDS in Uganda. *J Ethnopharmacol.* 2020;246:112205.
- [6] Omara T, Kiprop AK, Ramkat RC, Cherutoi J, Kagoya S, Nyangena DM et al (2020b) Medicinal plants used in traditional management of cancer in Uganda: a review of ethnobotanical surveys, phytochemistry, and anticancer studies. *Evid Based Complement Alternat Med.* 2020;2020:1-26.
- [7] Olila D, Ziraba BR, Kamoga D. Bio-prospective studies on medicinal plants used to manage poultry diseases in the Mount Elgon region of Uganda. *Res J Pharmacol.* 2007;1:56–60.
- [8] Orodho JA, Kirimuhuzya C, Otieno JN, Magadula JJ, Okemo P. Local management of tuberculosis by traditional medicine practitioners in Lake Victoria region. *Open Complement Med J.* 2011;3:1–9.
- [9] Schultz F, Anywar G, Wack B, Quave CL, Garbe L. Ethnobotanical study of selected medicinal plants traditionally used in the rural greater Mpigi region of Uganda. *J Ethnopharmacol.* 2020;256:112742.
- [10] Tugume P, Kakudidi EK, Buyinza M, Namaalwa J, Kamatenesi M, Mucunguzi P, Kalema J. Ethnobotanical survey of medicinal plant species used by communities around Mabira Central Forest Reserve, Uganda. *J Ethnobiol Ethnomed.* 2016;12(1):5.
- [11] Bunalema L, Obakiro S, Tabuti JRS, Waako P. Knowledge on plants used traditionally in the treatment of tuberculosis in Uganda. *J Ethnopharmacol.* 2014;151(2):999–1004.
- [12] Nanyunja KR. Indigenous knowledge of the abundance of medicinal and food plants in Mount Moroto forest reserve. Proceedings of the 11th World Forestry Congress. 2003.
- [13] Namukobe J, Lutaaya A, Asiimwe S, Byamukama R. An ethnobotanical study of medicinal plants used in the management of dermatological disorders in Buyende and Kayunga Districts, Uganda. *Eur J Med Plants.* 2021;32:15–40.
- [14] Kyazike E. Exploring the preference for indigenous medicinal plant medicine in Buliisa District, Western Uganda. *Inkanyiso.* 2021;13:77-104.
- [15] Katuura E, Omujal F, Tumusiime RH, Nabukalu D, Byamukama R. Documentation of indigenous knowledge on medicinal plants used to manage common influenza and related symptoms in Luwero district, central Uganda. *J Med Plants Res.* 2016;10(39):705–71.
- [16] Omara T, Kagoya S, Openy A, Omute T, Ssebulime S, Kiplagat KM, Bongomin O. Antivenin plants used for treatment of snakebites in Uganda: ethnobotanical reports and pharmacological evidences. *Trop Med Health.* 2020;48(1):6.
- [17] Ochwang'i DO, Kimwele CN, Oduma JA, Gathumbi PK, Mbaria JM, Kiama SG. Medicinal plants used in treatment and management of cancer in Kakamega County, Kenya. *J Ethnopharmacol.* 2014;151:1040–1055.
- [18] Olala CN. Identification of plants used for treatment of malaria and factors influencing their use in Boro division, Siaya county, Kenya. MSc Thesis, Kenyatta University, Nairobi, Kenya. 2014.
- [19] Omara T. Antimalarial plants used across Kenyan Communities. *Evid-Based Complement Altern Med.* 2020;2020:1-31.
- [20] Shiracko N, Owuor BO, Gakuubi MM, Wanzala W. A survey of ethnobotany of the AbaWanga people in Kakamega county, Western province of Kenya. *Indian J Tradit Knowl.* 2016;15:93–102.
- [21] Geissler PW, Harris SA, Prince RJ, Olsen A, Odhiambo RA, Oketch-Rabah H et al. Medicinal plants

- used by Luo mothers and children in Bondo district, Kenya. *J Ethnopharmacol.* 2002;83:39–54.
- [22] Jeruto P, Mutai C, Ouma G, Lukhoba C, Nyamaka RL, Manani SD. Ethnobotanical survey and propagation of some endangered medicinal plants from south Nandi district of Kenya. *J Anim Plant Sci.* 2010;8(3):1016–1043.
- [23] Johns T, Faubert GM, Kokwaro JO, Mahunnah RLA, Kimanani EK. Antigiardial activity of gastrointestinal remedies of the Luo of East Africa. *J Ethnopharmacol.* 1995;46:17–23.
- [24] Orwa JA, Mwitari PGM, Mtu E, Rukunga GM. Traditional healers and the management of malaria in Kisumu District, Kenya. *East Afr Med J.* 2007;84:51-55.
- [25] Araya YN. Contribution of trees for oral hygiene in east Africa. *Ethnobot Leaflets.* 2007;11:38–44.
- [26] ICRAF. A Selection of Useful Trees and Shrubs for Kenya. International Centre for Research in Agroforestry. Nairobi: Kenya. 1992.
- [27] Johns T, Kokwaro JO, Kimanani EK. Herbal remedies of the Luo of Siaya District, Kenya: establishing quantitative criteria for consensus. *Econ Bot.* 1990; 44(3):369–381.
- [28] Leiderer. La médecine traditionnelle chez les Bekpak (Bafia) du Cameroun. St. Augustin, Deutschland: Haus Volker und kulturen. 1982.
- [29] Tsabang N, Yedjou CG, Tchounwou PB. Phytotherapy of high blood pressure in three phytogeographic regions of Cameroon. *Pharmaceut Anal Acta.* 2017;7:530.
- [30] Abba A, Dogara AM. Ethnomedicinal Survey of Plants used for Management of Inflammatory Diseases in Ringim Local Government, Jigawa State, Nigeria. *Ethnobot Res Appl.* 2021;22:47.
- [31] Dharani N, Yenesew A, Aynekulu E, Tuei B, Jamnadas R (2015). Traditional ethnoveterinary medicine in East Africa: a manual on the use of medicinal plants. Dawson IK ed. The World Agroforestry Centre (ICRAF), Nairobi, Kenya, 199 pages.
- [32] Bossard E. Angolan medicinal plants used also as piscicides and/or soaps. *J Ethnopharmacol.* 1993;40:1-19.
- [33] Sirama VO. Ethnobotanical, Bioactivity and Phytochemical Evaluation of Anthelmintic Herbal Remedies of Migori County, Kenya. MSc Thesis, University of Nairobi, Nairobi, Kenya. 2014.
- [34] Mengesha H, Birrie H, Gundersen G. The molluscicidal effect of the bark of “Bitza” tree: a local discovery towards the control of schistosomiasis in the Abay river valley of Western Ethiopia. *Ethiopian J Health Dev.* 1997;11:89-92.
- [35] Akanga JO. Screening of antidiarrhoea medicinal plants for in vitro antimicrobial activity against clinical and environmental enteropathogens. MSc Thesis, Jomo Kenyatta University of Agriculture and Technology, Nairobi, Kenya. 2008.
- [36] Omara T, Kiprof AK, Kosgei VJ. Intraspecific Variation of Phytochemicals, Antioxidant, and Antibacterial Activities of Different Solvent Extracts of *Albizia coriaria* Leaves from Some Agroecological Zones of Uganda. *Evid-Based Complement Altern Med.* 2021;2021:1-14.
- [37] Omara T, Kiprof AK, Kosgei VJ. Phytochemical screening, total polyphenolic content, in vitro antioxidant and antibacterial activities of *Albizia coriaria* Welw ex. Oliver flowers. *Lett Appl NanoBiosci.* 2022 (in press).
- [38] Nalubega R, Kabasa JD, Olila D, Kateregga J. Antibacterial activity and phytochemical screening of eleven plants used as poultry ethnomedicines in Southern Uganda. *Agric J.* 2011;6:303–309.
- [39] Byamukama R, Ganza B, Namukobe J, Heydenreich M, Kiremire BT. Bioactive compounds in the stem bark of *Albizia coriaria* (Welw. ex Oliver). *Int J Biol Chem Sci.* 2015;9(2):1013–1024.

- [40] Schultz F, Anywar G, Tang H, Chassagne F, Lyles JT, Garbe L, Quave CL. Targeting ESKAPE pathogens with anti-infective medicinal plants from the Greater Mpigi region in Uganda. *Sci Rep.* 2020;10(1):11935.
- [41] India, J., Efficacy of some medicinal plants used in various parts of Kenya in treating selected bacterial and fungal pathogens. MSc Thesis. Nairobi: Kenyatta University, School of Pure and Applied Sciences. 2015: p. 153.
- [42] Luvonga MW. Evaluation of antimicrobial activity of some plants used by traditional healers for treatment of microbial infections in Kakamega district: Kenya. MSc Thesis, Kenyatta University, Nairobi, Kenya. 2007.
- [43] Noté OP, Mitaine-Offer AC, Miyamoto T, Paululat T, Mirjolet JF, Duchamp O, Pegnyemb DE, Lacaille-Dubois MA. Cytotoxic acacic acid glycosides from the roots of *Albizia coriaria*. *J Nat Prod.* 2009;72(10):1725–1730.
- [44] Muthaura CN, Keriko JM, Mutai C, Yenesew A, Gathirwa JW, Irungu BN, Nyangacha R, Mungai GM, Derese S. Antiplasmodial potential of traditional phytotherapy of some remedies used in treatment of malaria in Meru-Tharaka Nithi County of Kenya. *J Ethnopharmacol.* 2015;175:315–323.
- [45] Schultz F, Osuji OF, Wack B, Anywar G, Garbe LA. Antiinflammatory medicinal plants from the Ugandan greater Mpigi region act as potent inhibitors in the COX-2/PGH2 pathway. *Plants.* 2021;10:351.
- [46] Obakiro SB, Kiprop A, K'Owino I, Andima M, Owor RO, Chacha R, Kigundu E. Phytochemical, Cytotoxicity, and Antimycobacterial Activity Evaluation of Extracts and Compounds from the Stem Bark of *Albizia coriaria* Welw ex. Oliver. *Evid-Based Complement Altern Med.* 2022. 2022:1-20.
- [47] Oloya B, Namukobe J, Ssengooba W, Afayoa M, Byamukama R. Phytochemical screening, antimycobacterial activity and acute toxicity of crude extracts of selected medicinal plant species used locally in the treatment of tuberculosis in Uganda. *Trop Med Health.* 2022;50:16.
- [48] Note OP, Chabert P, Pegnyem DE, Weniger B, Lacaille-Dubois M, Lobstein A. Structure elucidation of new acacic acid-type saponins from *Albizia coriaria*. *Mag Res Chem.* 2010; 48(10):829–836.
- [49] Omara T, Kiprop AK, Kosgei VJ. Isolation and characterization of compounds in ethanolic extract of *Albizia coriaria* (Welw ex. Oliver) leaves: a further evidence of its ethnomedicinal diversity. *Bull Natl Res Cent.* 2022;46:17.
- [50] Chepkorir R, Matasyoh JC, Wagara IN. Two withanolides from *Withania somnifera* (solanaceae) and activity of methanolic extracts against fungal and bacterial pathogens that affects food crops. *Afr J Food Sci.* 2018;12:115-125.
- [51] Yetayih MM, Ravichandran YD. Extraction and GC-MS Analysis of the Essential Oil from the Peel of *Solanum incanum* and its Antibacterial Activity Studies. *Asian J Chem.* 2020;32:2001-06.
- [52] Müller M, Volmer D. Interpretation of mass spectra. Institute for Bioanalytical Chemistry, Saarland University, 66123 Saarbrücken, Germany. 2017. [https://www.uni-saarland.de/fileadmin/upload/lehrstuhl/jauch/An04\\_Massenspektroskopie\\_Skript\\_Volmer.pdf](https://www.uni-saarland.de/fileadmin/upload/lehrstuhl/jauch/An04_Massenspektroskopie_Skript_Volmer.pdf).
- [53] Das AJ, Khawas P, Miyaji T, Deka SC. Phytochemical Constituents, Attenuated Total Reflectance Fourier Transform Infrared Analysis and Antimicrobial Activity of Four Plant Leaves Used for Preparing Rice Beer in Assam, India. *Int J Food Prop.* 2016;19:2087-2101.
- [54] Ferreira ICC, Aguiar EMG, Silva ATF, Santos LLD, Cardoso-Sousa L, Araújo TG, et al. Attenuated Total Reflection-Fourier Transform Infrared (ATR-FTIR) Spectroscopy Analysis of Saliva for Breast Cancer Diagnosis. *J Oncol.* 2020;2020:1-11.

- [55] Gómez HG. Determinación De Estructuras Orgánicas (Organic Spectroscopy): IR Spectroscopy. 2021. <http://www.upv.es/herme/files/4a-ir-spectroscopy.pdf>.
- [56] Netala VR, Kotakadi VS, Nagam V, Bobbu P, Ghosh SB, Tartte V. First report of biomimetic synthesis of silver nanoparticles using aqueous callus extract of *Centella asiatica* and their antimicrobial activity. *Appl Nanosci*. 2015;5:801–807.
- [57] Janakiraman N, Johnson M. Functional Groups of Tree Ferns (*Cyathea*) Using FT-IR: Chemotaxonomic Implications. *Romanian J Biophys*. 2015;25:131-141.
- [58] Song M, Hang TJ, Wang Y, Jiang L, Wu XL, Zhang Z, et al. Determination of oleanolic acid in human plasma and study of its pharmacokinetics in Chinese healthy male volunteers by HPLC tandem mass spectrometry. *J Pharmaceut Biomed Anal*. 2006;40:190–6.
- [59] Eunyong K, Noh K, Lee SJ, Shin B, Hwang JT, Lee SW, et al. Simultaneous determination of 3-O-acetyloleanolic acid and oleanolic acid in rat plasma using liquid chromatography coupled to tandem mass spectrometry. *J Pharmaceut Biomed Anal*. 2016;118:96–100.
- [60] Sut S, Poloniato G, Malagoli M, Dall'Acqua S. Fragmentation of the main triterpene acids of apple by LCAPCI-MSn. *J Mass Spect*. 2018;53:882–892.
- [61] Wahyuono S. Phytochemical Investigation of *Amsonia randiflora* Family Apocynaceae. MSc Thesis, University of Arizona, Tucson, Arizona. 1985.
- [62] Hu X, Shen Y, Yang S, Lei W, Luo C, Hou Y, Bai G. Metabolite identification of ursolic acid in mouse plasma and urine after oral administration by ultrahigh performance liquid chromatography/quadrupole time-of-flight mass spectrometry. *RSC Adv*. 2018; 8: 6532.
- [63] Chen Q, Zhang Y, Zhang W, Chen Z. Identification and quantification of oleanolic acid and ursolic acid in Chinese herbs by liquid chromatography-ion trap mass spectrometry. *Biomed Chromatogr*. 2011;25:1381–88.
- [64] Bugeja ML. Comparison of gas chromatography-mass spectrometry and liquid chromatography-mass spectrometry for discrimination of *Salvia divinorum* from related *Salvia* species using chemometric procedures. MSc Thesis, Michigan State University, East Lansing, Michigan. 2012.
- [65] Baek MY, Cho JG, Lee DY, Ann EM, Jeong TS, Baek NI. Isolation of triterpenoids from the stem bark of *Albizia julibrissin* and their inhibition activity on ACAT-1 and ACAT-2. *J Korean Soc Appl Biol Chem*. 2010;53:310–315
- [66] Jesus JA, Lago JH, Laurenti MD, Yamamoto ES, Passero LF. Antimicrobial activity of oleanolic and ursolic acids: an update. *Evid-based Complement Altern Med*. 2015;2015:1-14.
- [67] Joshi H, Saxena GK, Singh V, Arya E, Singh RP. Phytochemical investigation, isolation and characterization of betulin from bark of *Betula utilis*. *J Pharmacog Phytochem*. 2013;8192(1):266–285.
- [68] Namboozee J. Isolation and structural elucidation of antibacterial triterpenoids compounds from stem bark of *Psorospermum febrifugum* (Spach var. febrifugum). MSc Thesis, Makerere University, Kampala, Uganda. 2019.
- [69] Oh HM, Lee SW, Yun BR, Hwang BS, Kim SN, Park CS, et al. *Vigna angularis* inhibits IL-6-induced cellular signalling and ameliorates collagen-induced arthritis. *Rheumatol*. 2014;53:56–64.
- [70] Kuse M, Yanagi M, Tanaka E, Tani N, Nishikawa T. Identification of a Fluorescent Compound in the Cuticle of the Train Millipede *Parafontaria laminata armigera*. *Biosci Biotechnol Biochem*. 2010;74:2307-2309.
- [71] Allegri G, Costa Netto HJ, Ferreira Gomes, LN, Costa de Oliveira ML, Scalco FB, de Aquino Neto FR. Determination of six pterins in urine by LC-MS/MS. *Bioanal*. 2012;4:1739–46.

- [72] Ameera O, Ghaidaa J, Mohammed Y. Phytochemical screening of methanolic dried galls extract of *Quercus infectoria* using gas chromatography-mass spectrometry (GC-MS) and Fourier transform-infrared (FT-IR). *J Pharmacog Phytother.* 2016;8:49-59.
- [73] Hussein HJ, Hadi MY, Hameed IH. Study of chemical composition of *Foeniculum vulgare* using Fourier transform infrared spectrophotometer and gas chromatography - mass spectrometry. *J Pharmacog Phytother.* 2016;8:60-89.
- [74] Mohammed GJ, Hameed IH, Kamal SA. Study of secondary metabolites produced by *Aspergillus flavus* and evaluation of the antibacterial and antifungal activity. *African Journal of Biotechnology.* 2021. in press. [https://academicjournals.org/journal/AJB/article-in-press-abstract/study\\_of\\_secondary\\_metabolites\\_produced\\_by\\_a\\_spergillus\\_flavus\\_and\\_evaluation\\_of\\_the\\_antibacterial\\_and\\_antifungal\\_activity](https://academicjournals.org/journal/AJB/article-in-press-abstract/study_of_secondary_metabolites_produced_by_a_spergillus_flavus_and_evaluation_of_the_antibacterial_and_antifungal_activity).
- [75] Shenta AA, Al-Maliki ADM. Isolation and Identification of Three Alkaloids Compounds from *Albizia lebbek* L. Leaves and study of Their Antimicrobial Activity Against Pathogenic Bacteria of Urinary Tracts Inflammatory *in vitro*. *Univ Thi-Qar J Sci.* 2013;3:99-111.
- [76] Hashem H. Principles of mass spectrometric analysis. 2008. <http://pharmgrads.blogspot.com/2008/04/lecture3.html>.
- [77] Nicolescu TO. Interpretation of Mass Spectra. In: *Mass Spectrometry.* IntechOpen. 2017. pp. 23-78.
- [78] Dunnivant and Ginsbach. Fragmentation of Alcohols. 2008. [http://people.whitman.edu/~dunnivfm/C\\_MS\\_Ebook/CH6/6\\_10.html](http://people.whitman.edu/~dunnivfm/C_MS_Ebook/CH6/6_10.html)
- [79] Habib R, Mohyuddin A, Khan Z, Mahmood T. Analysis of Non-polar Chemical Profile of *Melia azedarach* L. *Sci Inquiry Review.* 2017;1:49-54.
- [80] Oloyede GK, Ali M, Lateef M. Enzyme inhibition, Antioxidant and Insecticidal activities of flavonoids and fixed oil from *Albizia zygia* (J. F. Macbr). *Life Sci J.* 2019;16:33-40.
- [81] Ayeleso TB, Matumba MG, Mukwevho E. Oleanolic Acid and Its Derivatives: Biological Activities and Therapeutic Potential in Chronic Diseases. *Molecules.* 2017;22:1915.
- [82] Gao D, Li Q, Li Y, Liu Z, Fan Y, Liu Z, et al. Antidiabetic and antioxidant effects of oleanolic acid from *Ligustrum lucidum* Ait in alloxan-induced diabetic rats. *Phytother Res.* 2009;23:1257-1262.
- [83] Horiuchi K, Shiota S, Hatano T, Yoshida T, Kuroda T, Tsuchiya T. Antimicrobial activity of oleanolic acid from *Salvia officinalis* and related compounds on vancomycin-resistant enterococci (VRE). *Biol Pharmaceut Bull.* 2007;30:1147-49.
- [84] Togashi N, Shiraishi A, Nishizaka M, Matsuoka K, Endo K, Hamashima H, Inoue Y. Antibacterial activity of long-chain fatty alcohols against *Staphylococcus aureus*. *Molecules.* 2007;12:139-148.