

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/363890613>

Factors Influencing the Time to Presentation, Diagnosis, and Treatment of Patients with Esophageal Cancer at Moi Teaching and Referral Hospital, Eldoret, Kenya

Article · September 2022

DOI: 10.5923/j.rct.20221001.03

CITATIONS

0

READS

14

3 authors, including:



Tabitha Chepkemai

Kabarak University

1 PUBLICATION 0 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



HPV uptake in Rongai and Nakuru West Sub- Counties, Nakuru Kenya. [View project](#)

Factors Influencing the Time to Presentation, Diagnosis, and Treatment of Patients with Esophageal Cancer at Moi Teaching and Referral Hospital, Eldoret, Kenya

Tabitha Chepkemoi^{1,*}, Diana Menya², Naftali Busakhala²

¹Kabarak University, Kabarak, Kenya

²Moi University, Eldoret, Kenya

Abstract Esophageal Cancer (EC) is a malignancy with high incidence and mortality globally. Late presentation and diagnosis lead to poor outcomes and high cost of treatment. The aim of this study was to describe factors influencing the time to presentation, diagnosis and treatment of patients with esophageal cancer at Moi Teaching and Referral Hospital (MTRH). This was a cross-sectional study conducted on patients with esophageal cancer seeking treatment at MTRH during the study period. Consecutive sampling of the patients in oncology clinic and surgical wards was done until a sample size of 104 was achieved. Logistic regression was used to determine the associations between factors and timelines for presentation, diagnosis and treatment using R-4.1.0-win software. The association was measured using Odds Ratios (ORs) with 95% CI. Monthly income of < 20,000 Kenya shillings (OR=0.1, CI, 0.002-0.52) was associated with late presentation to the health facility after the onset of symptoms. Being widowed (OR 0.05, CI, 0.003-0.39) was associated with late diagnosis while referral from another facility to MTRH was associated with early diagnosis (OR=3.74 CI, 1.08-14.7). Participants who had no medical insurance were more likely to be diagnosed and treated late (OR =0.3 CI, 0.079-0.82) and (OR= 0.27, CI, 0.079-0.82) respectively. Low income status was associated with late presentation to a health facility. Being widowed and lack of medical insurance were associated with late diagnosis and treatment while referral to MTRH was associated with early diagnosis.

Keywords Esophageal cancer, Cancer presentation, Cancer diagnosis, Cancer treatment, Cancer stage at diagnosis, Patient -mediated factors, Health system mediated factors, Late presentation, Early presentation

1. Introduction

Esophageal cancer ranks seventh with reference to incidence (604,100 new cases) and sixth cause of mortality (544,076 deaths) globally [1]. The latter shows that esophageal cancer is responsible for an estimated 1 in every 18 cancer deaths. Of all the esophageal cancer incidence and deaths globally, 80% occur in low and middle -income countries [2,3]. According to GLOBOCAN report [4], esophageal cancer is the fourth most common cancer in Kenya. It causes the highest number of cancer-related deaths in the country killing over 4,000 people annually [5]. The factors influencing the time to presentation, diagnosis, and treatment are summed in a complex interplay between the patient- mediated and health system mediated. Patient-mediated factors are described as factors influencing the time from onset of symptoms to the time of the first

presentation to the health facility include demographics, socioeconomic issues, health beliefs, and psychosocial [6]. Health system-related factors include all factors that are influenced by health care providers or the health system in terms of access to healthcare, diagnostic and treatment cost, and referrals [7,8]. Despite being a deadly disease, there have no studies conducted in Western Kenya on the factors related to time of patients' presentation to health facility for diagnosis and treatment [9]. The aim of the study therefore was to identify factors influencing the time to presentation, diagnosis, and treatment of patients with esophageal cancer at MTRH.

2. Materials and Methods

Study design and Study population: The study design was a cross-sectional which involved the interviewing of patients with esophageal cancer using a semi-structured questionnaire and the review of their medical records to retrieve information on clinical data. The study population

* Corresponding author:

tabithakirulex@gmail.com (Tabitha Chepkemoi)

Received: Jul. 13, 2022; Accepted: Aug. 5, 2022; Published: Aug. 23, 2022

Published online at <http://journal.sapub.org/rct>

included patients with esophageal cancer seeking medical and surgical treatment at Moi Teaching and Referral Hospital. The study population was patients who were histologically confirmed to be having esophageal cancer seeking medical and surgical treatment at Moi Teaching and Referral Hospital within the study period while those whose records were incomplete and those who have been visiting the clinic for more than 6 months (to avoid recall bias) [10] were excluded.

Sample size determination and Sampling techniques:

A sample size of 104 patients was estimated using the formula for the finite population (Daniel 1999) with the following assumptions: 50% prevalence of late esophageal cancer diagnosis, 5% margin of error, Critical value associated with type 1 error 1.96 (95% level of confidence). Patients were recruited using consecutive sampling until the targeted sample size was achieved.

Data collection techniques: At the oncology and cardiothoracic clinics, the researcher started from the clinician desk to check on the registers for the patients who were identified as having esophageal cancer. Their names were written down and then they were approached physically on the waiting bay to give consent for the researcher to peruse through their files and to be interviewed. For those who met eligibility criteria, the researcher retrieved the information on clinical data from their files then proceeded to interview them in a secluded area within the hospital. For those who were admitted to the surgical wards cardiothoracic section, the researcher identified the patients with esophageal cancer with the assistance of the nurse on duty. They too were approached to give consent for their file to be perused and to be interviewed if they met eligibility criteria.

Data management and analysis: Data was reviewed after collection to check for missing data and unclear entries. Data captured using the structured interviewer-administered questionnaires were entered into an electronic database, MS. Excel. The database was encrypted with a password to ensure confidentiality. Data analysis was done using R-4.1.0-win software. Descriptive statistics was used for demographic characteristics where frequency tables and percentages were calculated. Logistic regression was also used to determine the associations between explanatory factors determining the stage at diagnosis and timelines of esophageal cancer presentation, diagnosis and treatment. The strength of the association was measured using ORs with 95% CI.

Ethics: The study sought ethical approval from the Institutional Research and Ethics Committee (IREC) of Moi University, IREC/2019/288. Informed written consent was obtained from all the eligible participants and those who consented were interviewed. Permission was also sought from the hospital administration before the commencement of the study. Informed written consent was obtained from all eligible patients, Information about the research was given verbally to each patient and those that gave a written consent were interviewed.

3. Results

Table 1. Socio-demographic characteristics

Variable	Categories	Frequency (%)
Age	Mean Age in years (SD)	57.2 (12.5)
Gender	Male	67 (65.7)
	Female	35 (34.3)
Marital Status	Married	87(85.3)
	Separated	2(2.0)
	Single	2(2.0)
	Widowed	11(10.7)
Regions of Residence	North Rift	41(40.2)
	Nyanza	30 (29.4)
	South Rift	7 (6.9)
	Western	24 (23.5)
Type of area lived	Rural	83(81.4)
	Suburban	13(12.7)
	Urban	6(5.9)
Level of Education	College/University	12 (11.8)
	Secondary	50(49.0)
	Primary	32(31.4)
	No formal education	8(7.8)
Religion	African traditional Religion	2(2.0)
	Catholic	38(37.3)
	protestants	62(60.8)
Occupation	Business	25(24.5)
	Farming	65(63.7)
	Home maker	2 (2.0)
	Office employment	7(6.9)
	Others	3(2.9)
Monthly Income	None	3(2.5)
	Less than 20,000	73(72.5)
	Above 20,000	26 (25.0)
Health Insurance	No Insurance	40 (39.2)
	NHIF	59(57.8)
	NHIF+ Private Health Insurance	3(3.0)

The mean age of the study participants was 57.2 years (SD, 12.5 years) at the time of study. A majority of participants 65.4% were male, 85.3% married, 81.4% living in rural setting and 40.2% were residents of north rift region. On education, 7.8% were illiterate, with over 60.0% having secondary and above as their highest level of education. On religion, a majority of participants were Protestants 60.8%. Majority of the participants 63.7%, were farmers with 72.5% earning less than 20,000 shillings per month. Most participants 60.8% had medical insurance Table 1. Participants with monthly income less than 20,000 Kenya shillings 0.1 (0.002-0.52) were more likely to present late to health facility after the onset of symptoms Table 2. Being widowed AOR, 0.05 CI (0.003-0.39) and having no medical insurance AOR 0.3 CI (0.07-0.76), were more likely to make participants delay in esophageal cancer diagnosis. Participants who were referred from another facility to MTRH were more likely to be diagnosed early as compared

to those not referred AOR 3.74, CI (1.08- 14.70) Table 3. more likely for a patient to delay for treatment after
Having no medical insurance AOR 0.27, CI (0.08-0.82), was histological confirmation of esophageal cancer Table 4.

Table 2. Factors influencing time to first presentation of patients with esophageal cancer

Independent Variables	Categories	COR(95% CI)	AOR (95% CI)
Age		1.0(0.97- 1.03)	0.9 (0.94-1.04)
Gender	Male	1.2(0.51-2.71)	1.4 (0.45-4.54)
	Female	ref	ref
Marital Status	Married	Ref	ref
	Single	1	1
	Separated	0.6 (0.02-16)	1.4 (0.04-45.95)
	widowed	0.5 (0.14-1.91)	0.8 (0.11-5.86)
Level of Education	No formal education	3.3(0.54-24.43)	15 (0.69-804)
	Primary	3.8 (0.98-17.10)	4.8 (0.63-45.7)
	Secondary	3.3(0.9-13.63)	3.7 (0.58-27.91)
	Tertiary	ref	ref
Monthly income	No Income	Ref	ref
	Less than 20,000	0.09(0.004-0.76)	0.1 (0.002-0.52)
	Above 20,000	0.4(0.11-1.30)	0.9(0.18-4.05)
Medical Insurance	No Insurance	1.6(0.93- 2.67)	0.9(0.27-2.69)
	NHIF	1	1
	Private Insurance	1	1
	NHIF+ Private Insurance	ref	ref
Use of traditional medicine	Yes	0.8 (0.37-1.85)	0.9 (0.34- 2.85)
	No	Ref	ref
Ethnicity	Kikuyu	2.8 (0.32-60.40)	7.8 (0.3-798)
	Kisii	1.6 (0.41-7.30)	3.6 (0.63-24.83)
	Luhya	1.5 (0.53- 4.36)	2.8 (0.81- 10.87)
	Luo	1.7 (0.60-5.45)	3.3 (0.78- 15.22)
	Pokot	1.4 (0.21- 11.78)	1.3 (0.03-51.11)
	Kalenjin	Ref	ref
Religion	Catholic	1	1
	Protestant	1	1
	African Tradition	Ref	ref
History of esophageal cancer	Yes	1.5 (0.57-0.39)	2.3 (0.68-8.72)
	No	Ref	Ref

Table 3. Factors influencing time to diagnosis of patients with esophageal cancer

Independent Variables	Categories	COR (95% CI)	AOR (95% CI)
Age		1(0.97-1.03)	1.02 (0.96-1.07)
Gender	Male	0.8 (0.36-1.85)	0.55 (0.15-1.83)
	Female	Ref	Ref
Marital Status	Married	Ref	Ref
	Single	1	1
	Separated	0.9(0.03-21.98)	0.39 (0.008-16.42)
	Widowed	0.2(0.03-0.79)	0.05 (0.003-0.39)
Level of Education	No formal education	2(0.03-21.98)	2.80 (0.15-64.34)
	Primary	1.2(0.23-8.89)	1.56 (0.19- 12.39)
	Secondary	2(0.50-6.75)	2.29 (0.34-15.03)
	Tertiary	ref	ref
Monthly income	None	ref	ref
	Less than 20,000	1.4(0.09-1.20)	0.9 (0.24-4.15)
	Above 20,000	0.8(0.30-2.20)	1.5 (0.14-21.31)
Medical Insurance	No Insurance	0.5(0.22-1.13)	0.3 (0.07-0.76)
	NHIF	1	1
	Private Insurance	1	1

Independent Variables	Categories	COR (95% CI)	AOR (95% CI)
Use of traditional medicine	Yes	1.2(0.53-2.55)	1.02 (0.36- 2.85)
	No	Ref	ref
Referral from another health facility	Yes	2.3(0.84-6.56)	3.74 (1.08- 14.70)
	No	ref	Ref
Ethnicity	Kikuyu	1.1(0.12-9.73)	1.40 (0.07-39.80)
	Kisii	0.6(0.14-2.41)	0.23 (0.036-1.27)
	Luo	1.4(0.52-4.14)	1.05 (0.26-4.13)
	Luhya	1.4(0.47-4.10)	2.35 (0.30-6.47)
	Pokot	0.7(0.08-4.82)	0.73 (0.015-18.91)
	Kalenjin	ref	ref
Religion	Catholic	1.4(0.04-25.29), 0.83	0.41 (0.003-47.80)
	Protestant	0.9(0.05-36.55), 0.93	0.29 (0.002-27.98)
	African Tradition	ref	Ref

Table 4. Factors influencing time to treatment of patients with esophageal cancer

Independent Variables	Categories	COR (95% CI)	AOR (95% CI)
Age	Age	0.9 (0.96-1.02)	0.97 (0.92-1.01)
Marital Status	Married	ref	ref
	Single	1.7(0.06-43.67)	1.2 (0.03-46.67)
	Separated	1.7(0.06-43.67)	1.8 (0.05-55.37)
	widowed	0.4(0.05-1.57)	0.2 (0.02-1.67)
Level of Education	No formal education	2(0.32-13.42)	16 (0.93-444.19)
	Primary	0.5(0.11-2.27)	0.9 (0.12-9.08)
	Secondary	1.6(0.43-6.51)	2.6 (0.39-22.51)
	Tertiary	ref	ref
Monthly income	None	ref	ref
	Less than 20,000	0.5(0.13-2)	2.58 (0.76-9.51)
	Above20,000	1.7(0.63-4.47)	0.67 (0.13-3.01)
Medical Insurance	No Insurance	0.6 (0.23-1.35)	0.27 (0.08-0.82)
	NHIF	1	1
	Private Insurance	1	1
	NHIF+ Private insurance	Ref	Ref
Use of traditional medicine	Yes	1.2(0.53-2.76)	1.36 (0.49-3.83)
	No	ref	ref
Referral from another health facility	Yes	1.7(0.59-5.68)	1.55 (0.44-6.15)
	No	ref	ref
Distance to MTRH	Less than 1Hr	0.9(0.31-2.81)	1
	1-2 Hrs.	0.6(0.12-2.40)	1.87 (0.29- 11.90)
	3-4 Hrs.	0.8(0.25-2.5)	1.57 (0.25-10.00)
	Above 4 Hrs.	ref	ref

4. Discussion

The study revealed that esophageal cancer is common among the older population; the results were consistent with other published reports [10,11]. The male: female ratio among the patients with esophageal cancers was 1.9: 1, a slightly higher ratio than 1.5:1 found by [12]. The higher male to female ratio may be attributable to higher risk among men such as drinking alcohol and smoking [13]. The distribution of patients with esophageal cancer was high in North rift region which can be explained by the close proximity and accessibility of MTRH. Nyanza region, however, recorded a significant figure supporting previous study that found high incidence rates of esophageal cancer

near Lake Victoria [14]. A relatively high number of the participants had low level of education, with a majority in low socio-economic status. Despite the efforts by Kenyan government to increase the NHIF enrollment, the study revealed most participants still had no medical insurance. However, there has been a significant improvement (57.8%) in NHIF coverage overall as compared to 2015-2016 NHIF coverage of about 19% [15].

Factors influencing time to first presentation of patients with esophageal cancer

The study revealed that monthly income was associated with time to presentation of patients with esophageal cancer. Those patients who earned an income of less than 20,000

shillings were 5 times more likely to present late as compared to their counterparts who earned more monthly income. This can be explained by structural inequalities in Kenya's health care, disproportionality influence accessibility where people with high income status seek health care more conveniently [16,17]. Transport cost and other related expenses to medical attention are some of the hindrances to early presentation [14]. Income status is a precursor for social economic status (SES), so people with low SES have poor financial support and less ability to afford medical care prolonging delay time. The results agreed with a study done in China that found an association between social economic status and health care delay in esophageal cancer [18,19,20].

Factors influencing time to diagnosis of patients with esophageal cancer

In our study, widowed patients were 6 times more likely to be diagnosed late in reference to their married counterparts, (OR 0.062, CI, 0.004-0.43). These results were consistent with other studies that have reported an association between being widowed and late esophageal cancer diagnosis as well as higher mortality from esophageal cancer [21,22]. This association may be attributable to social challenges resulting from loss of emotional support, less income, and fewer social relationships as a result of partner loss. The study found a > 2 folds increased risk of being diagnosed late with esophageal cancer among the patients with no medical insurance as compared to those participants who had either NHIF or private medical insurance (OR =0.228 CI, 0.057-0.80). Medical insurance cost in Kenya is high leaving the poor families to pay for diagnostic services out of pockets causing delay in healthcare finance mobilization and negating their health seeking behaviors for esophageal cancer for diagnosis. The government in the recent past has put efforts to improve the NHIF coverage for patients with terminal illness, [14] but more effort is still needed because the percentage of cancer patients who are uninsured is still high. The results were in agreement with a study done by [23], on cost and affordability of NCDs screening, diagnosis and treatment in Kenya who found out that the payments required for cancer diagnostic tests for patients without medical insurance were beyond their reach. The patients who were referred from other facilities to MTRH were 3.74 more likely to be diagnosed early as compared to those patients who were not referred (OR=3.74 CI, 1.08- 2.34). Delay in referral is often related to misdiagnosis of common symptoms, for instance treating heartburns, pneumonia, GERD, and esophagitis among others [24]. Strengthening the referral system has proven to reduce the delays in diagnosis thus improving the overall outcome of the disease. The use of telephone contact or digital referral system is one way to reduce the navigation pathway [25,26].

Factors influencing time to treatment of patients with esophageal cancer

Having no medical insurance, was statistically significant factor to delay in seeking treatment for esophageal cancer after histological confirmation (OR= 0.27, CI, 0.079-0.823).

The cost of cancer treatment in Kenya is very expensive, impeding many patients from seeking early treatment [27,28]. The government of Kenya has made positive strides on improving the NHIF care package on cancer treatment including ten chemotherapy sessions, oral and injectable anti-cancers drugs, inpatient and outpatient oncology services, twenty sessions for radiotherapy, and about two sessions for Brachytherapy for advanced cancer, per year [14,15]. It is therefore, evident that having no medical insurance is a huge barrier to early cancer treatment because the patients miss out on the NHIF package that would help in treatment to the cancer. In addition, a medical insurance is an indicator of social economic status aggravating financial constraints in seeking cancer treatment services [21].

5. Conclusions

Income status influenced the time to presentation of patients with esophageal cancer in MTRH, with low income associated with late presentation to first medical attention. Widowed individuals and lack of medical insurance were associated with late diagnosis while referral to MTRH was associated with early diagnosis of esophageal cancer. Lack of medical insurance was also associated with late treatment of esophageal cancer. These factors' influence on esophageal cancer burden and overall implications on patient health point to the need for tailored interventions to modify and support positive healthcare seeking behaviors and overall outcome of the disease in the population.

ACKNOWLEDGEMENTS

We thank God for giving us the strength to carry out the study; we also convey our gratitude to the study participants, and the School of Public Health Moi University for academic guidance.

Disclosure

Competing interests

The authors declare no competing interest.

REFERENCES

- [1] Sung, H., Ferlay, J., Siegel, R. L., Laversanne, M., Soerjomataram, I., Jemal, A., & Bray, F. (2021). Global Cancer Statistics . GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA: A Cancer Journal for Clinicians*. 2020; 71(3), 209–249. <https://doi.org/10.3322/caac.21660>.
- [2] International, T., Epidemiology, C., Cheng, M. L., Zhang, L., Borok, M., Chokunonga, E., Dzamamala, C., Korir, A., Wabinga, H. R., Hiatt, R. A., Parkin, D. M., & Loon, K. Van. *The incidence of oesophageal cancer in Eastern Africa:*

- Identification of a new geographic hot spot?* §. 2015; 39, 7821.
- [3] Mathew, C. G., Middleton, D., Murphy, G., & Dawsey, S. M. (2022). *The African Esophageal Cancer Consortium: A Call to Action*. <https://doi.org/10.1200/JGO.17.00163>.
- [4] Globocan. (2020). *Kenya Source*. 799, 2020–2021; <https://gco.iarc.fr/today/data/factsheets/populations/404-kenya-fact-sheets.pdf>.
- [5] Car, L. T., Urch, C., Majeed, A., Khatib, M. El, Aylin, P., Atun, R., Car, J., Vincent, C., & Car, L. T. *Correspondence to: Preventing delayed diagnosis of cancer: clinicians' views on main problems and solutions*. 2016; 6(2). <https://doi.org/10.7189/jogh.06.020901>.
- [6] Akuoko, C. P., Armah, E., Sarpong, T., Quansah, D. Y., Amankwaa, I., & Boateng, D. (2017a). *Barriers to early presentation and diagnosis of breast cancer among African women living in sub-Saharan Africa*. 1–18. <https://doi.org/10.1371/journal.pone.0171024>.
- [7] Espina, C., & McKenzie, F. *Annals of Epidemiology Delayed presentation and diagnosis of breast cancer in African women: a systematic review*. *Annals of Epidemiology*. 2017; 27(10), 659–671. e7. <https://doi.org/10.1016/j.annepidem.2017.09.007>.
- [8] Njuguna, F., Martijn, H., Langat, S., Musimbi, J., & Muliro, H. *FACTORS INFLUENCING TIME TO DIAGNOSIS AND TREATMENT AMONG PEDIATRIC ONCOLOGY PATIENTS IN KENYA*. n.d; .33(2016), 186–199.
- [9] Patel, K., Wakhisi, J., Mining, S., Mwangi, A., & Patel, R. *Esophageal Cancer, the Topmost Cancer at MTRH in the Rift Valley, Kenya, and Its Potential Risk Factors*. 2013.
- [10] Majur, M., & Mabior, C. *Time To Presentation And Diagnosis In Patients With Esophageal Cancer At Kenyatta National Hospital*. 2018.
- [11] Wakhisi, J., Patel, K., Buziba, N., & Rotich, J. *Esophageal cancer in north rift valley of western Kenya*. 2005; 157–163.
- [12] Wakhisi, J., Patel, K., Buziba, N., & Rotich, J. (2005). *Esophageal cancer in north rift valley of western Kenya*. *African Health Sciences*, 5(2), 157–163.
- [13] Middleton, D. R. S., Bouaoun, L., Hanisch, R., Bray, F., Dzamalala, C., Chasimpha, S., Menya, D., Mbalawa, C. G., N'Da, G., Woldegeorgis, M. A., Njie, R., Koulibaly, M., Buziba, N., Ferro, J., Nouhou, H., Ogunbiyi, F., Wabinga, H. R., Chokunonga, E., Borok, M. Z., ... McCormack, V. A. *Esophageal cancer male to female incidence ratios in Africa: A systematic review and meta-analysis of geographic, time and age trends*. *Cancer Epidemiology*. 2018; 119–128. <https://doi.org/10.1016/j.canep.2018.01.020>.
- [14] Barasa, E., Rogo, K., Mwaura, N., & Chuma, J. *Kenya national hospital insurance fund reforms: Implications and lessons for universal health coverage*. *Health Systems and Reform*. 2018; 4(4), 346–361. <https://doi.org/10.1080/23288604.2018.1513267>.
- [15] Greene, S. B., Othieno-abinya, N. A., & Bennett, A. V. *Improving Access to Cancer Testing and Treatment in Kenya*. 2018; 7. <https://doi.org/10.1200/JGO.2017.010124>.
- [16] Wheeler, S., Bennett, A. V., Makau-Barasa, L. K., Othieno-Abinya, N. A., Greene, S. B., & Skinner, A. *Improving Access to Cancer Testing and Treatment in Kenya*. *Journal of Global Oncology*. 2017; 7(4), 1–8. <https://doi.org/10.1200/jgo.2017.010124>.
- [17] Wambalaba, F. W., Son, B., Wambalaba, A. E., Nyong, D., & Nyong, A. (2019). *Prevalence and Capacity of Cancer Diagnostics and Treatment: A Demand and Supply Survey of Health-Care Facilities in Kenya*. 26, 1–12. <https://doi.org/10.1177/1073274819886930>.
- [18] Chen, H., Chen, I., Chen, Y., Chen, C., & Chuang, C. (2022). *The Influence of Socioeconomic Status on Esophageal Cancer in Taiwan: A Population-Based Study*.
- [19] Kou, K., Baade, P. D., Guo, X., Gattton, M., Cramb, S., Lu, Z., Fu, Z., Chu, J., Xu, A., & Sun, J. (2019). *Area socioeconomic status is independently associated with esophageal cancer mortality in*. *Scientific Reports, November 2018*, 1–8. <https://doi.org/10.1038/s41598-019-42774-x>.
- [20] Wang, N., Cao, F., Liu, F., Jia, Y., Wang, J., Bao, C., Wang, X., Song, Q., Tan, B., & Cheng, Y. *The effect of socioeconomic status on health - care delay and treatment of esophageal cancer*. *Journal of Translational Medicine*. 2015; 1–5. <https://doi.org/10.1186/s12967-015-0579-9>.
- [21] Islami, F., Kamangar, F., Nasrollahzadeh, D., Aghcheli, K., Sotoudeh, M., Abedi-Ardekani, B., Merat, S., Nasseri-Moghadam, S., Semnani, S., Sepehr, A., Wakefield, J., Møller, H., Abnet, C. C., Dawsey, S. M., Boffetta, P., & Malekzadeh, R. *Socio-economic status and oesophageal cancer: Results from a population-based case-control study in a high-risk area*. *International Journal of Epidemiology*. 2009; 38(4), 978–988. <https://doi.org/10.1093/ije/dyp195>.
- [22] Jansson, C., Johansson, A. L. V., & Nyre, O. *Socioeconomic Factors and Risk of Esophageal Adenocarcinoma: A Nationwide Swedish Case-Control Study*. 2005 July 14; 1754–1762.
- [23] Subramanian, S., Gakunga, R., Kibachio, J., Gathecha, G., Edwards, P., Ogola, E., Yonga, G., Busakhala, N., Munyoro, E., Chakaya, J., Ngugi, N., Mwangi, N., Rege, D. Von, Wangari, L. M., Wata, D., Makori, R., Mwangi, J., & Mwanda, W. *Cost and affordability of non-communicable disease screening, diagnosis and treatment in Kenya: Patient payments in the private and public sectors*. 2018; *PLoS ONE*, 13(1), 1–16. <https://doi.org/10.1371/journal.pone.0190113>.
- [24] Grotenhuis, B. A., van Hagen, P., Wijnhoven, B. P. L., Spaander, M. C. W., Tilanus, H. W., & van Lanschot, J. J. B. *Delay in diagnostic workup and treatment of esophageal cancer*. *Journal of Gastrointestinal Surgery*. 2010; 14(3), 476–483. <https://doi.org/10.1007/s11605-009-1109-y>.
- [25] van Erp, N. F., Helsper, C. W., Slottje, P., Brandenbarg, D., Büchner, F. L., van Asselt, K. M., Muris, J. W. M., Kortekaas, M. F., Peeters, P. H. M., & de Wit, N. J. *Time to diagnosis of symptomatic gastric and oesophageal cancer in the Netherlands: Where is the room for improvement?* *United European Gastroenterology Journal*. 2020; 8(5), 607–620. <https://doi.org/10.1177/2050640620917804>.
- [26] Gyenwali, D., Pariyar, J., & Onta, S. R. (2013). *Factors Associated with Late Diagnosis of Cervical Cancer in Nepal*. 14, 4373–4377.
- [27] Makau-Barasa, L. K., Greene, S., Othieno-Abinya, N. A., Wheeler, S. B., Skinner, A., & Bennett, A. V. *A review of Kenya's cancer policies to improve access to cancer testing*

and treatment in the country. *Health Research Policy and Systems*. 2020; 18(1), 1–10.
<https://doi.org/10.1186/s12961-019-0506-2>.

[28] Atieno, O. M., Opanga, S., Martin, A., Kurdi, A., Michelle,

O., Opanga, S., Martin, A., & Kurdi, A. (2018). Pilot study assessing the direct medical cost of treating patients with cancer in Kenya; findings and implications for the future. *Journal of Medical Economics*, 21(9), 878–887.
<https://doi.org/10.1080/13696998.2018.1484372>.

Copyright © 2022 The Author(s). Published by Scientific & Academic Publishing

This work is licensed under the Creative Commons Attribution International License (CC BY). <http://creativecommons.org/licenses/by/4.0/>