

## Ebola: The Bundibugyo virus is back

- Risk factors in glaucoma
- OV16 antibody seropositivity and onchocerciasis transmission
- Urinary schistosomiasis in the River Niger Basin areas
- Communicable and infectious diseases research agenda
- Disaster preparedness among public health facilities
- Health information system infrastructure and data quality
- Maternal knowledge and caregiving practices in stunting
- And more . . .

# SSMJ

SOUTH SUDAN MEDICAL JOURNAL

ISSN 2309 - 4605 eISSN 2309-4613 Volume 19. No. 2. May 2026

A Publication of the Health and Social Sciences Research Institute of South Sudan

Juba, South Sudan

Email: [southsudanmedicaljournal@gmail.com](mailto:southsudanmedicaljournal@gmail.com) Website: [www.southsudanmedicaljournal.com](http://www.southsudanmedicaljournal.com)

---

## EDITOR-IN-CHIEF

Dr Edward Eremugo Kenyi  
South Sudan Medical Journal  
Juba, South Sudan

## EDITORS

Prof John Adwok  
Dr Charles Bakhiet  
Dr Richard Bogle  
Dr Nyakomi Adwok  
Dr Justin Bruno Tongun  
Dr Boniface A.E Lumori  
Dr James Ayrton  
Dr David Tibbutt

## WEB TEAM

Dr Edward Eremugo Kenyi  
Dr Rachel Ayrton

## DESIGN AND LAYOUT

Dr Edward Eremugo Kenyi

## ASSOCIATE EDITORS

Dr Eluzai Abe Hakim  
Newport, UK

Dr Wani Gindala Mena  
Juba, South Sudan

Dr Koma Akim  
Juba, South Sudan

## MANAGING EDITOR


Ann Burgess

## EDITORIAL TEAM

Dr Grace Juan Soma  
Dr James Frater  
Dr Onyango J.O. Okech  
Dr Anthony Garang  
Dr Anil Cherian  
Dr Indranil Saha  
Nancy MacKeith  
James Beard

---

## Index and Copyright Information



Scopus

DOAJ  
DIRECTORY OF  
OPEN ACCESS  
JOURNALS

The *South Sudan Medical Journal* is a quarterly publication intended for Healthcare Professionals, both those working in the South Sudan and those in other parts of the world seeking information on health in South Sudan. The Journal is published in mid-February, May, August and November.

It is an Open Access Journal licensed under a [Creative Commons Attribution - Noncommercial Works License \(CC BY-NC 4.0\)](https://creativecommons.org/licenses/by-nc/4.0/).

## Editorial

- Ebola: The Bundibugyo virus is back *Edward Eremugo Kenyi* ..... 74

## Research Articles

- Risk factors in glaucoma: Insights from a prospective case–control study, Belagavi, India *Asawari Kabure, Bhushan Kulkarni, Shilpa Kodkany, Murigendra Hiremath* ..... 75
- OV16 antibody seropositivity reveals persistent onchocerciasis transmission and cross-border risk in Magwi County, South Sudan: a population-based survey *WApary et al* ..... 81
- Prevalence of urinary schistosomiasis in communities along the River Niger Basin areas, North-Central Nigeria *Nyamngee et al* ..... 87
- Developing a communicable and infectious diseases research agenda in South Sudan: A multi-stakeholder prioritisation approach *Jacob et al* ..... 93
- Determinants of disaster preparedness among public health facilities in Juba City, South Sudan *Dumba Samuel Kenyi, Muthoni Mwangi Eunice, Musa Olouch* ..... 99
- Evaluating the role of the health information system infrastructure in enhancing data quality in Central Equatoria State, South Sudan *Matur T. Tieng, Karani Magutah, Joseph Muchiri* ..... 107
- Impact of developmental stimulation education on maternal knowledge and caregiving practices in stunting: a longitudinal study *Ning Iswati et al* ..... 112
- Holding the line: lived experiences of mental health, moral injury, and organizational support among public-health professionals in the Rohingya Camp, Bangladesh — a phenomenological study *Md. Ruhul Amin* ..... 119

## Case Reports

- Endoscopic endonasal repair of post-traumatic sphenoid sinus cerebrospinal fluid leak with encephalocele *Justin Rubena, Khalid Coco, Francis Swaka, Othwonh Twong* ..... 126
- A rare pathological feature of portal vein thrombosis complicated by portal hypertension and haematemesis *Illa et al* ..... 130

## Review Article

- Interpersonal and community-level factors influencing intimate partner violence in South Sudan: a literature review *Nyinypiu Tong Chol Adong* ..... 135

## Short Communications

- Letters to the Editor *J. Clarke McIntosh and Eluzai A. Hakim* ..... 141
- News from the Gordon Memorial College Trust Fund *Eluzai A. Hakim* ..... 143

## BACKCOVER: CDC Poster: Flu or Ebola?

**FRONT COVER IMAGE:** Health workers using personal protective equipment (PPE) at a training during the Ebola outbreak in Guinea, 2014 (CDC/Lindsey Horton - Public Health Image Library ID #18706)

# Ebola: The Bundibugyo virus is back

On 17 May 2026, the World Health Organization (WHO) declared the new outbreak of Ebola Virus Disease (EVD) in the Democratic Republic of the Congo (DRC), a public health emergency of international concern.<sup>[1]</sup> As of 22 May 2026, there were 750 suspected cases, and 177 deaths had been reported in the DRC; two cases (with one death) had been reported in Kampala, Uganda. New cases and deaths are expected to rise as the virus spreads. This is the 17th EVD outbreak in the DRC since the first in 1976.<sup>[2]</sup>

This outbreak is caused by a rare strain of the Ebola virus, the Bundibugyo type, which had previously caused only two outbreaks: the 2007 outbreak in Bundibugyo province in western Uganda<sup>[3]</sup> and the 2012 outbreak in Isiro, Pawa, Dungu, and Province Orientale in the DRC.<sup>[4]</sup> There are two other types of the virus that caused most of the last outbreaks: the Ebola (Zaire) and Sudan viruses.

First transmitted from wild animals to humans, the virus can spread from person to person through close contact with an infected person's body fluids, leading to symptoms, appearing after two to 21 days, including fever, fatigue, myalgia, vomiting, diarrhoea, headache, and internal and external bleeding with high mortality up to 50%. Although two vaccine types have been approved for the Zaire strain of the virus, there is no vaccine for the Bundibugyo virus. We need one urgently.

Although the current outbreak is not considered a pandemic, the threat of serious regional spread of this virus is very high, thus necessitating the declaration of an international emergency. DRC and Uganda have activated their national disaster/emergency management mechanisms and establish emergency operation centers. The WHO and Africa CDC have sent teams into the countries to provide support. South Sudan has raised its emergency and surveillance levels amid ongoing monitoring of the situation in the DRC.

But why does the Ebola virus keep coming back? Is it because of the endemicity in fruit bats, which act as the reservoirs, and eating these bats allow the virus to cross into humans, makes it difficult to control? The DRC regions involved are rife with conflict, poverty, isolation, and inadequate health infrastructure. How can we conduct community prevention and disease surveillance more effectively in such conditions?

As the health authorities respond to the emergency, the outbreak reminds us to continually evaluate our work and develop new out-of-the-box, multipronged approaches to tackle this disease. Perhaps then we can break the cycle.

### Reference:

1. World Health Organization. Epidemic of Ebola Disease caused by Bundibugyo virus in the Democratic Republic of the Congo and Uganda determined a public health emergency of international concern, WHO: Geneva. 17 May 2026.
2. World Health Organization. Ebola virus disease - Democratic Republic of the Congo. Disease Outbreak News. WHO. 1 December 2025.
3. Wamala et al. "Ebola Hemorrhagic Fever Associated with Novel Virus Strain, Uganda, 2007–2008". *Emerging Infectious Diseases*. July 2010; 16 (7): 1087–1092.
4. Ebola outbreak in Democratic Republic of Congo (Update) – WHO Regional Office for Africa: WHO. 20 August 2012.

**Edward Eremugo Kenyi** 

**Editor-in-Chief**

**South Sudan medical Journal**

**Email:**

[southsudanmedicaljournal@gmail.com](mailto:southsudanmedicaljournal@gmail.com)

**Citation:** Kenyi. Ebola: The Bundibugyo virus is back. *South Sudan Medical Journal*. 2026;19(2):74 © 2026 The Author (s) **License:** This is an open access article under [CC BY-NC](https://creativecommons.org/licenses/by-nc/4.0/) DOI: <https://dx.doi.org/10.4314/ssmj.v19i2.1>

# Risk factors in glaucoma: Insights from a prospective case–control study, Belagavi, India

Asawari Kabure<sup>1</sup>, Bhushan Kulkarni<sup>2</sup>, Shilpa Kodkany<sup>3</sup>, Murigendra Hiremath<sup>1</sup> 

## Author Affiliation:

1. Department of Microbiology & Biotechnology, Karnatak University, Dharwad, Karnataka, India
2. Dr. Prabhakar Kore Basic Science Research Center & JNMC Women's & Children's Research Unit, Belagavi, Karnataka, India
3. Dr. Kodkany's Superspeciality Eye Care Centre, Belagavi, Karnataka, India

## Correspondence:

Murigendra Hiremath  
[murigendra@gmail.com](mailto:murigendra@gmail.com)

**Submitted:** January 2026

**Accepted:** March 2026

**Published:** May 2026

**Citation:** Kabure et al. Risk factors in glaucoma: Insights from a prospective case–control study, Belagavi, India. South Sudan Medical Journal, 2026;19(2):75-80 © 2026 The Author (s) **License:** This is an open access article under [CC BY-NC](https://creativecommons.org/licenses/by-nc/4.0/) DOI: <https://dx.doi.org/10.4314/ssmj.v19i2.2>

## ABSTRACT

**Introduction:** Glaucoma is an optic neuropathy and degeneration of retinal ganglion cells, causing irreversible visual damage and blindness. While raised intraocular pressure (IOP) is the most important established risk factor, the exact pathogenesis and other risk factors for glaucoma are not fully understood, particularly in developing countries. Early identification and management of risk factors are important for reducing incidence and burden. This study assessed the risk factors for glaucoma at Dr. Kodkany's Eye Center in Belagavi, India.

**Method:** This case-control study examined 24 cases with confirmed diagnosis of glaucoma and 20 age- and sex-matched controls. A pre-designed questionnaire was used to collect sociodemographic data along with detailed eye examination findings. All glaucoma participants underwent thorough ophthalmic examinations, including visual acuity, intraocular pressure (IOP), central corneal thickness, anterior segment abnormalities, anterior chamber angle, fundus photography, and visual field testing.

**Results:** Of the 24 cases, 14 (58.3%) were diagnosed with open-angle glaucoma, with bilateral involvement noted in 13(54.2%) cases. IOP was considerably higher in glaucomatous patients compared to non-glaucomatous (right eye:  $27.86 \pm 8.97$  vs  $15.85 \pm 2.84$  mmHg [Mean difference: 12.01 (95% CI: 8.24–15.78)]; left eye:  $28.27 \pm 8.75$  vs  $16.05 \pm 3.02$  mmHg; [Mean difference: 12.22 (95% CI: 8.49–15.95)]  $p < 0.001$  for both). None of the studied factors — age (OR=0.990, 95% CI: 0.937-1.045,  $p=0.705$ ), female gender (OR=1.667, 95% CI: 0.500-5.559,  $p=0.416$ ), or hypertension (OR=2.538, 95% CI: 0.697-9.240,  $p=0.160$ )—predicted glaucoma risk.

**Conclusion:** This study confirms that elevated IOP is a key characteristic of glaucoma. However, age, female gender, and hypertension were not recognised as important risk factors for glaucoma. This suggests that other risk factors might have contributed to the risk of glaucoma. Screening beyond traditional risk factors helps in identifying at-risk patients.

**Keywords:** case-control study, hypertension, incidence, open-angle glaucoma, risk factors

### Introduction

Glaucoma is a major cause of irreversible vision loss and blindness, marked by progressive damage to the optic nerve. In 2020, approximately 76 million individuals worldwide were affected, including 4.5 million with moderate to severe visual impairment and 3.2 million with blindness, and the total number of individuals affected with glaucoma is projected to rise to 111.8 million by 2040.<sup>[1,2]</sup> Because the disease is often asymptomatic in its early stages, diagnosis typically occurs only after substantial optic nerve damage, leading to compromised quality of life.<sup>[1]</sup> Early detection and timely management are therefore essential to prevent further vision loss.<sup>[1,2]</sup>

Glaucoma is classified into multiple subtypes, such as primary open-angle glaucoma (POAG), primary angle-closure glaucoma (PACG), secondary glaucoma, and congenital glaucoma,<sup>[3]</sup> with their prevalence varying across ethnic and geographic populations. Established risk factors include modifiable factors (elevated intraocular pressure, hypertension, steroid use) and non-modifiable factors (age, sex, family history, thinning of the central cornea).<sup>[1,4]</sup> Additionally, systemic conditions like diabetes mellitus, hypothyroidism, and corticosteroid use may predispose individuals to secondary glaucoma. Identifying and addressing modifiable risk factors is critical in reducing disease progression and raising awareness among both clinicians and patients.<sup>[1]</sup>

While the epidemiology and clinical impact of glaucoma are well studied in developed nations, limited data exist for developing countries such as India.<sup>[5,6]</sup> Evidence suggests that patients in these regions present late with more advanced disease, face a higher risk of blindness, and may have a greater age-adjusted incidence compared to those in developed countries, though the reasons remain unclear.<sup>[5]</sup> This study sought to address this gap by assessing the incidence of glaucoma and evaluating associated risk factors in developing countries.

### Method

This single-centre prospective case-controlled study was conducted over a one-year period in 2023 at the Department of Ophthalmology, Dr. Kodkany's Eye Center, Belagavi. It was designed to evaluate selected risk factors for glaucoma based on predefined objectives and feasibility, and does not aim to assess all known risk factors.

Adults aged  $\geq 40$  years with an intraocular pressure (IOP)

$> 21$  mmHg, vertical cup-disc ratio (VCDR)  $> 0.6$ , or other glaucomatous optic disc changes such as polar notch, glaucomatous field defects (Anderson criteria), deep cupping, disc haemorrhages, thinning of the neuroretinal rim, or rim change (characteristic glaucomatous alterations of the neuroretinal rim) in both eyes were included as cases. A diagnosis of glaucoma was confirmed if at least two of these three criteria were present. Controls were recruited from patients aged  $\geq 40$  years without glaucoma, excluding those incidentally diagnosed during the study period.

All participants underwent clinical evaluation using a standardized protocol including visual acuity assessment, IOP measurement, and detailed systemic and ocular history. Glaucoma cases underwent a comprehensive ophthalmic examination as part of the diagnostic assessment, including central corneal thickness (CCT), anterior segment abnormalities, anterior chamber angle, fundus examination, and visual field testing. These detailed investigations were not uniformly performed in control participants, as they were recruited as individuals without clinical evidence of glaucoma and underwent standardized screening evaluation, including IOP measurement and clinical assessment, to reasonably exclude the presence of glaucoma.

Written informed consent was obtained from all participants. Ethical approval was obtained from the Institutional Ethics Committee of Prabhakar Kore's Basic Science Research Center, Belagavi. Sociodemographic and clinical details were recorded using a predesigned, pretested questionnaire.

Data were analysed using SPSS software version 21. Categorical variables were presented as frequencies or proportions and were analysed using the Chi-square test. Continuous variables are presented as Mean  $\pm$  standard deviation (SD). Normality of the data was assessed using the Shapiro-Wilk test and Q-Q plots. Independent-samples t-tests or Mann-Whitney U tests were used to compare continuous variables between groups, depending on normality. A p-value less than or equal to 0.05 indicates statistical significance.

### Results

A total of 44 participants were included in the study, comprising 24 patients with glaucoma and 20 age- and sex-matched controls. Most participants in both groups were aged 56–65 years and predominantly male. Most were Hindus, married, had completed at least secondary

**Table 1. Sociodemographic characteristics of study participants**

| Variable              | Patients with glaucoma<br>(N=24) n (%) | Patients without glaucoma<br>(N=20) n (%) | p-value            |                     |           |           |
|-----------------------|----------------------------------------|-------------------------------------------|--------------------|---------------------|-----------|-----------|
| <b>Age</b>            |                                        |                                           |                    | Secondary           | 7 (29.2)  | 1 (5.0)   |
| 45-55                 | 5 (20.8)                               | 5 (25.0)                                  | 0.929 <sup>c</sup> | Secondary and above | 8 (33.3)  | 7 (35.0)  |
| 56-65                 | 10 (41.7)                              | 8 (40.0)                                  |                    | Graduation          | 2 (8.3)   | 3 (15.0)  |
| 66-75                 | 5 (20.8)                               | 5 (25.0)                                  |                    | <b>Occupation</b>   |           |           |
| 76-85                 | 3 (12.5)                               | 1 (5.0)                                   |                    | Doctor              | 1 (4.2)   | 0.0       |
| >85                   | 1 (4.2)                                | 1 (5.0)                                   |                    | Employed            | 1 (4.2)   | 1 (5.0)   |
| <b>Sex</b>            |                                        |                                           |                    | Farmer              | 3 (12.5)  | 2 (10.0)  |
| Female                | 9 (37.5)                               | 10 (50.0)                                 | 0.404 <sup>c</sup> | Housewife           | 5 (20.8)  | 6 (30.0)  |
| Male                  | 15 (62.5)                              | 10 (50.0)                                 |                    | Housewife/skilled   | 0.0       | 1 (5.0)   |
| <b>Religion</b>       |                                        |                                           |                    | Retired             | 6 (25)    | 3 (15.0)  |
| Hindu                 | 21 (87.5)                              | 20 (100.0)                                | 0.101 <sup>c</sup> | Semiskilled         | 0 (0.0)   | 1 (5.0)   |
| Muslim                | 3 (12.5)                               | 0                                         |                    | Shop                | 2 (8.3)   | 5 (25.0)  |
| <b>Marital status</b> |                                        |                                           |                    | Shop/Farmer         | 1 (4.2)   | 1 (5.0)   |
| Married               | 19 (79.2)                              | 16 (80.0)                                 | 0.945 <sup>c</sup> | Skilled/employed    | 2 (8.3)   | 0.0       |
| Widowed               | 5 (20.8)                               | 4 (20.0)                                  |                    | Skilled/retired     | 1 (4.2)   | 0.0       |
| <b>Education</b>      |                                        |                                           |                    | Unskilled           | 2 (8.3)   | 0.0       |
| Illiterate            | 3 (12.5)                               | 4 (20.0)                                  | 0.319 <sup>c</sup> | <b>Residence</b>    |           |           |
| Primary               | 4 (16.7)                               | 5 (25.0)                                  |                    | Rural               | 11 (45.8) | 14 (70.0) |
|                       |                                        |                                           |                    | Urban               | 13 (54.2) | 6 (30.0)  |

c - values calculated using the chi-square test.

**Table 2. Comparison of intraocular pressure and systemic disorders in cases and controls**

| Variable                        | Patients with glaucoma<br>(N=24) | Patients without glaucoma<br>(N=20) | Mean Difference and 95% CI                     | p-value             |
|---------------------------------|----------------------------------|-------------------------------------|------------------------------------------------|---------------------|
| Right eye (mmHg)                | 27.86 ± 8.97                     | 15.85 ± 2.84                        | Mean difference: 12.01<br>(95% CI: 8.24–15.78) | <0.001*             |
| Left eye (mmHg)                 | 28.27 ± 8.75                     | 16.05 ± 3.02                        | Mean difference: 12.22<br>(95% CI: 8.49–15.95) | <0.001*             |
| <b>Other systemic disorders</b> |                                  |                                     |                                                |                     |
| Present                         | 21 (87.5%)                       | 7 (35%)                             | NA                                             | 0.003* <sup>c</sup> |
| Absent                          | 3 (12.5%)                        | 13 (65%)                            |                                                |                     |

\*- statistically significant. c - values calculated using the chi-square test.

education, resided in rural areas, and were either housewives or retired (Table 1). However, no significant associations were observed between sociodemographic characteristics and glaucoma risk.

IOP was significantly higher among glaucoma patients compared to controls in both eyes (right eye:  $27.86 \pm 8.97$

vs  $15.85 \pm 2.84$  mmHg [Mean difference: 12.01 (95% CI: 8.24–15.78]; left eye:  $28.27 \pm 8.75$  vs  $16.05 \pm 3.02$  mmHg; [Mean difference: 12.22 (95% CI: 8.49–15.95)]  $p < 0.001$  for both) (Table 2). The presence of any systemic comorbidity (considered as a composite variable) was significantly associated with glaucoma ( $p < 0.05$ ). Cataract

**Table 3. Distribution of glaucoma types, affected eyes, and vision loss in the study participants**

| Variable                                        | N = 24 n (%)       | Cup to disc ratio - Left eye                                  | 0.719 $\pm$ 0.15 |
|-------------------------------------------------|--------------------|---------------------------------------------------------------|------------------|
| <b>Diagnosis</b>                                |                    | <b>Visual field changes<sup>#</sup></b>                       |                  |
| Disc suspect                                    | 2 (8.3)            | Arcuate defect (BE)                                           | 1 (4.2)          |
| Narrow angle glaucoma                           | 1 (4.2)            | Asteroid hyalosis (BE)                                        | 1 (4.2)          |
| Normal tension glaucoma                         | 1 (4.2)            | Centrocaecal scotoma (BE)                                     | 1 (4.2)          |
| Open angle glaucoma                             | 14 (58.3)          | Complete depressed field (BE)                                 | 1 (4.2)          |
| Ocular pulse amplitude                          | 1 (4.2)            | Congestion (BE), Deep cup thin (BE)                           | 1 (4.2)          |
| Primary open angle glaucoma                     | 3 (12.5)           | Deep cupping                                                  | 1 (4.2)          |
| Pseudo exfoliation glaucoma                     | 2 (8.3)            | Enlarged blind spot (RE), Vision loss (LE)                    | 3 (12.5)         |
| <b>Affected eye</b>                             |                    | Generalized depression (BE)                                   | 1 (4.2)          |
| Both eyes                                       | 13 (54.2)          | Inferior arcuate scotoma (BE)                                 | 1 (4.2)          |
| Left eye                                        | 7 (29.2)           | Mild congestion (BE)                                          | 1 (4.2)          |
| Right eye                                       | 4 (16.7)           | Notching in (LE)                                              | 1 (4.2)          |
| <b>Vision loss<sup>#</sup></b>                  |                    | Rim loss (RE), Mild congestion in (LE)                        | 1 (4.2)          |
| Left eye                                        | 3 (60)             | Shallow anterior, vision loss (LE)                            | 1 (4.2)          |
| Right eye                                       | 1 (20)             | Superior arcuate scotoma (BE)                                 | 2 (8.3)          |
| Steroid induced                                 | 1 (20)             | Superior arcuate scotoma Ben Notching (RE), Deep cupping (LE) | 1 (4.2)          |
| Central corneal thickness - Right eye (microns) | 541.08 $\pm$ 18.82 | Superior arcuate scotoma (RE)                                 | 1 (4.2)          |
| Central corneal thickness -Left eye (microns)   | 542.42 $\pm$ 18.39 | Superior hemifield defect (RE)                                | 1 (4.2)          |
| Cup to disc ratio -Right eye                    | 0.771 $\pm$ 0.14   | Total constriction (RE)                                       | 2 (8.3)          |

<sup>#</sup>, Data not available: data for vision loss and visual field changes were not available for all participants hence percentages are only calculated based on the number with available data.

**Table 4. Comparison of available clinical and systemic parameters between glaucoma cases and controls**

| Univariate analysis  |                   |                   |                       |         |
|----------------------|-------------------|-------------------|-----------------------|---------|
|                      | Glaucoma (n=24)   | Control (n=20)    | Unadjusted OR (95%CI) | p-value |
| <b>Hypertension</b>  | 11 (45.8%)        | 5 (25%)           | 2.538 (0.697, 9.240)  | 0.160   |
| <b>Age</b>           | 64.42 $\pm$ 11.17 | 63.15 $\pm$ 11.38 | 0.990 (0.937, 1.045)  | 0.705   |
| <b>Gender-Female</b> | 9 (37.5%)         | 10 (50%)          | 1.667 (0.500, 5.559)  | 0.416   |

OR – Odds Ratio, CI – Confidence interval

was found in both groups. Among glaucoma patients, six (25%) had only cataract, five (20.8%) had cataract with hypertension, and three (12.5%) had cataract, hypertension, and diabetes. Among non-glaucoma patients, two (10%) had only cataract, and one had cataract, hypertension, and diabetes mellitus.

IOP and systemic parameters were available for both cases and controls and were included in comparative analysis. However, detailed ophthalmic parameters, such as central corneal thickness and cup-to-disc ratio, were not uniformly recorded in control participants; therefore, comparative analysis of these variables could not be performed.

Regarding glaucoma subtypes, open-angle glaucoma was the most frequent diagnosis, observed in 14 (58.3%) patients. Other forms included primary open-angle glaucoma (12.5%), pseudo-exfoliation glaucoma (8.3%), and normal-tension glaucoma (4.2%). Bilateral involvement was noted in 13 (54.2%) patients, with the left eye more frequently affected than the right eye. (Table 3).

Central corneal thickness was comparable between right ( $541.08 \pm 18.82 \mu\text{m}$ ) and left eyes ( $542.42 \pm 18.39 \mu\text{m}$ ). The mean cup-to-disc ratio was  $0.771 \pm 0.14$  in the right eye and  $0.719 \pm 0.15$  in the left eye. Visual field changes varied, with superior arcuate scotoma (12.5%) and total field constriction (8.3%) being the most common defects (Table 3). Univariate analysis did not identify age, female gender, or hypertension as significant predictors of glaucoma risk ( $p > 0.05$  for all comparisons) (Table 4).

## Discussion

This case–controlled study demonstrated a significantly higher IOP in glaucoma patients compared with controls (show difference, CI, p-value) ( $p < 0.001$ ), reaffirming its central role in glaucoma pathogenesis. Although univariate analysis indicated that female patients and those with hypertension had higher odds of developing glaucoma (1.6-fold and 2.5-fold, respectively), these associations were not statistically significant.

The majority of glaucoma patients in this study were aged 55 years or older, with a predominance of males (62.5%), consistent with earlier reports.<sup>[1,7]</sup> Our study shows that hypertension and diabetes mellitus are common systemic risk factors, while Talaat et al<sup>[7]</sup> similarly reported hypertension (60.8%) and diabetes (58.3%) as frequent systemic comorbidities among glaucoma patients.

Cataract (49.1%) was the most common ocular condition.<sup>[7]</sup> In our cohort, primary open-angle glaucoma (POAG) was the

most prevalent subtype (58.3%), with more than half of the patients showing bilateral involvement. Diverse visual field abnormalities, including arcuate defects, scotomas, and generalized field depression, were observed, highlighting the heterogeneous presentation of glaucomatous damage.<sup>[8,9]</sup>

In the present study, IOP was significantly higher in glaucoma cases compared to controls. In the context of pseudo-exfoliation, previous studies have similarly demonstrated that a 5 mmHg increase in IOP may increase the risk of primary open-angle glaucoma by up to fourfold.<sup>[5]</sup>

Interestingly, our study did not confirm associations with other established risk factors such as age, sex, or hypertension. This contrasts with meta-analytic evidence suggesting that advancing age markedly increases POAG risk and that females may have a lower overall risk than males. One possible explanation for this difference could be the relatively small sample size and limited statistical power of the present study, which may have reduced the ability to detect such associations.

Other studies have identified ocular biometric features (axial length, corneal curvature), older age, and elevated IOP as significant contributors to glaucoma risk.<sup>[10]</sup> Furthermore, hormonal influences, such as reduced oestrogen in postmenopausal women, have been implicated in increasing glaucoma susceptibility.<sup>[11]</sup> Longitudinal studies also support associations between systemic conditions—including hypertension, diabetes, and hyperglycaemia—and glaucoma development.<sup>[12]</sup> Additionally, research among Bangladeshi populations indicated a modest association between hypertension and elevated IOP, possibly mediated through increased sympathetic activity.<sup>[13,14]</sup> In contrast, our findings did not demonstrate a statistically significant link between hypertension and glaucoma.

Overall, this study reinforces IOP as the key modifiable risk factor in glaucoma pathogenesis. Routine IOP screening may therefore be more effective in identifying individuals at risk than monitoring systemic comorbidities alone. Nonetheless, certain limitations should be acknowledged. The relatively small sample size restricts the statistical power and may have precluded the detection of meaningful associations. Larger, well-designed case–control studies that use multivariate analyses to adjust for confounders are needed to clarify the contributions of systemic and ocular risk factors. Future research should also evaluate the roles of lifestyle factors (such as alcohol consumption,

smoking, and diet) and genetic predisposition in glaucoma pathogenesis.

### Conclusion

This study reaffirms that IOP remains a defining and significant feature of glaucoma, underscoring its role as the most important modifiable risk factor in disease pathogenesis. Steroid use and other non-modifiable risk factors, including a VCDR >0.4 and inter-eye CDR asymmetry >0.2, should prompt physicians to further evaluate patients for glaucoma, even in the presence of normal intraocular pressure. Although age, female gender, and hypertension were evaluated, none showed a statistically significant association with glaucoma risk in this cohort. Early identification of individuals at risk, particularly in resource-limited settings where glaucoma often presents at an advanced stage, is critical to prevent irreversible vision loss.

**Funding:** None

**Conflict of interests:** None

### References

1. Shan S, Liu Y, Zhang Z, et al. Global incidence and risk factors for glaucoma: a systematic review and meta-analysis of prospective studies. *J Glob Health*. 2024;14:04252. doi: 10.7189/jogh.14.04252 <https://pubmed.ncbi.nlm.nih.gov/39513294/>
2. Delgado ME, Sánchez FG, Restrepo JC, Valencia M. Management of glaucoma in developing countries: challenges and opportunities for improvement. *Clinicoecon Outcomes Res*. 2019;11:591–604. doi: <https://doi.org/10.2147/CEOR.S218277> <https://pmc.ncbi.nlm.nih.gov/articles/PMC6776288/>
3. Dietze, J., Blair, K., & Havens, S. J. (2024). Glaucoma [Continuing education activity]. In StatPearls. StatPearls Publishing. PMID 30855805. <https://www.ncbi.nlm.nih.gov/books/NBK538217/>
4. Bae HW, Lee N, Kim CY, Seong GJ, Kim SH, et al. Systemic hypertension as a risk factor for open-angle glaucoma: a meta-analysis of population-based studies. *PLoS One*. 2014;9(9):e108226. doi: <https://doi.org/10.1371/journal.pone.0108226> <https://pubmed.ncbi.nlm.nih.gov/25254373/>
5. Ekici E, Turan N, Bayer A, Altinsoy HI. Central visual field defects in patients with distinct glaucomatous optic disc phenotypes. *Am J Ophthalmol*. 2021;223:229–40. doi: 10.1016/j.ajo.2020.10.015. <https://pmc.ncbi.nlm.nih.gov/articles/PMC7979468/>
6. Le A, Mukesh BN, McCarty CA, Taylor HR. Risk factors associated with the incidence of open-angle glaucoma: the visual impairment project. *Investigative ophthalmology & visual science* 2003; 44:3783-9. <https://doi.org/10.1167/iops.03-0077>
7. Talaat K, Althubaiti G, Alzahrani A, et al. Types of glaucoma and associated comorbidities among patients at King Abdulaziz Medical City, Jeddah. *Cureus*. 2021;13(6):e15934. doi: 10.7759/cureus.15574 <https://pubmed.ncbi.nlm.nih.gov/34277196/>
8. Ekici E, Moghimi S, Hou H, Proudfoot J, Zangwill LM, Do JL, et al. Central Visual Field Defects in Patients with Distinct Glaucomatous Optic Disc Phenotypes. *Am J Ophthalmol* 2021; 223:229-40. <https://doi.org/10.1016/j.ajo.2020.09.034>
9. Kim KE, Kim JM, Lee J, Lee MY, Park KH. Significant intraocular pressure associated with open-angle glaucoma: Korea National Health and Nutrition Examination Survey 2010-2011. *PLoS One* 2020; 15:e0235701. <https://doi.org/10.1371/journal.pone.0235701>
10. Hashemi H, Mohammadi M, Zandvakil N, Khabazkhoob M, Emamian MH, Shariati M, et al. Prevalence and risk factors of glaucoma in an adult population from Shahroud, Iran. *Journal of current ophthalmology* 2019; 31:366-72. <https://doi.org/10.1016/j.joco.2018.05.003>
11. Rizk M, Grise-Dulac A, Gatinel D. Glaucoma in Women: What do we Know so Far-A Systematic Review. *AJO International* 2024;100013. . <https://doi.org/10.1016/j.ajo.2024.100013>
12. Newman-Casey PA, Talwar N, Nan B, Musch DC, Stein JD. The relationship between components of metabolic syndrome and open-angle glaucoma. *Ophthalmology* 2011; 118:1318-26. <https://doi.org/10.1016/j.opth.2010.11.020>
13. Ramesh PV, Morya AK, Aradhya AK, Pannerselvam P, Gopalakrishnan ST, Ramesh SV, et al. Unveiling the silent link: Normal-tension glaucoma's enigmatic bond with cardiac blood flow. *World J Cardiol* 2024; 16:10-5. <https://doi.org/10.4330/wjc.v16.i1.10>
14. Ahmed MS, Ullah AY, Barman N, Ratan ZA, Mostafa S, Khaleque A, et al. Risk factors associated with elevated intraocular pressure: a population-based study in a rural community of Bangladesh. *BMJ Open Ophthalmology* 2023; 8:e001386. <https://doi.org/10.1136/bmjophth-2023-001386>

# OV16 antibody seropositivity reveals persistent onchocerciasis transmission and cross-border risk in Magwi County, South Sudan: a population-based survey

Ezbon WApary<sup>1</sup>,  Yak Yak<sup>2</sup>, Benson Ojara<sup>3</sup>, Wilson Ladu<sup>2</sup>, David Oguttu<sup>4</sup>, Ian Hennessee<sup>5</sup>, Pita Jane<sup>6</sup>, Johan Willems<sup>3</sup>, Thok Chol<sup>2</sup>, Kamulete Horasio<sup>3</sup>, Okot William<sup>2</sup>, Amany Jacob<sup>2</sup>

## Author Affiliation:

1. University of Juba, Juba, South Sudan
2. Ministry of Health, Juba, South Sudan
3. Christian Blind Mission, Juba, South Sudan
4. Ministry of Health, Kampala, Uganda
5. The Carter Center, Juba, South Sudan
6. World Health Organization, Juba, South Sudan

## Correspondence:

Ezbon WApary  
[aulyaiezbon@gmail.com](mailto:aulyaiezbon@gmail.com)

**Submitted:** March 2026

**Accepted:** April 2026

**Published:** May 2026

**Citation:** WApary et al. OV16 antibody seropositivity reveals persistent onchocerciasis transmission and cross-border risk in Magwi County, South Sudan: a population-based survey. *South Sudan Medical Journal*, 2026;19(2):81-86 © 2026 The Author (s) **License:** This is an open access article under [CC BY-NC](https://creativecommons.org/licenses/by-nc/4.0/) DOI: <https://dx.doi.org/10.4314/ssmj.v19i2.3>

## ABSTRACT

**Introduction:** Onchocerciasis, or river blindness, is the second leading infectious cause of blindness after trachoma. It is a parasitic disease caused by *Onchocerca volvulus*, a filarial worm transmitted by black flies. In 34 countries, 249.5 million people are at risk, of which 14.6 million are infected, including 1.15 million people who are visually impaired or have gone blind. Onchocerciasis remains endemic in many parts of South Sudan. Magwi is an onchocerciasis-endemic county in South Sudan bordering northern Uganda districts where onchocerciasis transmission has been suppressed. The study was conducted in Magwi County to determine the prevalence of onchocerciasis by testing for *Onchocerca volvulus*-16 (OV16) antibodies to estimate the risk of cross-border reintroduction of onchocerciasis into northern Uganda.

**Method:** The study adapted a community-based cross-sectional serological survey in Magwi County of South Sudan. The study recruited children aged 6-9 years. The study used a stratified cluster sampling approach, informed by data on population distribution and the proximity of communities to rivers infested with blackflies. Blood samples were collected using filter paper as dried blood spots. ELISA testing was done in the Uganda Onchocerciasis Reference Laboratory to confirm OV16 antibodies. Analysis software was Stata v17. The ethical standards were adhered to.

**Results:** 3,290 children participated in the study. 1,746 (53.1%) were males, and 6-7 years were 1,723 (52.4%). The overall OV16 seropositivity was 0.9%. Among the children, those aged 8-9 years had the highest seropositivity prevalence of 17 (1.1%). Across the clusters, transmission varied statistically ( $\chi^2 = 15.23$ ,  $p < 0.001$ ), with a small effect size (Cramer's  $V = 0.068$ ). Whereas most clusters had no positive cases, the highest seroprevalence was in Garamu (9.6%) which followed by Malakia Central (7.6%), Olikwi (4.2%), and Kamuli (2.7%).

**Conclusion:** Persistent onchocerciasis transmission in Magwi County poses a risk of cross-border transmission into Northern Uganda.

**Key words:** OV16 antibody seropositivity, onchocerciasis transmission, cross-border, population-based survey, Magwi County, South Sudan

### Introduction

Onchocerciasis, or river blindness, is the second leading infectious cause of blindness after trachoma. It is a parasitic disease caused by *Onchocerca volvulus*, a filarial worm transmitted by blackfly bites. In 34 countries, 249.5 million people are at risk of which 14.6 million are infected, 1.15 million people are visually impaired and/or have gone blind.<sup>[1]</sup> Twenty-six African countries shouldered 96% of all global cases.<sup>[2]</sup>

In South Sudan, Rapid Epidemiological Mapping of Onchocerciasis identified 48 endemic counties out of total 80 counties.<sup>[3]</sup> These areas are inhabited by communities classified as priority areas requiring Community-Directed Treatment with Ivermectin (CDTI), areas not requiring treatment, and possible endemic areas that need further investigation. The main endemic foci are in Western Bahr El-Ghazal, Western Equatoria, Central Equatoria, and Lakes states. In Unity State, the prevalence was <20%. Despite the country shifting focus from control to elimination, Mass Drug Administration (MDA) with Ivermectin was carried out in 2022 in all the 48 identified endemic counties for onchocerciasis, of which 34 were treated for onchocerciasis with >5 effective rounds of MDA (≥80% coverage) and lymphatic filariasis/Oncho-LF (co-endemic), while 14 were treated for onchocerciasis only.<sup>[3]</sup>

Onchocerciasis MDA has been implemented since 2005. The aim is to achieve ≥80% ivermectin coverage. The target set for 2027 on onchocerciasis is that 30% of counties should have suppressed transmission. The percentage of counties with suppressed transmission in 2023 was 0% and was used as the baseline; in 2025, it was 20% as a midterm.<sup>[3]</sup> Magwi County was selected for an onchocerciasis prevalence survey due to its proximity to northern Uganda districts where the transmission is suppressed. Additionally, population movements, farming near rivers, and environmental disruption maintain exposure to blackflies.

The *Onchocerca volvulus*-16 (OV16) antibody, specifically the human IgG4 subclass, is a serological biomarker used to detect exposure to *Onchocerca volvulus*. Monitoring this antibody is a critical component of elimination programs as it serves as an early marker of infection, often detectable months before the parasite's larvae (microfilariae) appear in the skin.<sup>[4]</sup>

The study was conducted in Magwi County to monitor the prevalence of onchocerciasis by testing for OV16 antibody

to estimate the risk of cross-border reintroduction of onchocerciasis into northern Uganda.

### Method

The study adapted a community-based cross-sectional serological survey design utilising a stratified cluster sampling technique.<sup>[5]</sup> The survey covered 32 clusters in Magwi County (Figure 1), selected based on proximity to Simulium breeding sites. The breeding sites were identified by walking along the rivers, locating fast-flowing sections with suitable submerged or trailing vegetation.

The survey included children aged 6-9 years, as they serve as a core indicator of recent transmission; resided in these clusters for at least 5 years; had guardians' or parents' consent; and assented. The exclusion criteria were individuals with temporary or recent (<5 years) residence in endemic areas; individuals with bleeding disorders or medical contraindications to finger-prick sampling.

The study engaged with community leaders and sensitisation on the survey objectives was carried out. Initial meetings were with community leaders and local government. Verbal and written sensitisation was done to inform communities about the survey. The study recruited children from house-to-house who met the inclusion criteria, recorded their demographic data, pricked their fingers to collect blood samples on filter paper, safely discarded sharps into safety boxes, and disposed daily other biohazardous waste in the nearest health facility's incinerators. Anti-OV16 ELISA testing was done in the Uganda Onchocerciasis Reference Laboratory.

The survey teams were trained on the study protocol with emphasis on community entry, sensitisation technique, ethical guidelines, tools for data collection, DBS sampling procedures, and sample handling. Indoor and outdoor practicums were conducted prior to the survey's commencement to ensure the quality of the survey team members.

### Ethical consideration

The ethical standards were adhered to, with reference number RERB/A/75/2025 obtained from the Research and Ethical Review Board, Ministry of Health, South Sudan. Written consent was obtained from the responsible party, with assent from minors (children). As the names were de-identified, the participant codes and sample codes were the same for each participant.




Figure 1. Location of sites within Magwi county, Eastern Equatoria State. Background data from the Humanitarian Data Exchange (<https://data.humdata.org>).

## Data analysis

Field data were gathered using ESPEN Collect installed on mobile devices. Data collected included coded informed consent and demographic characteristics, and were submitted daily into the ESPEN-hosted server. The data were exported into Stata v17 software for Analysis. Descriptive statistics summarised participant characteristics, including age, sex, and cluster distribution. Seroprevalence of OV16 antibodies was calculated as proportions with corresponding percentages. Bivariate analysis was conducted to assess associations between OV16 seropositivity and categorical variables (age group, sex, and geographic clusters) using the chi-squared ( $\chi^2$ ) test. Effect sizes were estimated using Cramer's V to assess the strength of associations. Cluster-level seroprevalence was analysed to identify spatial heterogeneity and potential transmission hotspots. Statistical significance was set at  $p < 0.05$ . Results were presented in tables and interpreted in the context of onchocerciasis elimination thresholds.

## Results

Of the 3,290 children who participated in the study,

males comprised 53.1%, 6-year-olds comprised 33.3%, and 42.1% were from Nimule Payam. The overall OV16 seropositivity was 0.9%. Among the children, those aged 8-9 years had the highest seroprevalence of 1.1%. Age and sex showed no statistical significance, indicating even exposure to the transmission. The effect size (Cramer's V) further confirms the absence of a strong relationship (Table 1).

Children residing in the Madi Corridor clusters had a higher seropositivity of 1.5% compared to 0.02% in the Acholi Corridor (Table 1). The highest seroprevalence was in Garamu (9.6%), followed by Malakia Central (7.6%), Olikiwi (4.2%), and Kamuli (2.7%) (Table 2).

## Discussion

Analysis of the ELISA findings revealed that 0.9% of the children aged 6-9 years tested positive for OV16 antibodies. This low seroprevalence aligns with results from similar surveys in hypo-endemic settings, indicating limited ongoing transmission of *Onchocerca volvulus* in the county, or, rather, in South Sudan.<sup>[6,7]</sup> The 0.9%

**Table 1. Seropositivity of onchocerciasis in children by age, sex and location**

|                     | ELISA Test Results |               | Total<br>n (%) | Chi-squared<br>(p-value) | Cramer's<br>V |
|---------------------|--------------------|---------------|----------------|--------------------------|---------------|
|                     | - ve<br>n (%)      | + ve<br>n (%) |                |                          |               |
| <b>Age in years</b> |                    |               |                | 0.99 (0.319)             | 0.017         |
| 6-7                 | 1710 (99.2)        | 13 (0.8)      | 1723 (52.4)    |                          |               |
| 8-9                 | 1550 (98.9)        | 17 (1.1)      | 1567 (47.6)    |                          |               |
| <b>Sex</b>          |                    |               |                | 0.01 (0.977)             | 0.001         |
| Male                | 1730 (99.1)        | 16 (0.9)      | 1746 (53.1)    |                          |               |
| Female              | 1530 (99.1)        | 14 (0.9)      | 1544 (46.9)    |                          |               |
| <b>Clusters</b>     |                    |               |                | 15.23 (<.001)            | 0.068         |
| Acholi Corridor     | 1601 (99.8))       | 4 (0.2)       | 1605 (48.8)    |                          |               |
| Madi Corridor       | 1659 (98.5)        | 26 (1.5)      | 1685 (51.2)    |                          |               |
| <b>Total</b>        | 3260 (99.1)        | 30 (0.9)      | 3290 (100)     |                          |               |

OV16 seropositivity exceeds the 0.1% WHO threshold for stopping MDA with ivermectin and indicates ongoing transmission in Magwi County.<sup>[5]</sup> Therefore, CDTI should continue in Magwi County and, where appropriate, be intensified, especially in identified hotspots, to prevent cross-border transmission into northern Uganda.

Children aged nine years exhibited the highest seropositivity compared to those aged 6-8 years. Despite this trend, the results showed no statistically significant differences, implying that age may not be a strong predictor of exposure in this population.<sup>[8]</sup>

Even though overall seropositivity is minimal, the detection of seropositive cases among these age groups reflects residual pockets of transmission. In East African foci, studies have revealed that low seropositivity can continue post-ivermectin MDA, necessitating targeted follow-up interventions.<sup>[9,10]</sup> The OV16 ELISA remains a sensitive and specific tool for detecting antibodies in these children, providing vital data to inform onchocerciasis elimination efforts.<sup>[11,12]</sup>

Seropositivity was homogeneously low across these age groups, extending from 0.4% to 1.4%. Similarly, the effects of age have been weak or absent in some endemic settings when analysis inquiry is restricted to a limited age group, whereas broader age ranges in adult populations, such as livestock or humans, show more often strong age gradients in seropositivity.<sup>[13,14]</sup>

Male and female children had similar seroprevalence (0.9%) with no statistically detectable association between ELISA status and sex. This contrasts with many occupationally mediated or sexually transmitted infections, where men often show higher seroprevalence or clustering,<sup>[13,15,16]</sup> and with brucellosis in cattle in South

Sudan, where females had substantially greater likelihood of seropositivity than males.<sup>[14]</sup> The absence of variation suggests that exposure is likely transmitted through shared household or environmental factors rather than through sex-related behaviours.

Clusters were observed as strong determinants of transmission homogeneity (e.g. Garamu, Malakia Central, Malakia West A, Olikwi) showed seroprevalence between 3.1% and 9.6%. The higher seropositivity in the Madi Corridor may indicate favourable localised ecological factors for vector breeding or transmission. This is analogous to riverside villages in Cameroon, Nigeria, and Ethiopia, where higher biting and infection are documented.<sup>[17-20]</sup> This pattern is consistent with focal "hotspots" of transmission embedded in a largely low-risk landscape, as seen for HIV, brucellosis, and other infections where particular communities or locations account for a disproportionate share of ongoing transmission.<sup>[13-16]</sup> In such settings, spatially targeted interventions in high-prevalence clusters may be more efficient than uniform county-wide measures.

Several limitations were considered. OV-16 antibodies indicate exposure; however, they may persist after treatment or infection. Entomological investigations were not concurrently conducted. This limits direct inference about the intensity of the transmission. Mobility of population data may also affect interpretation. Despite these limitations, the significance of seroprevalence strongly indicates that transmission of onchocerciasis has not yet been interrupted.

## Conclusion

Persistent transmission of onchocerciasis in Magwi

**Table 2 Seropositivity of onchocerciasis in children by clusters within Magwi county**

|                          | ELISA Test Results |               | Total<br>n (%) |
|--------------------------|--------------------|---------------|----------------|
|                          | - ve<br>n (%)      | + ve<br>n (%) |                |
| Acholi Corridor Clusters |                    |               |                |
| Angok                    | 110 (100)          | 0 (0)         | 110 (3.3)      |
| Ayii                     | 109 (100)          | 0 (0)         | 109 (3.3)      |
| Buchingo                 | 111 (100)          | 0 (0)         | 111 (3.4)      |
| Chama                    | 110 (100)          | 0 (0)         | 110 (3.3)      |
| Gara                     | 109 (99.1)         | 1 (0.9)       | 110 (3.3)      |
| Kamuli                   | 107 (97.3)         | 3 (2.7)       | 110 (3.3)      |
| Kilio                    | 110 (100)          | 0 (0)         | 110 (3.3)      |
| Lawaci                   | 112 (100)          | 0 (0)         | 112 (3.4)      |
| Lerwa                    | 111 (100)          | 0 (0)         | 111 (3.4)      |
| Loudo                    | 110 (100)          | 0 (0)         | 110 (3.3)      |
| Obokolot                 | 110 (100)          | 0 (0)         | 110 (3.3)      |
| Ogwal-Woo                | 110 (100)          | 0 (0)         | 110 (3.3)      |
| Opamo                    | 110 (100)          | 0 (0)         | 110 (3.3)      |
| Opari                    | 62 (100)           | 0 (0)         | 62 (1.9)       |
| Palwar                   | 110 (100)          | 0 (0)         | 110 (3.3)      |
| Madi Corridor Clusters   |                    |               |                |
| Abila West               | 123 (100)          | 0 (0)         | 123 (3.7)      |
| Avumadrici               | 42 (100)           | 0 (0)         | 42 (1.3)       |
| Beka-Angwa               | 64 (100)           | 0 (0)         | 64 (1.9)       |
| Coyi                     | 33 (100)           | 0 (0)         | 33 (1)         |
| Ganzi                    | 52 (100)           | 0 (0)         | 52 (1.6)       |
| Garamu                   | 75 (90.4)          | 8 (9.6)       | 83 (2.5)       |
| Hai Kenisa West          | 138 (100)          | 0 (0)         | 138 (4.2)      |
| Kololo/Jelie             | 99 (100)           | 0 (0)         | 99 (3)         |
| Central                  |                    |               |                |
| Malakia Central          | 109 (92.4)         | 9 (7.6)       | 118 (3.6)      |
| Malakia West A           | 125 (96.9)         | 4 (3.1)       | 129 (3.9)      |
| Masindi                  | 56 (100)           | 0 (0)         | 56 (1.7)       |
| Moi-Oci-Kwira            | 34 (100)           | 0 (0)         | 34 (1)         |
| Motoyo West A            | 148 (100)          | 0 (0)         | 148 (4.5)      |
| Motoyo West B            | 149 (100)          | 0 (0)         | 149 (4.5)      |
| Olikwi                   | 113 (95.8)         | 5 (4.2)       | 118 (3.6)      |
| Rei West                 | 150 (100)          | 0 (0)         | 150 (4.6)      |
| Sekuwa East              | 149 (100)          | 0 (0)         | 149 (4.5)      |
| Total                    | 3260 (99.1)        | 30 (0.9)      | 3290 (100)     |

County, South Sudan, poses a risk of cross-border transmission into Northern Uganda districts. The study underscores the importance of shifting programmatic interventions from broad, uniform approaches toward more specific, spatially updated strategies that address focal transmission, thereby consolidating gains accomplished and preventing resurgence in high-risk clusters.

**Conflict of interest:** None

**Sources of funding:** The Carter Center, Reaching the Last Mile Fund

**Acknowledgments:** We are grateful to the SSMJ Editorial Team and especially to James Beard for improving the map, to the Uganda Onchocerciasis Reference Laboratory for testing the dried blood spot samples, and to the Magwi County Community Leaders and Director for providing a supportive environment to this survey.

#### References

1. World Health Organization. Onchocerciasis (river blindness) [Internet]. 2025 [cited 2025 Nov 19]. Available from: <https://www.who.int/news-room/fact-sheets/detail/onchocerciasis>
2. Institute for Health Metrics and Evaluation. Global Burden of Disease Study 2017 (GBD 2017) Data Resources | GHDx [Internet]. 2017 [cited 2025 Nov 19]. Available from: <https://ghdx.healthdata.org/gbd-2017>
3. Ministry of Health, Republic of South Sudan. South Sudan Neglected Tropical Diseases (NTD) Master Plan 2023 - 2027 [Internet]. 2023 [cited 2025 Nov 19]. Available from: <https://espen.afro.who.int/tools-resources/documents/south-sudan-ntd-master-plan-2023-2027-en>
4. Lont YL, Coffeng LE, Vlas SJ de, Golden A, Santos T de los, Domingo GJ, et al. Modelling Anti-Ov16 IgG4 Antibody Prevalence as an Indicator for Evaluation and Decision Making in Onchocerciasis Elimination Programmes. PLoS Negl Trop Dis. 2017;11(1):e0005314. doi: <https://doi.org/10.1371/journal.pntd.0005314>
5. World Health Organization. Onchocerciasis elimination mapping [Internet]. World Health Organization; 2025 [cited 2026 Mar 23]. Available from: <https://www.who.int/publications/b/67137>
6. World Health Organization. Onchocerciasis [Internet]. 2022 [cited 2026 Feb 23]. Available from: <https://www.who.int/news-room/fact-sheets/detail/onchocerciasis>

7. Tekle AH, Zouré HGM, Noma M, Boussinesq M, Coffeng LE, Stolk WA, et al. Progress towards onchocerciasis elimination in the participating countries of the African Programme for Onchocerciasis Control: epidemiological evaluation results. *Infect Dis Poverty*. 2016;5(1):66. doi: <https://doi.org/10.1186/s40249-016-0160-7>
8. Borenstein M, Hedges LV, Higgins JP, Rothstein HR. *Introduction to meta-analysis*. John Wiley & sons; 2021.
9. Amazigo U, Boatin B. The future of onchocerciasis control in Africa. *The Lancet*. 2006;368(9551):1946–7. doi: [https://doi.org/10.1016/S0140-6736\(06\)69786-9](https://doi.org/10.1016/S0140-6736(06)69786-9) PubMed PMID: 17141687.
10. Chihoma E, Mbwambo SG, Moirana E, Masayi C, Masanika J, Lalji S, et al. Onchocerciasis Elimination Mapping in Tanzania Mainland: A case of Kilindi DC, Handeni DC and Kilolo DC, Iringa DC. *medRxiv*. 2024;2024–03.
11. Cupp EW, Sauerbrey M, Richards F. Elimination of human onchocerciasis: history of progress and current feasibility using ivermectin (Mectizan®) monotherapy. *Acta Trop*. 2011;120:S100–8.
12. Diawara L, Traoré MO, Badji A, Bissan Y, Doumbia K, Goita SF, et al. Feasibility of onchocerciasis elimination with ivermectin treatment in endemic foci in Africa: first evidence from studies in Mali and Senegal. *PLoS Negl Trop Dis*. 2009;3(7):e497.
13. Monod M, Brizzi A, Galiwango RM, Ssekubugu R, Chen Y, Xi X, et al. Longitudinal population-level HIV epidemiologic and genomic surveillance highlights growing gender disparity of HIV transmission in Uganda. *Nat Microbiol*. 2024;9(1):35–54. doi: <https://doi.org/10.1038/s41564-023-01530-8>
14. Lita EP, Ochi EB, Misinzo G, Heerden H van, Katani R, Godfroid J, et al. Seroprevalence and risk factors of brucellosis in pastoralists and their livestock in Central Equatoria State, South Sudan. *PLoS Negl Trop Dis*. 2024;18(12):e0012144. doi: <https://doi.org/10.1371/journal.pntd.0012144>
15. Pang X, Ma J, He Q, Tang K, Huang J, Fang N, et al. Analysis of HIV transmission characteristics and intervention effects in Guangxi based on molecular networks. *AIDS*. 2025;39(6):719. doi: <https://doi.org/10.1097/QAD.0000000000004123>
16. Hu Z, Liu Y, Wang J, Meng Z, Leuba SI, Wei J, et al. Frequently Transmission and Close Relationship Among Immigrants in the China–Myanmar Border Region Indicated by Molecular Transmission Analysis From a Cross-Sectional Data. *Front Med*. 2022;8. doi: <https://doi.org/10.3389/fmed.2021.693915>
17. Fodjo JS, Ngarka L, Njamnshi W, Enyong P, Bissek ACZK, Njamnshi A. Onchocerciasis in the Ntui Health District of Cameroon: epidemiological, entomological and parasitological findings in relation to elimination prospects. *Parasit Vectors*. 2022;15. doi: <https://doi.org/10.1186/s13071-022-05585-0>
18. Hendy A, Krit M, Pfarr K, Laemmer C, De Witte J, Nwane P, et al. Onchocerca volvulus transmission in the Mbam valley of Cameroon following 16 years of annual community-directed treatment with ivermectin, and the description of a new cytotype of Simulium squamosum. *Parasit Vectors*. 2021;14. doi: <https://doi.org/10.1186/s13071-021-05072-y>
19. Chikezie F, Opara K, Ubulom P, Yaro CA, Al-Akeel R, Osei-Atweneboana M, et al. Onchocerciasis transmission status in some endemic communities of Cross River State, Nigeria after two decades of mass drug administration with ivermectin. *Sci Rep*. 2023;13. doi: <https://doi.org/10.1038/s41598-023-31446-6>
20. Yosef DK, Awil BS, Ismail A, Hassan MA. Epidemiological assessment of onchocerciasis in previously unmapped villages for ivermectin distribution in Fer-Fer District, Somali Regional State, Eastern Ethiopia. *BMC Infect Dis*. 2025;25. doi: <https://doi.org/10.1186/s12879-025-11164-2>

# Prevalence of urinary schistosomiasis in communities along the River Niger Basin areas, North-Central Nigeria

Nyamngee Amase<sup>1</sup>, Ikpe Raphael Terlumun<sup>1</sup>, Kayode Yusuf Abel<sup>1</sup>, Sulaiman Mariam Kehinde<sup>1</sup>, Akanbi Il Aliu Ajibola<sup>2</sup>

## Author Affiliation:

1. Department of Medical Microbiology and Parasitology, Faculty of Basic Clinical Science, College of Health Sciences, University of Ilorin, Nigeria.
2. Department of Microbiology, University of Ilorin Teaching Hospital, Ilorin, Nigeria.

## Correspondence:

Nyamngee Amase  
[anyamngee2010@yahoo.com](mailto:anyamngee2010@yahoo.com)

**Submitted:** March 2026

**Accepted:** April 2026

**Published:** May 2026

**Citation:** Amase et al. Prevalence of urinary schistosomiasis in communities along the River Niger Basin areas, North-Central Nigeria. South Sudan Medical Journal, 2026;19(2):87-92 © 2026 The Author (s) **License:** This is an open access article under [CC BY-NC](https://creativecommons.org/licenses/by-nc/4.0/) DOI: <https://dx.doi.org/10.4314/ssmj.v19i2.4>

## ABSTRACT

**Introduction:** Urinary schistosomiasis is a chronic, progressive disease caused by the parasitic bloodworms (trematodes) *Schistosoma haematobium*. Following infection, the adult worm develops in humans and resides in the veins supplying the bladder, leading to haematuria. Globally, there are approximately 240 million infected individuals with an estimated 534,000 fatalities annually. Urinary schistosomiasis has a higher prevalence in remote areas of Nigeria. This study aimed to determine the prevalence and intensity of urinary schistosomiasis in the riverine communities along the River Niger Basin Areas in North-Central Nigeria.

**Method:** Urine samples were collected from 3,475 people aged between 10 and 55 years from five communities along the basin areas of the River Niger and analysed using microscopy to detect the presence of *Schistosoma haematobium* eggs.

**Results:** The prevalence of urinary schistosomiasis was high (72.66%), and more males were significantly infected ( $p < 0.05$ ). The egg counts revealed moderate-to-heavy infections, with farmers and miners having a significantly higher prevalence than other occupations. This infection affects all ages, peaking between ages 25 and 29.

**Conclusion:** The prevalence of urinary schistosomiasis in this study is high, predominantly among farmers and gold miners, affecting all ages, primarily due to climate change and gold mining activities among participants. Therefore, there is a need for targeted interventions, including health education about schistosomiasis and mass drug administration to mitigate the burden of the disease in such communities across Nigeria and elsewhere.

**Keywords:** prevalence, intensity, schistosomiasis, riverine communities, Nigeria

## Introduction

Schistosomiasis is a chronic, progressive disease caused by the parasitic bloodworms (trematodes). Urinary schistosomiasis is caused by *Schistosoma haematobium*, transmitted from one person to another when an infected person passes the ova

from urine into water bodies. The ova hatch immediately into miracidia, which are then ingested by a snail (in this case, *Bulinus spp*). Inside the snail intermediate host, the miracidia undergo metamorphosis into cercariae, which are then released and become free-swimming in the water, serving as the infective stage to man. The cercariae penetrate humans through the skin and the infective cercariae metamorphose and become adult worms in the human host and move to the predestined site, where, in this case, they reside in the veins supplying the bladder. There they continue to lay eggs, and these eggs are released into the bladder and do pass out with the urine into a water body, hence the life cycle continues.<sup>[1]</sup> The main presenting symptom of urinary schistosomiasis is haematuria.

Schistosomiasis poses a public health challenge second only to malaria globally. The prevalence of schistosomiasis infections ranks second among the major Neglected Tropical Diseases linked to poverty.<sup>[2]</sup> Globally, there are approximately 240 million infected individuals with an estimated 534,000 fatalities annually.<sup>[3]</sup> Although not causing early death, this disease has a severe, debilitating effect on individuals.<sup>[4]</sup> As in other African countries, urinary schistosomiasis has a higher prevalence in remote areas of Nigeria.<sup>[5]</sup> The prevalence within communities is linked to the presence of water reservoirs such as dams, rivers, or streams, which are crucial for meeting the domestic and recreational water needs of the local population.<sup>[6]</sup> The cercariae directly penetrate the skin of individuals engaged in water-related activities such as fishing, rice or other fadama (dry season irrigation) farming, gold washing (mining process), laundry, bathing, and swimming.<sup>[7]</sup> Individuals at the highest risk are those residing in or visiting areas where the disease is widespread.<sup>[8]</sup> In sub-Saharan Africa, more than 90% of schistosomiasis cases occur in Nigeria, with an estimated 29 million cases reported.<sup>[9]</sup> Data published in 2021 indicate approximately 29 million infected individuals, including 16 million children and young adults, with 101 million at risk in the country.<sup>[9]</sup> Given the identification of urinary schistosomiasis across various regions in Nigeria, with estimates of intensity (i.e., the number of eggs in urine samples) and prevalence rates, coupled with indications of increasing incidence,<sup>[10]</sup> the absence of current estimates underscores the need for updated information.

Nigerian school children and active young adults continue to face significant morbidity from schistosomiasis.<sup>[11]</sup> The highest infection intensities are typically observed in children aged 8 to 20 years in endemic areas but not

restricted to these age groups in such endemic areas. Localized studies, such as those in Kwara State, have consistently reported high prevalence and intensity with no particular age pattern.<sup>[11]</sup> This study aimed to determine the prevalence and intensity of urinary schistosomiasis in riverine communities along the River Niger Basin Areas in North-central Nigeria.

### Method

This study was carried out in riverine communities along the River Niger Basin Areas through Kwara and Niger states in North-central Nigeria. It lies within latitude 8°30'00"N and longitude 4°35'00"E, and the inhabitants in the study area are mainly civil servants, farmers, fishermen, traders, artisans, and individuals from the organized private sector.

This cross-sectional study included schools and communities and used a quantitative data collection method to assess potential predisposing risk factors. Individuals aged 10 to 55 years were randomly selected using the open research balloting system (to maintain objectivity and reduce bias in subject selection).

In addition, conditions, characteristics, outcomes, and exposure of pregnant women to malaria were studied.

The sample size was determined as described by Abdulkareem et al.<sup>[4]</sup>

Ethical approval was obtained from the Ethical Review Committees of the Kwara State and Niger State Ministries of Health.

### Data collection

Semi-structured and standardized proforma were administered to participants with assistance from research staff. This proforma aimed to collect demographic data, including names, ages, genders, and educational backgrounds, prior to sample collection.

### Sample collection and transportation

Sample collection was conducted between 10:00 am and 2:00 pm, a period when *Schistosoma haematobium* egg excretion peaks. Participants were then provided with disposable plastic cups labelled with a code and serial number. Hand soap and sanitizers were provided for all participants to disinfect their hands before and after sample collection. From the urine specimens, 10 ml was transferred into labelled universal bottles using a syringe. The samples were preserved by adding 10 ml of 70% alcohol and carefully stored in sturdy paper cartons,

secured with adhesive tape to prevent spills or displacement during transport to the laboratory.

### Laboratory analysis and interpretation of results

#### *Microscopic examination procedure for the urine filtration technique*

The preserved urine samples were examined using the filtration technique with a 325 µm wire mesh. The samples were passed through the wire mesh filter, folded conically, and placed in small plastic funnels atop empty beakers or flasks. The sediment was transferred to a cross-hatched Petri dish and examined under a dissecting microscope at 20x magnification to identify characteristic *Schistosoma haematobium* eggs. Eggs were counted using a hand tally counter, and infection intensity was expressed as eggs per

10 ml of urine. The number of eggs per 10 ml of urine was counted and graded as follows: light (50-149), moderate (150- 499), and heavy (≥500). For quality control, precision, and accuracy, egg counts were performed twice. Only ova with a conspicuous terminal spine were counted.

### Data analysis

All data were analysed using SPSS version 27.00, with descriptive statistics used to determine associations between observed variables, and conclusions were drawn at the 95% confidence level.

### Results

This study examined 3,475 active young and old people in five communities along the River Niger in North-

**Table 1. Prevalence of schistosomiasis by age, sex and other demographic variables in the study areas**

| Parameters         | Indicators               | No. Examined | Positive n (% of 3,475) | p-Value |
|--------------------|--------------------------|--------------|-------------------------|---------|
| Age (years)        | 10-14                    | 222          | 210 (6.04)              | p>0.05  |
|                    | 15-19                    | 425          | 392 (11.28)             |         |
|                    | 20-24                    | 617          | 486 (13.99)             |         |
|                    | 25-29                    | 624          | 501 (14.42)             |         |
|                    | 30-34                    | 611          | 422 (12.14)             |         |
|                    | 35-39                    | 398          | 210 (6.04)              |         |
|                    | 40-44                    | 308          | 196 (5.64)              |         |
|                    | 45-49                    | 200          | 82 (2.36)               |         |
|                    | ≥50                      | 70           | 26 (0.75)               |         |
|                    | <b>Total</b>             | <b>3,475</b> | <b>2,525 (72.66)</b>    |         |
| Sex                | Male                     | 2,611        | 2003 (57.64)            | p<0.05  |
|                    | Female                   | 864          | 522 (15.02)             |         |
|                    | <b>Total</b>             | <b>3,475</b> | <b>2,525 (72.66)</b>    |         |
| Educational Levels | Non-formal               | 428          | 264 (7.60)              | p<0.05  |
|                    | Primary                  | 921          | 620 (17.84)             |         |
|                    | Secondary                | 1,281        | 1,016 (29.24)           |         |
|                    | Tertiary                 | 845          | 625 (17.98)             |         |
|                    | <b>Total</b>             | <b>3,475</b> | <b>2,525 (72.66)</b>    |         |
| Occupation         | Trading                  | 312          | 122 (4.83)              | p<0.05  |
|                    | Farming                  | 1,212        | 980(38.81)              |         |
|                    | Fishing                  | 423          | 386(15.29)              |         |
|                    | Mining                   | 420          | 411 (16.28)             |         |
|                    | Civil Service            | 416          | 201 (7.96)              |         |
|                    | No particular occupation | 692          | 425 (16.83)             |         |
|                    | <b>Total</b>             | <b>3,475</b> | <b>2,525 (72.66)</b>    |         |

**Table 2. Intensity of infection by egg count among participants across communities in the study areas**

| Intensity   |                   |                            |                             |                          | ANNOVA<br>(p-value) |
|-------------|-------------------|----------------------------|-----------------------------|--------------------------|---------------------|
| Level       |                   | Light                      | Moderate                    | Heavy                    |                     |
| Score range | Infected<br>(+ve) | 1-149 eggs/<br>10 mL urine | 150-499 eggs/<br>10mL urine | ≥500 eggs/<br>10mL urine |                     |
| Community A | 503 (19.92)       | 184(7.29)                  | 137(5.43)                   | 182(7.21)                | 0.01196             |
| Community B | 506 (20.04)       | 167(6.61)                  | 167(6.61)                   | 172(6.81)                |                     |
| Community C | 504(19.96)        | 161(6.38)                  | 171(6.77)                   | 172(6.81)                |                     |
| Community D | 504 (19.96)       | 101(4.00)                  | 199(7.88)                   | 204(8.08)                |                     |
| Community E | 508 (20.12)       | 198(7.84)                  | 166(6.57)                   | 144(5.70)                |                     |
| Total       | 2,525 (100)       | 811 (32.12)                | 840(33.27)                  | 874(34.61)               |                     |

central Nigeria, and 2,525 (72.66%) tested positive for schistosomiasis, as shown in Table 1.

The infection rate peaked in the 25-29 age group (14.42%), followed closely by the 20-24 (13.99%) and 30-34 (12.14%) age groups. The age group with the lowest prevalence of infection was ≥50 years (0.75%). However, the differences in the prevalence across age groups were not statistically significant. Conversely, more males (57.64%) were infected than females (15.02%) ( $P<0.05$ ). The prevalence by participants' educational level showed that individuals with secondary education had the highest prevalence (29.24%), followed closely by those with tertiary (17.98%) and primary (17.84%) education, with the lowest prevalence among those with non-formal education (7.60%).

Again, in Table 1, the difference in prevalence by participants' occupation was statistically significant, with the highest prevalence of 38.81% among farmers, 16.28% among miners, and 15.29% among fishermen, and the lowest (3.51%) among traders ( $P<0.05$ ).

In all five communities where the study was conducted, the prevalence of infection was relatively similar, ranging from 5.43% to 7.84%. The small differences observed among the communities were not statistically significant ( $p>0.05$ ). Also, there was an even distribution of infection intensity across the study areas (Table 2).

Figure 1 shows activities predisposing community members to schistosomiasis infection in the study areas.

In Figure 2, the numbers from 0 to 5 indicate the range of urine colours and haematuria levels across different areas of the study.

## Discussion

### Prevalence of schistosomiasis

The study finds that, proportionately, individuals in the communities have a significant schistosomiasis infection rate. The prevalence of urinary schistosomiasis among community inhabitants in the riverine areas was high, at 72.66%. This aligns with the World Health Organization's classification of the area as heavily endemic, highlighting the urgent need for intensified control measures.<sup>[3]</sup> This suggests that environmental and socio-economic factors significantly influence the transmission of urinary schistosomiasis in these communities, as previously reported,<sup>[5]</sup> emphasising that children and young adults living near rivers and lakes are more susceptible to infection. Joseph et al.<sup>[6]</sup> reported a prevalence rate of 8.9%, which is lower but still comparable in their study area.

This high prevalence rate from this study compares with previous findings in Kwara State, where Abdulkareem et al.<sup>[4]</sup> recorded a high prevalence of 45.6% and Bolaji<sup>[7]</sup> recorded 58.7% in similar communities, attributing the higher rate to the presence of extensive irrigation projects, water reservoirs, and frequent river contact, which significantly impact local socio-economic practices.

### Prevalence by age groups

Although prevalence rates varied across age groups, the differences were not significant. The higher rate of infection seen among the 25-29 age group may be explained by this group being the most active in the study population,




Figure 1. Activities of participants in infected water that favour the spread of schistosomiasis (Credit: Amase Nyamngee)




Figure 2. The appearances of different urine samples collected from participants

compared with the lower prevalence observed in the  $\geq 50$  age group, who apparently have reduced water-contact activities. Similarly, findings from Muhammad et al.<sup>[8]</sup> recorded a high prevalence of 62.86%, and Obonogbigho et al.<sup>[9]</sup> also recorded a prevalence of 19.3% among similar age groups. According to them, the high prevalence was attributed to more frequent engagement in water-related activities such as swimming, mining, and playing in rivers.

### Prevalence by sex

This study found that more males (57.64%) were infected than females (15.02%), and this difference was statistically significant. These data indicate that social stigma associated with open bathing may hinder females from actively participating in high-water-contact activities. The higher prevalence among males aligns with findings from Abdullahi & Ramatu,<sup>[10]</sup> who reported 38.2% among males and 15.0% among females. However, the prevalence observed among females may be due to indiscriminate farming activities in these communities that involve both males and females.

### Quantification of eggs by schools in the communities

The quantification of *Schistosoma* eggs in urine samples provides critical insight into infection intensity among participants in these communities. This distribution of egg intensity was categorized as light, moderate, or heavy, with most infections falling into the heavy category. This high intensity in our findings is attributable to a lack of understanding of the biology of schistosomiasis transmission and of water-contact activities as a means of livelihood in these communities, which predisposes community members to repeated infection, resulting in a heavy egg load in their urine. The high egg-count intensity

observed in our findings indicates that the transmission of infection to others, or even reinfection within the same group, is more likely, posing a severe risk of schistosomiasis spread in these communities.

### Conclusion

There is a very high prevalence of urinary schistosomiasis among community inhabitants in riverine areas along the river Niger in North-Central Nigeria. The intensity of this infection is also very high, affecting people of all ages, both males and females. We recommend targeted interventions to reduce water contact, improve sanitation, and introduce health education.

**Conflict of interest:** none

**Authors' contribution:** AN: Conception and design of the study, analysis and interpretation of data; drafting of the article. MJS: Literature search and assemblage of data. IRT: study design, field work, laboratory work. KYA: Provision, Collection, and assemblage of data. SMK: Literature search, Provision, Collection, and assemblage of data. AAA: Study design, revising the write-up, and administrative, technical, and logistic support.

**Acknowledgments:** We acknowledge the authorities of the University of Ilorin Teaching Hospital for allowing us access to their Laboratory.

**Study Limitation:** This was a community-based study; thus, there is a possibility of community overlap. Several of the participants did not know their exact age. However, we implemented a calendar of events to overcome this challenge.

**Funding:** None from any institution. All forms and

sources of assistance were personal contributions from the researchers.

### References

1. Vennervald B. Schistosomiasis and Other Trematode Infections. In: Loscalzo J, Fauci A, Kasper D, Hauser S, Longo D, Jameson J, eds. *Harrison's Principles of Internal Medicine*. 21st ed. McGraw-Hill Education; 2022.
2. Folahan FF, Edungbola LE, & Folahan JT. Prevalence of urinary schistosomiasis among primary school pupils. *Journal of Microbiology and Infectious Diseases*. 2021;11(1):95-104. doi: 10.5799/jmid.951609
3. Brunn B, & Aagaard-Hansen J. The social context of schistosomiasis and its control: An introduction and annotated bibliography (1st ed.). Geneva: World Health Organization, 2008.
4. Abdulkareem BO et al. Urogenital Schistosomiasis among Schoolchildren and the Associated Risk Factors in Selected Rural Communities of Kwara State, Nigeria. *Journal of Tropical Medicine*, 2018.
5. Zamdayu NM, Qadeer MA, Chessed G & Francis MI. Prevalence of urinary schistosomiasis among children from three selected local government areas in Adamawa State, Nigeria. *Dutse Journal of Pure and Applied Sciences*, 2023; 9 (3a):168–178.
6. Joseph SO, Abdulkareem BO & Samuel UU. Distribution pattern of human urinary schistosomiasis in Kwara State, Nigeria. *American Journal of Infectious Diseases*, 2017;13(4):38–44.
7. Bolaji OS. Prevalence and intensity of *Schistosoma haematobium* among school children in Ajase-Ipo, Kwara State, Nigeria. *Asian Journal of Biomedical and Pharmaceutical Sciences*, 2015;5(43):06–11. <https://doi.org/10.15272/ajbps.v5i43.685>
8. Muhammad IA, Abdullahi K, Bala AY & Shinkafi SA. Prevalence of urinary schistosomiasis among primary school pupils in Wamakko Local Government, Sokoto State, Nigeria. *The Journal of Basic and Applied Zoology*, 2019; 80(1):7-11.
9. Obonogbigho Boih L, Okaka CE & Igetei JE. A survey of urinary schistosomiasis among school aged children of ten communities in South-South, Nigeria. 2021;57(1):1-13. DOI: 10.4103/0972-9062.308793 <http://www.jvbd.org>
10. Abdullahi M & Ramatu MA. Prevalence of Urinary Schistosomiasis Among Pupils Attending Angwan Lambu Primary School, Keffi, Nasarawa State, Nigeria. *Fudma Journal of Sciences*, 2023; 6(6):146–149. <https://doi.org/10.33003/fjs-2022-0606-1134>
11. Balogun JB, et al. Prevalence and Associated Risk Factors of Urinary Schistosomiasis among Primary School Pupils in the Jidawaand Zobiya Communities of Jigawa State, Nigeria. *Annals of Global Health*, 2022; 88(1):7-11.

## Programme Guidance to Protect the Nutrition of Women and Adolescent Girls in Humanitarian Settings

This guidance has been created for nutrition cluster/sectors, programme staff of UNICEF and the World Food Programme (WFP), and other practitioners who are designing and/or implementing a nutrition response in humanitarian settings. These nutrition responses may include assessments and/or interventions for pregnant and breastfeeding women and adolescent girls in humanitarian settings.

[Watch the webinar recording here.](#)

[https://www.unicef.org/documents/guidance-protect-nutrition-women-and-adolescent-girls-humanitarian-settings?utm\\_campaign=WINS%20%7C%20Maternal%20Nutrition%20ESARO%20%28Partners%29&utm\\_medium=email&utm\\_source=Mailjet](https://www.unicef.org/documents/guidance-protect-nutrition-women-and-adolescent-girls-humanitarian-settings?utm_campaign=WINS%20%7C%20Maternal%20Nutrition%20ESARO%20%28Partners%29&utm_medium=email&utm_source=Mailjet)

# Developing a communicable and infectious diseases research agenda in South Sudan: A multi-stakeholder prioritisation approach

Amanya Jacob<sup>1,2,3</sup>, Ezbon WApary<sup>1,2</sup>, Dhieu Daniel<sup>1,2</sup>, Atemthi Dau<sup>1,2</sup>, Matur T. Tieng<sup>1,2</sup>, John Rumunu<sup>1</sup>, Cicilia Konga<sup>1</sup>, Bol Atem<sup>2</sup>, Betty Eyobo<sup>1</sup>, Robert Napoleon<sup>1</sup>, Joseph Panyuon<sup>1</sup>, Kenneth L Sube<sup>2</sup>, Kon Alier<sup>2</sup>, Joseph Lako<sup>2</sup>, John Suraj<sup>3</sup>, Abuelguasim M. Hassan<sup>3</sup>, Mapuor Areu<sup>2</sup>, Lazarus Atem<sup>2</sup>, Aja Kuol<sup>1</sup>, Pizzi Wilson<sup>1</sup>, Achcer Adup Mou<sup>1</sup>

## Author Affiliation:

1. Department of Research, Survey and Ethics, Ministry of Health, Juba, South Sudan
2. Applied & Industrial Sciences, Schools of Medicine and Public Health, University of Juba, Juba, South Sudan
3. Faculty of Public and Environmental Health, Upper Nile University, Juba, South Sudan

## Correspondence:

Amanya Jacob  
[amanyajazy@yahoo.com](mailto:amanyajazy@yahoo.com)

**Submitted:** February 2026

**Accepted:** April 2026

**Published:** May 2026

**Citation:** Jacob et al. Developing a communicable and infectious diseases research agenda in South Sudan: A multi-stakeholder prioritisation approach. South Sudan Medical Journal, 2026;19(2):93-98 © 2026 The Author (s) **License:** This is an open access article under [CC BY-NC](#) DOI: <https://dx.doi.org/10.4314/ssmj.v19i2.5>

## ABSTRACT

**Introduction:** South Sudan faces a considerable health burden, including high rates of communicable, non-communicable, and maternal and child health concerns within a constrained health system and challenging humanitarian context. Our aim was to engage stakeholders to identify major health research gaps, apply the Africa Centres for Disease Control & Prevention (CDC) prioritization tool to rank disease programme areas and the health system based on context-specific criteria, and produce a ranked agenda to inform implementation plans.

**Method:** A multi-stakeholder, criteria-driven approach was used, adapting the Africa CDC research prioritisation tool. This involved a situational analysis, the identification of potential research questions, the scoring of topics against established criteria such as burden, equity, feasibility, impact, and cost-effectiveness, and a consensus-building workshop. The disease themes and health system components were on high, medium, and low set-scores.

**Results:** Several high-priority research themes and questions were identified. South Sudan remains heavily burdened by infectious diseases, with malaria a leading cause of morbidity and mortality. The highest probability conditions in this category were Human papillomavirus (HPV), respiratory syncytial virus, and staphylococcal infections. Health System Strengthening emerged as a cross-cutting challenge due to workforce shortages, weak infrastructure, and disease surveillance gaps.

**Conclusion:** Initiating the health research agenda with a national research plan that identifies lead institutions, potential funding sources, and timelines was considered essential. Investing in health research capacity strengthening and leveraging on stakeholders such as the Africa CDC is a call to action. Institutionalising and harmonising priority-setting mechanisms through a national health research prioritisation technical committee to ensure ongoing responsiveness to health threats and shocks was also realised as important.

**Keywords:** communicable, research agenda, Africa CDC prioritisation tool, South Sudan

### Introduction

An effective health research agenda is fundamental for aligning research investments with national health priorities, particularly in developing and fragile contexts such as South Sudan. Research prioritisation informs the allocation of resources to health needs, supports the generation of appropriate evidence, and facilitates strategic coordination among stakeholders.<sup>[1]</sup> South Sudan faces acute and chronic burdens of communicable and non-communicable diseases. This report documents the process, prioritised health research themes and questions, and indicates the next priority steps in South Sudan, adapting, applying, and contextualising the Africa CDC research prioritisation tool.<sup>[2]</sup>

South Sudan faces numerous health system challenges: high maternal and newborn mortality, a scarcity of skilled health workers, disease outbreaks - for instance, cholera, Hepatitis E, and Mpox. There are only 7.6 skilled health workers per 10,000 population, when the minimum for universal health coverage (UHC) is 44 per 10,000 persons.<sup>[3]</sup> Priorities to achieve UHC include strengthening health security, improving population well-being, and reinforcing health information systems.<sup>[4]</sup>

These essential pillars necessitate targeted research to inform decision-making and interventions. Priority-setting processes are heavily influenced by external donor-driven agendas. Power asymmetries between donors and national health governance exist, affecting how priorities were set within the Health Pooled Fund (HPF models 1, 2, & 3) service-delivery contract framework, as well as the Health Sector Transformation Project (HSTP), a multi-donor funded project.<sup>[5]</sup>

Health financing and implementation gaps persist due to weak priority-setting mechanisms and governance.<sup>[6]</sup> A structured activity to prioritise health research in South Sudan is critical to ensure alignment of research with national priorities. The principal objectives were to engage key stakeholders to identify major health research priorities and apply and contextualise the Africa CDC prioritisation tool to rank research topics according to context-specific criteria, produce a ranked health research agenda that inform national research funding, partnerships, and implementation plans.

### Method

The stakeholders from the Ministry of Health (MoH), academia, partners' agencies, national research

institutions, civil society organisations and researchers, and health service delivery partners were brought together for a consultative workshop held 1-5 September 2025 in Juba, with 35 invited, 53 attended. Multidisciplinary experts from biomedical, public health, laboratory, epidemiology, biostatisticians, monitoring and evaluation, ethics and clinical fields to (a) review health systems and research needs in South Sudan; (b) apply clear and agreed criteria to prioritise research themes and questions; (c) agree on a ranked list of priority research topics; and (d) propose mechanisms for follow-up, including funding coordination, capacity-building, and knowledge translation. This aligns with the Africa CDC's emphasis on inclusive, multi-stakeholder consultative approaches.<sup>[7]</sup>

The Africa CDC Research prioritisation tool was adapted, with a framework developed from a discrete choice experiment, a conditional utility model, and weighted criteria to guide our research agenda. The tool was used to conduct a situational analysis of the research landscape and burden of disease; identify candidate research questions; apply layout criteria and a weighted scoring and ranking scheme for themes; and validate and build consensus.<sup>[7]</sup> Stakeholders were consulted, and invitations extended to MoH technical staff, implementing partners, academia, and researchers. A desk analysis was made of strategic documents and previous research to identify gaps and weaknesses. The participants agreed on prioritisation criteria and contextualized the Africa CDC research prioritisation tool.

Each proposed research topic was scored and ranked against the criteria. Topics were ranked by aggregate probability scores, then discussed to reach consensus. During the workshop, participants reviewed existing national health indicators, national databased (DHIS2), recent studies, and stakeholder inputs to develop a long list of potential research questions within thematic domains such as communicable diseases, eye defects, fungal infections, injuries, maternal and child health, mental health, neglected tropical, non-communicable, and zoonotic diseases. The list was then scored according to agreed-upon weighting and criteria, using disease thematic groups to score them, and a plenary session was convened to build consensus. Finally, the results were consolidated into a priority ranking.

A plenary session reviewed the ranked list, presented by each thematic area team lead, and proposed mechanisms for monitoring and translating research outputs into policy and practice. The invited health sector stakeholders endorsed the plan for coordinating research

implementation, capacity building for local researchers, and mechanisms for dissemination and uptake. An agreement was reached to consolidate the research priority framework, aligning short-term (1-2 years) and medium-term (3-5 years) research initiatives with the national health strategic plan. The Africa CDC prioritisation tool is not publicly available; its details can be found on their websites and in recent communications.

## Results

### Prioritised research themes and questions

Infectious diseases and outbreak preparedness are important pillars for global health security. Repeated outbreaks expose the vulnerability of public health systems and the need for investment in disease surveillance, prevention, and rapid-response mechanisms. Integrating infectious disease management within the broader health-security framework emphasizes not only biomedical interventions but also governance, coordination, partnerships, and resilience-building. Hence, epidemic preparedness is a key component of international health policy and regulation, demonstrating that the control and threats of infectious conditions are a scientific and strategic imperative for safeguarding population health.




Figure 1. Situational context in Continental Africa: Diseases by thematic area




Figure 2. Situational context in South Sudan: Diseases by thematic area

### Context in Continental Africa

Figure 1 shows Communicable and Infectious Diseases (CID) 120 and emphasises that, in Africa, infectious disease outbreaks remain substantial. For example, from July 2021 to June 2022, 112 of 130 acute infectious disease conditions reported in Africa were infectious disease outbreaks. The non-communicable disease (NCD) burden is rising, and so a “double burden” of infectious and non-infectious conditions demands attention.

### Context in South Sudan

Health research in South Sudan is limited by low capacity and the translation of evidence into policy and practice. Many member states still under-invest in research.

The disease burden is complex, reflecting longstanding communicable and maternal/child health challenges, emerging non-communicable, zoonotic, injury, ocular, and mental-health conditions. Figure 2 provides the thematic breakdown of disease burden by proportion, starting from communicable and infectious diseases (70%), maternal and child health (47%), non-communicable (32%), neglected tropical diseases (27%), injuries (21%), eye defects (19%), zoonotic diseases (15%), mental health disorders (11%), and fungal infections (4).

Infectious diseases are a heavy burden in South Sudan. Malaria is a leading cause of morbidity and mortality. In 2021, it was estimated at ~3 million cases (incidence ~280 per 1,000) with ~7,344 deaths. In addition, there is a high incidence of HIV/AIDS, lower respiratory infections and diarrhoeal diseases. Thus, the figure “70” in Figure 2 shows the dominance of infectious-diseases. The latter is exacerbated by humanitarian crises, displacement, weak infrastructure, and outbreak risk.

The maternal, neonatal, and child health theme is an area of concern. The link between infections, poor immunisation coverage, malnutrition and maternal/child outcomes is clear in South Sudan. Poor vaccination coverage and disruption of services are major public health issues. The figure “47” in the Figure 2 reflects these matters.

NCDs are increasingly recognised in South Sudan, accounting for 28% of deaths in 2019. The probability of premature death from NCDs for ages 30-70 years was ~20% in 2020, higher than the sub-Saharan African average of ~13.7%. The figure “32” in Figure 2 signals the growing relevance of NCDs. There is limited surveillance of NCDs with low diagnostic capacity, shortages of essential medicines, and health information systems that focus on acute infections and maternal and child conditions. Thus,

while the number (“32”) appears lower than infectious burdens, the impact of NCDs is under-estimated.

South Sudan has a challenging problem of neglected tropical diseases (NTDs). “It is endemic for all five NTDs amenable to preventive chemotherapy through mass drug administration (MDA) in 2021, reached 91% of the 6.6 million people targeted for treatment.”<sup>[8]</sup> Number “27” in Figure 2 highlights the impact of NTDs.

In a country with decades of conflict and instability, injuries are a notable burden (“21” in Figure 2). Data on national injuries are sparse and represent a key area for public-health programming.

Ocular disorders receive poor attention in national health statistics but are considerable for disability-adjusted life years (DALYs). In South Sudan, eye defects are recognised as a substantial thematic area (see “19” in Figure 2). The broader disease burden, including sensory impairments and congenital anomalies, is important for integrated primary care and rehabilitative service planning.

Zoonotic diseases are an under-resourced area, given the livestock-based livelihoods, displacement, and outbreak risk. Figure “15” (Figure 2) indicates that zoonotic diseases require serious attention.

Although the number “11” (Figure 2) is relatively low, and probably an underestimation, mental health disorders are serious in the context of chronic and acute conflict, trauma, and limited-service availability. Specialised mental-health clinics are rare, aggravating the unmet need.

Fungal infections are frequently overlooked, particularly in settings like South Sudan. Figure “4” in Figure 2 is likely another underestimate, bearing in mind immunosuppression, malnutrition, and poor hygiene that predispose to fungal infections.

The coexistence of major communicable disease burden (“70”) with an increasing NCD burden (“32”) reflects the “double burden” of disease in low-income, fragile settings. The health system must adapt to address these changing circumstances.

Many thematic categories are under-reported. Health-information systems focus on infectious, maternal, and reproductive health, with chronic conditions under-recognised and underestimated and lack integration in healthcare provision.

The “neglected” categories (NTDs “27”, eye defects “19”, zoonoses “15”, fungal infections “4”) need a cross-cutting approach such as “One-Health” (a multi-professional body

addressing zoonotic diseases). In this context improvement in cost-effective laboratory capacity is crucial.

In the South Sudan context, preventable infectious and




Figure 3. Diseases and health conditions in South Sudan context




Figure 4. Priority communicable and infectious diseases




Figure 5. Priority fungal infectious diseases & conditions

maternal–child conditions, particularly pneumonia, diarrhoeal disease, vaccine-preventable bacterial infections, and the combination of malnutrition and maternal complications, represent the highest health threats in 2020-2025 (EWARS/DHIS2 report). These threats are compounded by NTDs and episodic viral haemorrhagic disease risks, all amplified by conflict, displacement, food insecurity and weak health systems.

Pneumonia causes more under-5-year-old deaths than any other disease. High malnutrition and low vaccine coverage are serious issues in South Sudan. In addition, there is a high maternal mortality and low service coverage. Hypertensive disorders in pregnancy contribute substantially to maternal deaths. Addressing maternal health and nutrition is central to reducing maternal and child mortality.

Conditions such as cysticercosis and occasional human cases of dracunculiasis place NTD control on the priority list. A small number of human guinea-worm cases were documented in the early 2020s. The emergence of viral haemorrhagic fevers highlights ongoing outbreak risk and the need for cross-border preparedness.

### Priority research questions

#### *Communicable and infectious diseases (CID)*

The bar chart presents a comparative analysis of various diseases and conditions according to their associated probabilities, expressed as percentages (Figure 4). The probabilities range from 41.3% to 94.2%, with the diseases grouped by colour to indicate relative priority or likelihood.

At the highest probability level (red), Human papillomavirus (HPV) ranks first with 94.2%, followed by Respiratory syncytial virus and Staphylococcal infections, both at 91.2%. Other diseases include pneumococcal meningitis, Hepatitis B, Herpes simplex 2, and botulism, all exceeding 84%.

The intermediate probability group (yellow) includes measles (79.4%), Zika (69.4%), and diphtheria, reflecting moderate probability/priority. The lower probability group (green), includes diseases such as gonorrhoea, SARS, and impetigo (each at 62.3%), as well as Marburg (56.6%), HIV/AIDS (55.2%), rubella (47.3%), trichomoniasis (41.3%), Mpox, and tetanus.

Figure 5 presents four distinct medical conditions with similar probability percentages: Madura foot, *Tinea corporis* (ringworm), onychomycosis, and candidiasis.

## Discussion

### Broad and specific strategic research questions for CIDs

What is known about the distribution and determinants of communicable and infectious diseases in South Sudan?

The detailed, specific strategic research questions for different thematic areas are found in the health/diseases research prioritization agenda for 2025-2030. These research questions are categorized as short-, medium-, and long-term. The short-term questions include, but are not limited to, the following: (1) What are the structural and systemic factors influencing South Sudan's emergency preparedness and response capacity at national and subnational levels? (2) How do governance, financing, and coordination mechanisms influence the effectiveness of South Sudan's emergency health preparedness system? (3) What lessons can be drawn from South Sudan's responses to past public health emergencies (e.g., cholera, Ebola alerts, yellow fever, Mpox, Hepatitis-E virus, COVID-19) to strengthen resilience against future outbreaks? (4) How do humanitarian crises, displacement, and conflict dynamics shape emergency preparedness planning and implementation in South Sudan? For medium and long-term strategic research questions, four are selected, with two questions from each category. For example: (5) What are the cost-effectiveness outcomes of universal Hepatitis B vaccination programmes in South Sudan? (6) How does climate change influence the seasonal distribution of typhoid outbreaks? (7) What are the barriers to effective implementation of oral cholera vaccines in high-risk populations? (8) What is the role of antimicrobial resistance in the management of cholera cases?

### Reflections on the prioritisation process

Through engaging stakeholders, broad participation ensured that multiple perspectives and opinions informed the agenda. However, some sub-national groups were under-represented. Feasibility and sustainability were prominent concerns. Participants emphasised that research should be adaptable to changing circumstances. It is acknowledged that donor funding and technical capacity drive research priorities.<sup>[5]</sup> The process uses explicit criteria and ranking as well as resources from Africa CDC, but the country's sustainability and ownership remained a challenge.

Aligning research priorities with national health policy and the health sector strategic plan ensures relevance.<sup>[4]</sup>

A recurrent theme was investment in national research capacity, data systems, and institutional infrastructure. The exercise confirms the applicability of the Africa CDC prioritisation approach in a developing context. The multi-stakeholder and criteria-driven process lends transparency to the setting of research priorities. Translating prioritized research into policy and practice requires institutional linkages, which should be deliberately constructed. A similar approach used in developing newborn and child health guidelines in African countries highlighted the importance of context-specific stakeholder engagement and consensus processes.<sup>[9]</sup>

### Conclusion

The prioritisation activity represents a meaningful step toward systematically aligning health research with national health priorities in South Sudan. Hence, engaging diverse stakeholders, applying transparent criteria, and producing a ranked research agenda linked to policy documents, the groundwork has been laid for more strategic research investment. Nonetheless, the success of the agenda depends on sustained governance, capacity investment, funding alignment and translation of research into actionable health policies and programmes.

### Recommendations

Operationalise the prioritized health research agenda by developing a national research plan identifying lead institutions, funding sources, timelines and responsible parties for each high-priority topic.

Invest in research capacity strengthened by leveraging relevant stakeholders, such as the Africa CDC's capacity-mapping initiatives, to build infrastructure, train researchers, strengthen ethical/regulatory frameworks, and embed research within service delivery.

Institutionalise priority-setting mechanisms by establishing a standing national research prioritisation committee that uses the Africa CDC tool to review and update across cycles, ensuring responsiveness to emerging health threats.

Strengthen translation of research into policy by creating pathways for uptake of research findings by the MoH and partner agencies, including policy briefs, knowledge-translation workshops, and institutional linkages.

Monitor and evaluate the impact of prioritised research by developing indicators to track funding flows, research outputs, policy uptake, and health outcomes tied to each priority topic.

**Conflict of interest:** Nil

**Funding:** Nil.

### References

1. Widdig H, Tromp N, Lutwama GW, Jacobs E. The political economy of priority-setting for health in South Sudan: a case study of the Health Pooled Fund. *Int J Equity Health* 2022;21(1):68.
2. Africa CDC. Risk Ranking and Prioritization of Epidemic-Prone Diseases [Internet]. Afr. CDC2023 [cited 2025 Dec 8]; Available from: <https://africacdc.org/download/risk-ranking-and-prioritization-of-epidemic-prone-diseases/>
3. Loi G. Review Article: Scaling up human resources for health in South Sudan: A strategic imperative for achieving universal health coverage. *South Sudan Med J* 2025;18(3):124–6.
4. World Health Organization. Country Cooperation Strategy 2022–2026 South Sudan. WHO. Regional Office for Africa [Internet]. 2022 [cited 2025 Dec 8]; Available from: <https://www.afro.who.int/countries/south-sudan/publication/country-cooperation-strategy-2022-2026-south-sudan>
5. Widdig H, Tromp N, Lutwama GW, Jacobs E. The political economy of priority-setting for health in South Sudan: a case study of the Health Pooled Fund. *Int J Equity Health* 2022;21(1):68.
6. Alier K, Kozuki N, Mothupi M. Stakeholder perspectives of maternal and newborn health prioritization in South Sudan: a policy analysis. *Discov Public Health* 2025;22(1):5.
7. The African Health Research Agenda Should Focus on Delivering More Health Outcomes [Internet]. Afr. CDC [cited 2025 Dec 8].
8. Sanders AM, Stewart AEP, Makoy S, Chebet JJ, Magok P, Kuol A, et al. Burden of trachoma in five counties of Eastern Equatoria state, South Sudan: Results from population-based surveys. *PLoS Negl Trop Dis* 2017;11(6):e0005658.
9. Durão S, Effa E, Mbeye N, Mthethwa M, McCaul M, Naude C, et al. Using a priority setting exercise to identify priorities for guidelines on newborn and child health in South Africa, Malawi, and Nigeria. *Health Res Policy Syst* 2024;22(1):48.

# Determinants of disaster preparedness among public health facilities in Juba City, South Sudan

Dumba Samuel Kenyi<sup>1</sup>, Muthoni Mwangi Eunice<sup>2</sup>, Musa Olouch<sup>2</sup>

## Author Affiliation:

1. Kenya Methodist University,  
Nairobi, Kenya
2. The Aga Khan University,  
Nairobi, Kenya

## Correspondence:

Dumba Samuel Kenyi  
[dumbasam@gmail.com](mailto:dumbasam@gmail.com)

**Submitted:** December 2025

**Accepted:** April 2026

**Published:** May 2026

**Citation:** Kenyi et al. Determinants of preparedness for disasters among public health facilities in Juba City, South Sudan. South Sudan Medical Journal, 2026;19(2):99-106 © 2026 The Author (s) **License:** This is an open access article under [CC BY-NC](https://creativecommons.org/licenses/by-nc/4.0/) DOI: <https://dx.doi.org/10.4314/ssmj.v19i2.6>

## ABSTRACT

**Introduction:** In times of disaster, the primary duties of the health sector are to save lives, minimize permanent disability, deliver medical care to the injured, and mitigate the risks of disease and mortality from health hazards. The success in delivering these responsibilities relies on the uninterrupted functionality of health services. The study aimed to evaluate the influence of resource availability, health workers' capacity, technological integration, and institutional policy on disaster preparedness in public health facilities in Juba City, South Sudan.

**Method:** This study used a cross-sectional research design. The target population comprised healthcare and administrative workers at three major public health facilities in Juba City. The study used descriptive and inferential statistics to analyse the quantitative data.

**Results:** 148 participants responded to the study questionnaire out of a sample size of 162 respondents. Most respondents 123, (83%) disagreed that their facility had sufficient funding for disaster preparedness activities. Furthermore, 105 (71%) disagreed that specific budgets were allocated to improve disaster preparedness, and 99 (67%) believed that financial constraints significantly impact their facility's ability to prepare for and respond to disasters. A small proportion (71; 48%) reported that staff were regularly trained on the guidelines. The reported insufficient funding for preparedness activities was significantly associated with a facility's inability to initiate a response plan within the recommended timeframe ( $\chi^2=15.24$ ,  $p<0.001$ ).

**Conclusion:** The overall state of readiness is significantly hampered by systemic deficiencies, with the absence of formal institutional policies and severe financial constraints as the most profound challenges. These foundational weaknesses created a cascading effect, leading to inadequate material resources, challenges in maintaining stable, motivated human resources, and a disconnect in general information.

**Keywords:** disaster, health sector, resources, preparedness, Juba City, South Sudan

## Introduction

Globally, health facilities are fundamental components of emergency response plans and must enhance their capabilities and capacities for emergency preparedness well before any disaster.<sup>[1]</sup> In times of disasters, the primary duties of the health sector include saving lives, minimizing permanent disability, delivering medical care to the injured, and mitigating the risks of disease and mortality resulting from illness or other health hazards.<sup>[2]</sup> Unfortunately, in many instances, health facilities are inadequately prepared for unforeseen disasters.

In Juba City, public health facilities were inadequately prepared for unforeseen disasters.<sup>[3]</sup> Chronic underfunding of the health sector by the government has worsened significantly over the past decade, negatively affecting the performance of public health facilities.<sup>[4]</sup> This study examined the factors influencing disaster preparedness in public health facilities in Juba City, South Sudan, which is a critical step towards enhancing health system resilience. The objectives of the study were to evaluate the influence of resource availability, health workers' capacity, technological integration, and institutional policy on disaster preparedness.

## Method

This was a cross-sectional study. The target population was 270 healthcare and administrative workers at three public health facilities in Juba City. Yamane's formula, with a 5% margin of error, was used to calculate a sample size of 162 respondents.

A structured self-administered questionnaire was used to collect data on the study variables (independent variables: availability of resources, health workers' capacity, technology integration, and institutional policy; and dependent variables: Time to Initiate Response, Adherence to Regulatory and Standard Guidelines, and Performance in Simulations and Training Drills).

The collected data were sorted, cleaned, coded, and entered into SPSS version 25; p-values < 0.05 were considered significant.

The results were presented in descriptive tables and the results of bivariate and logistic regression analysis. Since the targeted population was heterogeneous, a stratified sampling method was used to select the three health facilities (PHCU, PHCC, and a hospital), thereby capturing variation across healthcare levels.

## Results

148 participants responded to the study questionnaire. The largest proportion of respondents, 61 (41%), was within the 26-35 years age bracket. Disease outbreaks emerged as the most frequently reported event, cited by 76 (51%) respondents. A substantial proportion of health workers, 46 (31%), had less than one year of work experience. Most respondents 123, (83%) disagreed that their facility had sufficient funding for disaster preparedness activities. Furthermore, 105 (71%) disagreed that specific budgets were allocated for improving disaster preparedness, and 99 (67%) agreed that financial constraints significantly impact their facility's ability to prepare and respond during disasters (Table 1).

Regarding response initiation, although 99 (67%) of staff were aware of critical time objectives, only 70 (47%) facilities could initiate a response within the seven-day timeframe for detection, one day for notification, and seven days for response (Table 2). Regarding compliance with guidelines, a substantial majority, 108 (73%) of

**Table 1. Demographic characteristics of respondents (N=148)**

| Characteristic          | Category           | n (%)   |
|-------------------------|--------------------|---------|
| Sex                     | Male               | 70 (47) |
|                         | Female             | 78 (53) |
| Age (years)             | 19 – 25            | 18 (12) |
|                         | 26 – 35            | 61 (41) |
|                         | 36 – 45            | 42 (28) |
|                         | Above 45           | 27 (18) |
|                         | Certificate level  | 33 (22) |
| Education               | Diploma level      | 52 (35) |
|                         | Graduate/Degree    | 44 (30) |
|                         | Postgraduate       | 16 (11) |
|                         | Others (First Aid) | 3 (2)   |
| Work experience (years) | Less than 1        | 46 (31) |
|                         | 1 – 5              | 40 (27) |
|                         | 6 – 10             | 38 (26) |
|                         | Over 10            | 24 (16) |
| Types of disasters      | Flooding           | 25 (17) |
|                         | Disease outbreaks  | 76 (51) |
|                         | Heat wave          | 21 (14) |
|                         | Drought            | 2 (1)   |
|                         | Wildfire           | 0 (0)   |
|                         | Hailstorms         | 0 (0)   |
|                         | Landslides         | 0 (0)   |
|                         | Earthquakes        | 0 (0)   |
|                         | Civil war          | 24 (16) |
|                         | Others (Specify)   | 0 (0)   |

respondents, agreed their facility consistently adhered to local and national Disaster Preparedness and Response (DPR) guidelines. Despite this high level of adherence, a smaller proportion, 71 (48%), reported that staff were regularly trained on these guidelines (Table 2).

Concerning training and drills, only 63 respondents (43%) agreed that staff were trained in disaster response and drills (Table 2). A notable 98 (66%) of staff members

felt that participating in simulations and training drills significantly improved their preparedness for an actual disaster situation (Table 2). This observation suggests that, despite a relatively low frequency of reported training and drills, those that are conducted are highly valuable and effective for staff, indicating that increasing the frequency and scope of the simulations and drills could boost overall preparedness and staff confidence, even if other systemic issues persist.

**Table 2. Descriptive statistics of the dependent variables on disaster preparedness (N=148)**

| Statements                                                                                                                                                                                                | Disagree<br>n (%) | Agree<br>n (%) |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------|----------------|
| <b>Time to initiate response</b>                                                                                                                                                                          |                   |                |
| The facility can initiate its DPRP within seven days, as recommended by local disaster management authorities 7-1-7 (investigate within 7days, report within a day, and respond within 7 days) timelines. | 78 (53)           | 70 (47)        |
| The time it takes for our facility to mobilize resources for disaster response is adequate to meet the needs of the situation.                                                                            | 88 (59)           | 60 (41)        |
| The facility staff members are aware of the critical time objectives for initiating different phases of the disaster response.                                                                            | 49 (33)           | 99 (67)        |
| Communication with external emergency services is established promptly according to our disaster response plan.                                                                                           | 86 (58)           | 62 (42)        |
| <b>Compliance with guidelines</b>                                                                                                                                                                         |                   |                |
| The facility consistently adheres to local and national disaster preparedness and response guidelines.                                                                                                    | 40 (27)           | 108 (73)       |
| The facility staff members are regularly trained on regulatory and standard guidelines relevant to disaster preparedness and response.                                                                    | 77 (52)           | 71 (48)        |
| The disaster preparedness and response plans are regularly audited for compliance with regulatory and standard guidelines.                                                                                | 93 (63)           | 55 (37)        |
| Feedback from compliance audits is effectively used to improve our disaster preparedness and response plans.                                                                                              | 69 (47)           | 79 (53)        |
| <b>Training and drills</b>                                                                                                                                                                                |                   |                |
| Staff are trained in disaster response and in drills.                                                                                                                                                     | 85 (57)           | 63 (43)        |
| Participants in simulations and training drills demonstrate a high level of competence in their designated roles.                                                                                         | 67 (45)           | 81 (55)        |
| Simulations and training drills are realistic and cover a wide range of potential disaster scenarios.                                                                                                     | 72 (49)           | 76 (51)        |
| The performance in simulations and training drills is systematically evaluated to identify areas for improvement.                                                                                         | 69 (47)           | 79 (53)        |
| Lessons learned from simulations and training drills are effectively integrated into our disaster preparedness and response plans.                                                                        | 80 (54)           | 68 (46)        |
| Staff members feel that participating in simulations and training drills significantly improves their preparedness for an actual disaster situation.                                                      | 50 (34)           | 98 (66)        |

Regarding staffing levels, only 69 (47%) agreed that there was sufficient staffing for emergencies (Table 3). Although a higher proportion, 85 (57%), indicated an adequate number of specialized healthcare workers for disaster response, a significant majority, 98 (66%), disagreed that staffing levels remain stable during emergencies, and only 57 (39%) agreed that staff allocation was effectively adjusted during a disaster (Table 3).

The material resources assessment indicated significant deficiencies, with 98 (66%) respondents disagreeing that their facility possessed adequate physical infrastructure to manage large-scale health emergencies (Table 3). A notable positive outlier in resource availability of reliable access to utilities (water and electricity), with the majority 101 (68%) agreeing. However, this contrasts sharply with the significant 94 (64%) disagreement on the availability

**Table 3. Descriptive statistics of the independent variables on disaster preparedness (N=148)**

| Statements                                                                                                                                       | Disagree<br>n (%) | Agree<br>n (%) |
|--------------------------------------------------------------------------------------------------------------------------------------------------|-------------------|----------------|
| <b>Health workers capacity</b>                                                                                                                   |                   |                |
| The current staffing level at our health facility is sufficient to manage emergencies and disasters effectively.                                 | 79 (53)           | 69 (47)        |
| The facility has an adequate number of specialized healthcare workers (e.g., Lab techs, Paediatricians, health experts) for disaster response.   | 63 (43)           | 85 (57)        |
| The staffing levels in our facility remain stable and unaffected by emergencies or disasters.                                                    | 98 (66)           | 50 (34)        |
| The allocation of staff across various departments within our facility is effectively adjusted in response to a disaster.                        | 91 (62)           | 57 (39)        |
| <b>Resources Availability</b>                                                                                                                    |                   |                |
| The physical infrastructure of our facility (e.g., buildings, beds, wards) is sufficient to accommodate an influx of patients during a disaster. | 98 (66)           | 50 (34)        |
| The facility has reliable access to utilities (e.g., water, electricity) essential for disaster response.                                        | 47 (32)           | 101 (68)       |
| <b>Technology Integration</b>                                                                                                                    |                   |                |
| The technology infrastructure in our facility adequately supports disaster preparedness efforts                                                  | 94 (64)           | 54 (36)        |
| The facility has sufficient access to the technological resources necessary for responding to disasters.                                         | 102 (69)          | 46 (31)        |
| Technological tools are readily available in our facility for staff to use in disaster preparedness activities.                                  | 105 (71)          | 43 (29)        |
| The facility has Data management systems in place that allow for efficient tracking and monitoring of public health risks.                       | 44 (30)           | 104 (70)       |
| Technology has improved the decision-making process in disaster preparedness and response at our facility                                        | 58 (39)           | 90 (61)        |
| Technological integration supports a more coordinated approach to disaster management across different departments.                              | 49 (33)           | 99 (67)        |
| The use of technology facilitates the quicker adoption of public health strategies in the aftermath of a disaster.                               | 63 (43)           | 85 (57)        |
| <b>Institutional policy</b>                                                                                                                      |                   |                |
| The facility has a disaster preparedness committee.                                                                                              | 100 (68)          | 47 (32)        |
| The committee has a budget allocated for disaster preparedness.                                                                                  | 109 (74)          | 35 (24)        |
| The facility has a Disaster Preparedness and Response Plan (DPRP)                                                                                | 89 (60)           | 59 (40)        |

**Table 4. Bivariate analysis**

| Independent Variable (Key Indicator)      | Dependent Variable (Aspect of Preparedness) | Chi-square ( $\chi^2$ ) | p-value |
|-------------------------------------------|---------------------------------------------|-------------------------|---------|
| Financial Resources (Sufficient Funding)  | Ability to initiate response within 7 days  | 15.24                   | <0.001  |
| Institutional Policy (Has DPRP)           | Adherence to guidelines                     | 18.51                   | <0.001  |
| Health Worker Capacity (Staff Motivation) | Effectiveness of training drills            | 9.87                    | 0.002   |

of reliable communication systems specifically for disaster response (Table 3).

While a high majority, 104 (70%), agreed that their facility had data management systems for tracking public health risks, only 54 (36%) agreed that the current technological infrastructure adequately supports preparedness, and even fewer, 46 (31%), agreed that access to technological resources, such as communication devices and early warning systems, was sufficient (Table 3). Additionally, only 43 (29%) agreed that technological tools were readily available for staff use. Furthermore, 90 (61%) agreed that technology had improved decision-making, 99 (67%) agreed that it supported a more coordinated approach, and 85 (57%) agreed that it contributed to the quicker adaptation of strategies (Table 3).

Regarding institutional policy, a significant majority, 100 (68%) of respondents disagreed that their facility has a disaster preparedness committee, and even fewer, 35 (24%) agreed that such a committee has a budget allocated for disaster preparedness (Table 3). Furthermore, 89 (60%) disagreed that their facility had a formal DPRP Plan, indicating that disaster preparedness is largely informal or ad hoc rather than systematically integrated into the facility's operations (Table 3).

Both descriptive and inferential statistical analyses showed institutional policy and financial constraints had a strong, statistically significant negative association with a facility's inability to initiate a timely disaster response and mobilize resources adequately. This finding provides statistical evidence for the link between financial resources and preparedness outcomes. The reported insufficient funding for preparedness activities was significantly associated with a facility's inability to initiate a response plan within the recommended timeframe ( $\chi^2=15.24$ ,  $p<0.001$ ). Similarly, the lack of formal policies was statistically linked to lower preparedness ( $\chi^2=18.51$ ,  $p<0.001$ ).

#### Multiple binary logistic regression analysis

Multivariable binary logistic regression was employed to

identify independent predictors of disaster preparedness, while controlling for other variables (Tables 4 and 5). For this analysis, the dependent variable, "Disaster Preparedness," was converted into a binary outcome (e.g., "Prepared" vs. "Not Prepared" using the Likert scale from the "Time to Initiate Response," "Compliance with Guidelines," and "Training and Drills" categories. Similarly, the independent variables from the Likert scale were transformed into binary categories (e.g., "Sufficient Funding" vs. "Insufficient Funding," "Has DPRP" vs. "No DPRP"). The Likert scale responses "strongly disagree and strongly disagree" were considered as disagree, while "agree, moderately agree, and strongly agree" were regarded as agree.

The descriptive and inferential statistics confirmed that institutional policies and financial resources were among the strongest independent predictors of preparedness. The Odds Ratios (OR) quantified the magnitude of influence for each determinant. The analysis revealed that facilities with a Disaster Preparedness and Response Plan (DPRP) had an Odds Ratio of 3.49 ( $p=0.001$ ), indicating they were nearly 3.5 times more likely to be prepared than those without one, holding other factors constant. Similarly, the Odds Ratio for "Sufficient Funding" was 2.66 ( $p=0.002$ ), indicating that facilities with adequate financial resources were 2.66 times more likely to be prepared than those with insufficient funding.

#### Discussion

**Demographic characteristics:** The largest proportion of respondents fell within the 26-35 age bracket, suggesting a workforce predominantly in its younger-to-middle career stages, balancing experience with potential for long-term development. In terms of education, diploma holders represented the largest group, 52 (35%), followed closely by graduate/degree holders, 44 (30%), indicating a varied educational background. A notable observation was that a substantial proportion of health workers, 46 (31%), have less than one year of experience, which could be indicative of a high turnover rate or a recent expansion

**Table 5. Multivariate analysis**

| Variable                                                       | Odds Ratio | p-value |
|----------------------------------------------------------------|------------|---------|
| <b>Institutional Policy: Has DPRP</b>                          |            |         |
| No Disaster Preparedness & Response Plan (DPRP) (ref)          | 1          |         |
| Has DPRP                                                       | 3.49       | 0.001   |
| <b>Financial Resources: Sufficient Funding</b>                 |            |         |
| Insufficient Funding (ref)                                     | 1          |         |
| Sufficient Funding                                             | 2.66       | 0.002   |
| <b>Health Worker Capacity: Highly Motivated Staff</b>          |            |         |
| Low Motivation (ref)                                           | 1          |         |
| Highly Motivated Staff                                         | 2.05       | 0.013   |
| <b>Technological Integration: Adequate Tech Infrastructure</b> |            |         |
| Inadequate Tech Infrastructure (ref)                           | 1          |         |
| Adequate Tech Infrastructure                                   | 1.57       | 0.072   |
| Significance P<0.05                                            |            |         |

of the workforce.

**Influence of resource availability on preparedness:** Financial resources emerged as a major constraint on effective disaster preparedness, with 123 (83%) respondents reporting insufficient funding. This appeared to directly contribute to deficiencies in physical infrastructure and the procurement of adequate medical supplies. The bivariate and multivariate analyses confirm that financial deficiency is a significant negative predictor of overall preparedness, with facilities that have sufficient funding significantly more likely to be prepared. This aligns with a study conducted in Nigerian healthcare facilities that revealed inadequate funding, poor allocation, and mismanagement of available funds dedicated to disaster preparedness as critical barriers to acquiring necessary supplies and implementing disaster preparedness programmes. These findings suggest an urgent need for increased financial investment and improved financial management systems. <sup>[5,6]</sup>

**Influence of institutional policies on preparedness:** The absence of an institutional disaster preparedness policy is another critical finding, with 89 (60%) disagreeing that their facility has a formal Disaster Preparedness and Response Plan (DPRP). The binary logistic regression analysis confirmed institutional policies as one of the strongest independent determinants of disaster preparedness, with the presence of formal plans that provide a clear framework to drive preparedness efforts. This finding is supported by a study conducted in Ethiopian hospitals, which identified that the lack of formal plans and committees hindered disaster preparedness. <sup>[7,8]</sup>

**Influence of health workers' capacity on preparedness:** A significant concern is the overall staffing levels and their stability during emergencies, suggesting that staff retention during crises is lacking. Furthermore, the low reported motivation among health workers to participate in preparedness activities indicates that even when staff possess skills, a lack of motivation can lead to reduced engagement and reluctance to participate effectively. This is consistent with a study that identified insufficient staffing as a significant challenge in disaster management and maintaining quality of care. <sup>[9]</sup>

**Influence of technological integration on preparedness:** Despite the existence of data management systems and data security, a significant deficiency in overall technological infrastructure, access to communication gadgets and early warning systems, and the general availability of technological tools for staff use in disaster preparedness. This "digital divide" implies that while internal data management might be available, the ability to disseminate information in a timely manner, communicate in real time during crises, and provide proactive early warning is severely hindered. Other studies report disparities between basic ICT infrastructure and specialized disaster technologies in low- and middle-income countries. <sup>[10]</sup> Despite these infrastructure limitations, most respondents recognized that technology had improved decision-making, indicating that even the limited technological integration that exists is highly valued and recognized for its positive impact. This suggests that targeted investment in disaster-specific technological infrastructure could

significantly amplify preparedness benefits.<sup>[11,12]</sup>

## Recommendations

The study first recommends an urgent need for policy formalization and the establishment of disaster-preparedness committees within each public health facility, with clearly defined roles and responsibilities. This recommendation aligns with the foundational principles of effective disaster management advocated in their guide for developing hospital disaster preparedness plans.<sup>[13]</sup>

Secondly, ensure dedicated financial investment for disaster preparedness activities. Highlighted financial constraints as a major barrier to disaster risk reduction in low- and middle-income countries.<sup>[14]</sup>

Thirdly, implement comprehensive workforce planning to ensure adequate staffing and retention, preventing the system from being overwhelmed during disasters. A study found insufficient staffing to be a major gap in disaster management.<sup>[15]</sup>

## Conclusion

The overall state of readiness is significantly hampered by systemic deficiencies, with the absence of formal institutional policies and severe financial constraints as the most profound challenges. These foundational weaknesses created a cascading effect, leading to inadequate material resources, challenges in maintaining stable, motivated human resources, and a disconnect in general information.

**Source of funding:** This research was funded by Dumba Samuel Kenyi, without additional external financial support.

**Conflict of interests:** None

## References

- World Health Organization. (2022). Health systems resilience toolkit: A WHO global public health goal to support building and strengthening sustainable health systems resilience in countries with various contexts. <https://www.who.int/publications/i/item/9789240048751>
- Gillani AH, Izham M, Ibrahim M, Akbar J. (2020). Evaluation of disaster medicine preparedness among healthcare profession students. New York: Crown Press.
- Martinez D, Talbert T, Romero-Steiner S, Kosmos C, Redd S. Evolution of the Public Health Preparedness and Response Capability Standards to Support Public Health Emergency Management Practices and Processes. *Health Security* 2019;17(6):430–438
- Nyirenda T, Bockarie M, Machingaidze S. et al. Strengthening capacity for clinical research in sub-Saharan Africa: partnerships and networks. *International Journal of Infectious Diseases* 2021; 110: 54–61.
- Odigwe CC, Okoli UC, Onwuegbuzie AJ. Challenges of disaster preparedness in healthcare facilities in Nigeria: A cross-sectional study. *International Journal of Disaster Risk Reduction* 2021;59: 114.
- Barclay LE, Gimble AM. The impact of resource availability on hospital disaster preparedness in low- and middle-income countries: A systematic review. *Prehospital and Disaster Medicine* 2022;37(2):182-189.
- Deressa A, Ali J. Assessment of disaster preparedness in public hospitals in Addis Ababa, Ethiopia. *Ethiopian Journal of Health Development* 2020;34(2):147-154.
- Deressa A, Ali J, Alem A. Assessment of communication preparedness for disasters in public health facilities in Addis Ababa, Ethiopia. *Ethiopian Journal of Health Development* 2020;34(6):576-583.
- Carter H, Thompson J. Challenges for nurses in disaster management: A scoping review. *Journal of Nursing Management* 2021;29(5):1024-1034.
- Owolabi OO, Aderounmu GA, Adeyemi OS. Challenges and opportunities for using information and communication technology for disaster preparedness in health facilities in low- and middle-income countries: a systematic review. *International Journal of Disaster Risk Reduction* 2021; 58:193-205
- Bruns A, Meltzer DE. The Role of Social Media in Hospital Communication During Disasters. *Journal of Health Communication* 2021;26(1):74-82
- Ibrahim MH, Hawryluk AM. The Role of Information and Communication Technologies in Disaster Risk Management in Healthcare

- Facilities. *Journal of Hospital Administration* 2019;64(2):102-111.
13. Lewis DD, Woloshin C J, Bates DW. Effective communication strategies for disaster preparedness in healthcare settings. *Disaster Medicine and Public Health Preparedness* 2022;16(2):290–297.
14. Awokola OS, Fewtrell L, Nguyen VC. (2020). Financing disaster risk reduction in the health sector of low- and middle-income countries: A systematic review. *International Journal of Disaster Risk Reduction* 2020; 49:3
15. Carter H, Thompson J. Challenges for nurses in disaster management: A scoping review. *Journal of Nursing Management* 2021;29(5):1024–1034.

### Why sleeping sickness pill is raising hopes in Africa

- One-dose oral treatment for sleeping sickness gets European approval
- High success rate in human trials in Democratic Republic of Congo and Guinea
- The pill could enable “test-and-treat” strategies in remote African communities, say researchers

[NAIROBI, SciDev.Net] Sleeping sickness has been killing people across Africa for generations. A [disease](#) with no vaccine, spread by the bite of a tsetse fly, it was once treated with injections of an arsenic derivative, a cure that could be as dangerous as the illness itself. By 1998, it was infecting [40,000 people](#) a year.

Last month, the single oral-dose [medicine](#) Acoziborole Winthrop (acoziborole) became the newest and simplest weapon against the disease when it was given the nod for approval by European regulators.

The treatment was developed by French pharmaceutical company Sanofi in partnership with the Geneva-based [Drugs for Neglected Diseases Initiative \(DNDi\)](#), with human trials in West and Central Africa.

Now researchers are pushing further: a [paediatric trial, ACOZI-KIDS](#), is now underway in Democratic Republic of Congo and Guinea to find out whether the same drug can work for children aged one to 14, the one group the current approval does not yet cover.

<https://www.scidev.net/global/explainer/why-sleeping-sickness-pill-is-raising-hopes-in-africa/>

# Evaluating the role of the health information system infrastructure in enhancing data quality in Central Equatoria State, South Sudan

Matur T. Tieng<sup>1</sup>,  Karani Magutah<sup>2</sup>, Joseph Muchiri<sup>3</sup>

## Author Affiliation:

1. Department of Public Health, School of Public Health, Mount Kenya University.
2. Department of Medical Physiology, School of Medicine, Moi University, Eldoret, Kenya
3. Department of Public Health, School of Public Health, Mount Kenya University, Thika Campus, Kenya

## Correspondence:

Matur T. Tieng  
[themdit.matur@gmail.com](mailto:themdit.matur@gmail.com)

**Submitted:** January 2026

**Accepted:** March 2026

**Published:** May 2026

**Citation:** Tieng et al Evaluating the role of the Health Information System infrastructure in enhancing data quality in Central Equatoria State, South Sudan. South Sudan Medical Journal, 2026;19(2):107-111 © 2026 The Author (s) **License:** This is an open access article under [CC BY-NC](#) DOI: <https://dx.doi.org/10.4314/ssmj.v19i2.7>

## ABSTRACT

**Introduction:** A health information system (HIS) infrastructure is a critical foundation to produce high-quality routine health data for planning, monitoring, and decision-making, particularly in low-resource and fragile settings such as South Sudan. The aim of this study was to assess the influence of HIS infrastructure on the quality of the Health Management Information System (HMIS) data in public health facilities in Central Equatoria State, South Sudan.

**Method:** A cross-sectional study was conducted using a self-administered, structured questionnaire completed by health workers at the public health facilities. Descriptive statistics and Multivariate Analysis of Variance were produced using Stata software version 15.

**Results:** 139 health workers participated in the study. About half (50.4%) of respondents reported routine use of the District Health Information System version 2 (DHIS2); 30.2% of facilities had functional computers or tablets; and 53.2% lacked reliable internet connectivity. The HIS infrastructure factors collectively affect HMIS data quality, as shown by Wilks' Lambda = 0.507 ( $p < 0.001$ ).

**Conclusion:** Inadequate digital infrastructure, inconsistent training coverage, and limited system utilization undermine HMIS data quality in Central Equatoria State. Strengthening HIS infrastructure, alongside standardized capacity-building and improved reporting processes, is essential to improving routine health data quality in South Sudan.

**Keywords:** health information system, infrastructure, data quality, health facilities, South Sudan

## Introduction

Healthcare systems depend on reliable information from a well-established Health Management Information System (HMIS). As one of the health system building blocks, HMIS plays a significant role in collecting, storing and analysing data to support better decision-making.<sup>[1,2]</sup> For enhanced quality data, HIS infrastructures, such as availability of computers, Internet connectivity, and software, need to be adequate. This is because data quality impacts all aspects of a business, including operational efficiency, compliance, and customer engagement. In healthcare, poor-quality data leads to misallocation of resources, wrong decision-making,

and risks to patients' lives and safety.<sup>[3]</sup>

The HIS infrastructure refers to all components, including, but not limited to, hardware, software, networking, services, and policies that enable the collection, storage, and sharing of data.<sup>[4]</sup> The HIS infrastructure has two components: the Information Technology (IT), which deals with hardware and software, and the human resource infrastructure, which encompasses skills, training, and motivation.<sup>[5]</sup>

The rollout of the District Health Information System version 2 in 2019 was critical, establishing a robust HIS infrastructure that drives digital innovation.<sup>[6]</sup> Empirical evidence shows that robust HIS infrastructure has modernized the health service delivery in Ethiopia by enhancing health workers' response to patients and improving data quality.<sup>[7]</sup> This study examined how the HIS infrastructure, including the availability of computers/tablets, training on the HIS, and software, influences data quality at the public health facilities in Central Equatoria, South Sudan.

## Method

### Study population

This was a cross-sectional quantitative study. The target population included data clerks/monitoring and evaluation (M&E) officers, data managers, and other health cadres who were working at public health facilities in Central Equatoria State. We sampled 148 respondents. A self-administered questionnaire was completed by the health care workers between June and December 2025.

### Data collection

Quantitative data were collected using a structured self-administered questionnaire covering demographic characteristics, training exposure, infrastructure availability, system utilisation, and perceived challenges in data entry and reporting.

### Data analysis

Data were entered and analysed using STATA version 15. Descriptive statistics (frequencies and percentages) were generated. Additionally, a Multivariate Analysis of Variance (MANOVA) was performed to determine whether there were significant differences among infrastructural variables, such as facility electricity access and Internet connectivity, in the HMIS data quality dimensions (completeness, consistency, and timeliness).

## Ethical considerations

Ethical approval was obtained from the Mount Kenya University Institutional Scientific and Ethics Review Committee (Reference: MKU/ISERC/4883) and the Ministry of Health Research Ethics and Review Board, South Sudan (Reference: MOH/RERB/P/A/15/8/2024-MOH/REBR/A/19-N/2024). Written, informed consent was obtained from all participants, and confidentiality was maintained throughout.

## Results





Figures 1-4 show the demographic characteristics of the 139 respondents out of 148 sampled (93.9% response rate): 65.5% were male, and 35% were female, with 45% aged 26-33 years, followed by 34-41 years (32.4%). The respondents' educational attainment varied significantly, with most holding higher education certificates (61.9%), followed by secondary education (38.1%). Most of the respondents studied social sciences (21.6%), health informatics (19.4%), clinical medicine (18.7%), and IT (17.3%).

The descriptive statistics in Table 1 show the HIS infrastructure factors that influence data quality in the District Health Information Software (DHIS) system. The findings revealed that only 50.4% were using DHIS2, indicating a suboptimal rollout across health facilities. 63.3% of those using DHIS2 received training on the system, yet only 46% of those trained rated the training as adequate. Additionally, 50% of those who received training on DHIS2/HMIS covered all aspects of the training.

The system's ease of use was rated positively by most respondents, with 82.7% finding it easy to use. However, a small proportion (16.1%) of respondents struggled with the system, possibly due to interface complexities or technical limitations. The availability of computers or tablets in health facilities was reported to be limited: only 30.2% of respondents confirmed their presence, and 81% reported functionality.

The major challenge reported by 53.2% of the respondents was poor internet connectivity and a lack of data bundles.

The MANOVA shown in Table 2 examines the HIS infrastructural determinants of HMIS data quality, assessing factors such as the DHIS2 system, training, tablets, reasons for delayed on-time data entry, and challenges facing the data entry process. The overall model was statistically significant (Wilks'  $\lambda = 0.5072$ ,  $p$



Figures 1-4. Demographic information of the respondents

= 0.0005), indicating that the HIS infrastructural factors jointly significantly affect the quality of HMIS data.

However, at the individual factor's level such as availability of DHIS 2 (Wilks'  $\lambda = 0.9562$ ,  $p = 0.2707$ ), training on the DHIS2/HMIS (Wilks'  $\lambda = 0.9394$ ,  $p = 0.1405$ ), the availability of computers/tablets (Wilks'  $\lambda = 0.9607$ ,  $p = 0.3196$ ) were not statistically significant, indicating that the above factors cannot individually impact on the quality of HMIS data.

Nevertheless, other factors, such as reasons for the delay in data entry, perhaps related to motivation, data entry protocols, or reporting procedures (Wilks'  $\lambda = 0.8082$ ,  $p = 0.0266$ ), challenges, such as internet connectivity (Wilks'  $\lambda = 0.7231$ ,  $p = 0.0042$ ), were statistically significant, confirming their potential for impacting the quality of HMIS data.

## Discussion

In this study, there were more males than females. This sex disparity reflects employment ratios within the public health sector.<sup>[8]</sup> Conversely, a similar study in Benin found that 81.7% of respondents were female.<sup>[9]</sup> Our findings revealed that respondents' fields of study varied widely, with most having studied social sciences, health informatics, clinical medicine, and IT. This diversity emphasizes the multidisciplinary nature of, and the need required in, the public health workforce. A study conducted in Ethiopia in 2015 found that those tasked with data collection and reporting were nurses, accounting for 44% of the total respondents,<sup>[10]</sup> unlike in South Sudan, where the responsibilities for data recording and reporting rested with data clerks and M&E officers.

The use of DHIS2 in South Sudan, like other countries such as Uganda<sup>[6]</sup> is mandatory for all health facilities owned by the government, and is one of the HIS infrastructures upon which this study focused. The infrastructural determinants

**Table 1. Descriptive statistics for HIS infrastructural determinants of quality of DHIS (N = 139)**

| Variable                                            | Responses                     | n (%)     |
|-----------------------------------------------------|-------------------------------|-----------|
| Datasets using DHIS 2 system for reporting?         | Yes                           | 70 (50.4) |
|                                                     | No                            | 69 (49.6) |
| DHIS/HMIS training received                         | Yes                           | 88 (63.3) |
|                                                     | No                            | 51 (36.7) |
| Topics covered during the training                  | Data entry                    | 20 (22.7) |
|                                                     | Data Analysis                 | 8 (9.1)   |
|                                                     | Data Quality Check            | 1 (1.1)   |
|                                                     | All HMIS tools and DHIS2      | 18 (20.5) |
|                                                     | All of them                   | 38 (43.2) |
|                                                     | Others                        | 3 (3.4)   |
| Adequacy of the HMIS/DHIS2 training                 | Inadequate                    | 10 (11.5) |
|                                                     | Somehow adequate              | 22 (25.3) |
|                                                     | Adequate                      | 40 (46.0) |
|                                                     | Very adequate                 | 15 (17.2) |
| System use after HMIS/DHIS2 training                | Very difficult                | 5 (5.8)   |
|                                                     | Difficult                     | 9 (10.3)  |
|                                                     | Easy                          | 59 (67.8) |
|                                                     | Very easy                     | 13 (14.9) |
|                                                     | Missing value                 | 1 (1.2)   |
| Computers/tablets available at the facility         | Yes                           | 42 (30.2) |
|                                                     | No                            | 97 (69.8) |
| Functionality of Computers/tablets                  | Yes                           | 34 (81.0) |
|                                                     | No                            | 8 (19.1%) |
| Other Means of reporting in case no computer/tablet | Reporting through CHD         | 74 (76.3) |
|                                                     | Using mobile phone            | 18 (18.6) |
|                                                     | Borrow computer from friends  | 5 (5.2)   |
| Challenges facing data entry process                | Poor system design            | 16 (11.5) |
|                                                     | Incomplete data system        | 20 (14.4) |
|                                                     | Inaccurate data               | 18 (13.0) |
|                                                     | Poor internet/lack of bundles | 74 (53.2) |
|                                                     | Others                        | 11 (7.9)  |

assessed included the availability of HMIS reporting tools, such as HMIS registers, reporting forms, computers/tablets, and the overall database. The investigation encompassed the training on DHIS2 or HMIS tools, the topic covered, and whether the training was adequate. The influence of infrastructural factors on the quality of HMIS data was assessed using logistic regression and MANOVA, with a significance level of 5%. The results showed that about 50% of respondents used the DHIS2 system and were familiar with HMIS tools, of whom about 60% had received training on the HMIS/DHIS2 system.

Despite many respondents having received training on DHIS2 and HMIS tools, the training was inadequate or not well integrated, as trainees didn't cover all essential topics as per the HMIS training manual, which could lead to differing levels of proficiency among users. A study on DHIS2 utilization in Uganda and Ethiopia reported utilization rates above 90% and 80%, respectively,<sup>[7,8]</sup> far higher than in South Sudan. Only 46% of those who received training found it adequate, casting doubt on its efficiency and effectiveness. The robust HMIS system is anchored in the skilled human resources trained on the system and its tools, a comprehensive system design, and reliable internet connectivity.<sup>[9,10]</sup>

As computers or tablets play a critical role in reporting and the utilization of the DHIS2 in general, their availability and functionality remain a major concern, as many health facilities do not have computers/tablets. This highlights a critical infrastructural gap that hinders data reporting and compromises the quality of datasets collected through the DHIS2 system.

## Conclusion

HIS infrastructure was a key determinant of the quality of data generated at health facilities. Despite the adoption of DHIS2 as the national HIS, its utilization at the subnational level remains suboptimal, with 50% of the sampled facilities reporting through the county health department. It was also noted that most health workers who have received training on HMIS/DHIS2 have not been trained on all aspects of the curriculum, suggesting gaps in their knowledge. The limited availability of computers/tablets or HMIS tools for reporting and poor Internet connectivity further constrained the timely reporting of data into DHIS2. We recommend completing the rollout of the DHIS2 system, standardizing training, and increasing investment in HIS infrastructure.

## References

**Table 2. Multivariate analysis of HIS Infrastructural determinants of quality of DHIS datasets**

| Source            | Wilks' $\lambda$ | df | F (df1, df2) | F    | p-value |
|-------------------|------------------|----|--------------|------|---------|
| Model             | W 0.5072         | 10 | (30;256)     | 2.22 | 0.0005  |
| Residual          |                  | 89 |              |      |         |
| DHIS2             | W 0.9562         | 1  | (3;87)       | 1.33 | 0.2707  |
| Training          | W 0.9394         | 1  | (3;87)       | 1.87 | 0.1405  |
| Tablets           | W 0.9607         | 1  | (3;87)       | 1.19 | 0.3196  |
| Reasons for delay | W 0.8082         | 3  | (9;211.9)    | 2.15 | 0.0266  |
| Challenges        | W 0.7231         | 4  | (12;230.5)   | 2.5  | 0.0042  |

- Dagnew E, Woreta SA, Shiferaw AM. Routine health information utilization and associated factors among health care professionals working at public health institution in North Gondar, Northwest Ethiopia. *BMC Health Serv Res*. 2018;18:685. <https://doi.org/10.1186/s12913-018-3498-7>
- Kebede M, Adeba E, Chego M. Evaluation of quality and use of health management information system in primary health care units of east Wollega zone, Oromia regional state, Ethiopia: *BMC Med Inform Decis Mak*. 2020;20:107. <https://doi.org/10.1186/s12911-020-01148-4>
- Omer MK. The Contribution of Health Management Information Systems to Enhancing Healthcare Operations. *Int J Adv Comput Sci Appl [Internet]*. 2023 [cited 2025 June 3];14. <https://doi.org/10.14569/IJACSA.2023.0141263>
- Ministry of Health, South Sudan. Towards health system resilience Key Health Management Information System (HMIS) interventions and progress in South Sudan [Internet]. 2025 [cited 2025 Dec 6]. <https://moh.gov.ss/>. Accessed 6 Dec 2025
- Zerfu TA, Asressie M, Begna Z, Habtamu T, Werkneh N, Nigatu T, et al. Unveiling the role of DHIS2 in enhancing data quality and accessibility in primary healthcare facilities: Evidence from Ethiopia. *Demsash AW*, editor. *PLOS ONE*. 2024;19:e0314505. <https://doi.org/10.1371/journal.pone.0314505>
- Kiwanuka SN, Kabwama SN, Namuhani N, Gupta S, Layer E, Mechael P, et al. Barriers and facilitators of the effective use of DHIS2 data to improve program planning and monitoring in Uganda: a sequential mixed methods study. *Oxf Open Digit Health*. 2026;4:oqag002. <https://doi.org/10.1093/oodh/oqag002>
- Basajja M, Paul-Chima UO. Comparative Effectiveness of DHIS2 and FAIR Data Approaches for Privacy-Preserving Health Data Analytics in Uganda: A Systematic Review. *Clin Outcomes Res CEOR*. 2025;17:849–64. <https://doi.org/10.2147/CEOR.S560265>
- Teklegiorgis K, Tadesse K, Terefe W, Mirutse G. Level of data quality from Health Management Information Systems in a resources limited setting and its associated factors, eastern Ethiopia : original research. *South Afr J Inf Manag. AOSIS*; 2016;18:1–8. <https://doi.org/10.4102/sajim.v17i1.612>
- Global Strategy on Digital Health 2020-2025. 1st ed. Geneva: World Health Organization; 2021.
- WHO Toolkit for Routine Health Information Systems Data <https://www.who.int/data/data-collection-tools/health-service-data/toolkit-for-routine-health-information-system-data/modules>. Accessed 14 Feb 2026

# Impact of developmental stimulation education on maternal knowledge and caregiving practices in stunting: a longitudinal study

Ning Iswati<sup>1,2</sup>,  Azizan Omar<sup>3</sup>,  Sukhbeer Kaur Darsin Singh<sup>4</sup>,  Putra Agina Widyaswara Suwaryo<sup>5</sup> 

## Author Affiliation:

1. Faculty of Medicine and Health Science, Universiti Malaysia Sabah, Kinabalu, Malaysia
2. Department of Paediatric Nursing, Faculty of Health Science, Universitas Muhammadiyah Gombong, Kebumen, Indonesia
3. Department of Public Health, Faculty of Medicine and Health Science, Universiti Malaysia Sabah, Kinabalu, Malaysia
4. Department of Nursing, Faculty of Medicine and Health Science, Universiti Malaysia Sabah, Kinabalu, Malaysia
5. Department of Nursing, Faculty of Health Science, Universitas Muhammadiyah Gombong, Kebumen, Indonesia.

## Correspondence:

Ning Iswati  
[ningiswati@unimugo.ac.id](mailto:ningiswati@unimugo.ac.id)

**Submitted:** January 2026

**Accepted:** May 2026

**Published:** February 2026

---

## ABSTRACT

**Introduction:** Stunting is associated not only with impaired physical growth but also with suboptimal child development, partly influenced by inadequate caregiving and insufficient developmental stimulation. This study aimed to evaluate the impact of a developmental stimulation education programme on maternal knowledge and caregiving practices among mothers of children with stunting.

**Method:** A longitudinal quasi-experimental study with individual-level random allocation was conducted between January and November 2025 involving 87 mothers of children with stunting, of whom 80 completed follow-up and were included in the final analysis. Both groups received routine child health services; however, only the intervention group received a structured developmental stimulation education programme, which included age-appropriate guidance on motor, language, cognitive, and social stimulation integrated into daily caregiving activities and delivered through standardised sessions by trained facilitators. Maternal knowledge and caregiving practices were assessed at baseline, 1 month, 3 months, and 6 months. Data were analysed using generalised linear mixed models.

**Results:** Maternal knowledge and caregiving practices improved over time in both groups; however, the intervention group demonstrated significantly greater and more sustained improvements. The group-by-time interaction was statistically significant for both outcomes, indicating that the intervention altered the trajectory of maternal knowledge and caregiving practices beyond changes observed with routine care.

**Conclusion:** The developmental stimulation education programme produced sustained improvements in maternal knowledge and caregiving practices over six months. Integrating caregiver-focused educational interventions into routine maternal and child health services and community-based programmes may strengthen comprehensive stunting management and support early child development.

**Keywords:** stunting, developmental stimulation, maternal knowledge, caregiving practices, longitudinal study.

---

**Citation:** Iswati et al. Impact of developmental stimulation education on maternal knowledge and caregiving practices in stunting: a longitudinal study. *South Sudan Medical Journal*, 2026;19(2):112-118 © 2026 The Author (s) **License:** This is an open access article under [CC BY-NC](#) DOI: <https://dx.doi.org/10.4314/ssmj.v19i2.8>

## Introduction

Stunting remains a major public health problem with long-term consequences for child health and development.<sup>[1,2]</sup> In addition to impaired physical growth, stunting is associated with delays in cognitive, motor, and socio-emotional development that may persist into adulthood.<sup>[3,4]</sup> Although undernutrition is a key determinant, inadequate psychosocial stimulation and suboptimal caregiving practices also contribute substantially to poor developmental outcomes.<sup>[5]</sup> Therefore, nutrition-focused interventions alone may be insufficient to address the broader developmental risks associated with stunting.

Caregiver-based developmental stimulation is particularly important during early childhood, a period of high neural plasticity.<sup>[6]</sup> Mothers, as primary caregivers, play a key role in providing stimulation through everyday interactions such as play, communication, and responsive care. However, limited maternal knowledge and suboptimal caregiving practices may reduce the effectiveness of such stimulation, especially among children affected by chronic undernutrition. Educational interventions targeting maternal knowledge and caregiving practices, therefore, represent a promising strategy to improve developmental environments for children with stunting.<sup>[7]</sup>

Stunting remains highly prevalent in low- and middle-income countries, where socioeconomic constraints and limited access to early childhood interventions persist.<sup>[8]</sup> Given its scale, even modest improvements in caregiving practices at the household level may produce meaningful population-level benefits. This underscores the need for interventions that are feasible, scalable, and capable of generating sustained developmental gains beyond short-term nutritional recovery. Previous studies on caregiver education for children with stunting have reported promising but inconclusive results, largely due to cross-sectional designs, short follow-up periods, and limited analytical approaches.<sup>[9]</sup> In addition, many studies emphasized child outcomes without adequately examining maternal knowledge and caregiving practices as proximal determinants of developmental change, thereby limiting causal interpretation.

This study evaluated the impact of a structured developmental stimulation education programme on maternal knowledge and caregiving practices among mothers of children with stunting using a longitudinal design with repeated measurements over six months. By applying a mixed-effects analytical approach, the study examined changes over time while accounting for individual variability, providing robust evidence on the effectiveness of caregiver-focused educational interventions in stunting-related developmental care.

## Method

### Study Design

This study used a longitudinal quasi-experimental design with individual-level random allocation, conducted between January and November 2025. Participants were recruited from multiple community health centres and randomly assigned to intervention and control groups; however, due to the absence of full trial control procedures (e.g., blinding), the study is described as quasi-experimental. The intervention group received a structured developmental stimulation education programme, while the control group received routine

care. Maternal knowledge and caregiving practices were assessed at baseline, one month, three months, and six months after the intervention (Figure 1).

## Population and sample

The study included mothers of children with stunting registered at community health centres, defined as height-for-age below -2 standard deviations according to World Health Organisation growth standards. Eligible participants were primary caregivers who provided informed consent, while those with children with congenital anomalies, chronic illness, severe acute conditions, or multiple births (e.g., twins), as well as mothers with cognitive or severe mental health impairments, were excluded. A total of 87 eligible mother–child dyads were recruited and randomly allocated on an individual basis in a 1:1 ratio to the intervention ( $n = 44$ ) and control ( $n = 43$ ) groups, irrespective of community health centre. During follow-up, 7 participants were lost to follow-up, resulting in complete data for 40 participants in each group included in the final analysis.

## Instruments

Maternal knowledge and caregiving practices were measured using structured questionnaires developed for this study based on established child development and stunting-related care guidelines. Content validity was confirmed through expert review (I-CVI 0.83–1.00; S-CVI/Ave 0.92), and both instruments demonstrated good internal consistency (Cronbach's  $\alpha \geq 0.80$ ). The questionnaires assessed maternal knowledge of stunting and developmental stimulation, as well as caregiving practices across motor, language, cognitive, and social domains, and were administered at baseline and at one, three, and six months after the intervention.

## Data collection

Data were collected sequentially, beginning with baseline assessment following recruitment and eligibility screening. Mothers completed the questionnaires under the supervision of trained research assistants, after which the intervention group received structured developmental stimulation education delivered by trained facilitators using standardized materials. Follow-up assessments were conducted at one, three, and six months using identical procedures in both groups to ensure measurement consistency. Both groups continued to receive the same routine child health services, including growth monitoring and standard community-based care. The only difference between groups was the provision of the structured




Figure 1. Adapted CONSORT-style flow diagram of participant recruitment, allocation, follow-up and analysis

developmental stimulation education programme to the intervention group.

The educational intervention was delivered over four sessions within a one-month period, with each session lasting approximately 30–45 minutes. Sessions were conducted weekly by trained facilitators using interactive methods, including demonstrations, guided practice, and simple visual aids (e.g., flipcharts). The content focused on age-appropriate stimulation activities across motor, language, cognitive, and social domains, which could be integrated into daily caregiving routines. The sessions were delivered in the local language to ensure comprehension, and mothers were encouraged to practice the activities at home. No formal standardized external programme was adopted; however, the intervention was developed based

on established child development and early stimulation guidelines.

### Data analysis

Data were analysed using SPSS version 31, with descriptive statistics summarizing participant characteristics and outcome measures. Longitudinal changes in maternal knowledge and caregiving practices were examined using generalized linear mixed models with fixed effects for group, time (treated as a continuous variable coded as 0, 1, 3, and 6 months), and their interaction. The inclusion of both main effects and the interaction term followed the hierarchical modelling principle. A random intercept was specified to account for between-subject variability. Statistical significance was set at  $p < 0.05$ , with 95% confidence intervals reported where appropriate.

### Ethical approval

Ethical approval was obtained from an institutional health research ethics committee (No. 248.6/II.3.AU/F/KEPK/VIII/2024), and the study was conducted in accordance with the Declaration of Helsinki; written informed consent was obtained from all participants, with confidentiality and the right to withdraw assured.

## Results

Baseline characteristics of participants were comparable between the intervention and control groups (Table 1). Maternal age, child age, and educational levels were similar across groups, indicating no substantial baseline differences. As shown in Figure 1 (CONSORT flow diagram), 87 eligible mother–child dyads were recruited and randomly allocated to the intervention ( $n = 44$ ) and control ( $n = 43$ ) groups. Seven participants were lost to follow-up, resulting in complete data from 40 participants in each group at six months. These participants were included in the final longitudinal analysis.

Maternal knowledge scores increased over time in both groups, with greater and more sustained improvements observed in the intervention group compared to the control group (Table 2). Similarly, caregiving practices improved progressively in the intervention group, whereas changes in the control group were modest, indicating that the educational intervention was associated with concurrent improvements in maternal knowledge and caregiving practices (Table 3).

The mixed-effects analysis (Table 4) showed a statistically significant group-by-time interaction for both maternal

**Table 1. Baseline characteristics of participants**

| Variable                            | Intervention (n = 40) | Control (n = 40) |
|-------------------------------------|-----------------------|------------------|
| Maternal age (years), mean $\pm$ SD | 29.4 $\pm$ 5.8        | 28.9 $\pm$ 6.1   |
| Child age (months), mean $\pm$ SD   | 24.7 $\pm$ 8.3        | 25.1 $\pm$ 7.9   |
| Maternal education, n (%)           |                       |                  |
| – No formal education               | 8 (20.0)              | 9 (22.5)         |
| – Primary education                 | 18 (45.0)             | 17 (42.5)        |
| – Secondary or higher               | 14 (35.0)             | 14 (35.0)        |

**Table 2. Maternal knowledge scores by group and time**

| Time point | Intervention group (Mean $\pm$ SD) | Control group (Mean $\pm$ SD) |
|------------|------------------------------------|-------------------------------|
| Baseline   | 7.00 $\pm$ 0.00                    | 6.50 $\pm$ 0.51               |
| 1 month    | 7.60 $\pm$ 0.78                    | 7.18 $\pm$ 0.38               |
| 3 months   | 8.60 $\pm$ 0.84                    | 7.83 $\pm$ 0.71               |
| 6 months   | 10.00 $\pm$ 0.00                   | 7.83 $\pm$ 1.24               |

**Table 3. Maternal caregiving practices scores by group and time**

| Time point | Intervention group (Mean $\pm$ SD) | Control group (Mean $\pm$ SD) |
|------------|------------------------------------|-------------------------------|
| Baseline   | 6.00 $\pm$ 0.00                    | 5.90 $\pm$ 0.45               |
| 1 month    | 7.80 $\pm$ 0.79                    | 6.20 $\pm$ 0.41               |
| 3 months   | 9.00 $\pm$ 0.85                    | 6.80 $\pm$ 0.60               |
| 6 months   | 10.00 $\pm$ 0.00                   | 7.10 $\pm$ 0.74               |

**Table 4. Generalised linear mixed model analysis of maternal outcomes**

| Fixed effect                    | Estimate ( $\beta$ ) | SE   | 95% CI        | p-value |
|---------------------------------|----------------------|------|---------------|---------|
| <b>Maternal Knowledge</b>       |                      |      |               |         |
| Intercept                       | 6.82                 | 0.09 | 6.64 to 7.00  | <0.001  |
| Group (Intervention vs Control) | 0.24                 | 0.13 | -0.01 to 0.49 | 0.060   |
| Time                            | 0.20                 | 0.02 | 0.16 to 0.25  | <0.001  |
| Group $\times$ Time             | 0.29                 | 0.03 | 0.23 to 0.35  | <0.001  |
| <b>Caregiving Practices</b>     |                      |      |               |         |
| Intercept                       | 5.91                 | 0.11 | 5.69 to 6.13  | <0.001  |
| Group (Intervention vs Control) | 0.27                 | 0.14 | -0.01 to 0.55 | 0.058   |
| Time                            | 0.31                 | 0.03 | 0.25 to 0.37  | <0.001  |
| Group $\times$ Time             | 0.42                 | 0.04 | 0.34 to 0.50  | <0.001  |




Figure 2a and b. Estimated marginal means from the generalized linear mixed models showing trajectories of maternal knowledge and caregiving practices over time. Note the non-linear scale of the X-axes

knowledge and caregiving practices, indicating different trajectories between the intervention and control groups. Mothers in the intervention group demonstrated greater and more sustained improvements over time compared with the control group, even after accounting for individual variability.

Figure 2 presents the model-predicted trajectories derived from the generalized linear mixed models rather than raw observed means. The mixed-effects models indicated meaningful between-subject variability in maternal outcomes, supporting the inclusion of random intercepts to account for individual differences (Table 4). This confirmed the appropriateness of the generalized linear mixed modelling approach for analysing longitudinal changes over time. Although the theoretical maximum score for both outcome measures was 10, model-predicted values from the generalized linear mixed models are not mathematically constrained to the instrument's scale limits. Values exceeding ten in the graphical representation reflect extrapolated model estimates rather than observed raw scores.

## Discussion

This study demonstrated that a developmental stimulation education program produced sustained improvements in maternal knowledge and caregiving practices among mothers of children with stunting, consistent with previous evidence on the importance of caregiver-focused educational interventions.<sup>[3,10]</sup>

While earlier studies often relied on short-term or cross-sectional designs, the longitudinal findings of this study indicate that improvements in maternal knowledge were maintained over six months. These results suggest that structured caregiver education may support enduring cognitive and behavioural change rather than transient learning effects.<sup>[2,11]</sup>

The observed improvements in caregiving practices support previous evidence that caregiver knowledge can translate into behavioural change when educational interventions are practical, contextually relevant, and reinforced over time.<sup>[6,12]</sup> The significant group-by-time interaction demonstrated that the intervention altered the trajectory of maternal knowledge and caregiving practices rather than producing short-term effects, extending earlier findings by capturing longitudinal change and individual variability.<sup>[5,6,13]</sup> Overall, these findings corroborate existing literature by showing that structured developmental stimulation education can produce sustained improvements in maternal capacity, highlighting the importance of caregiver-focused strategies in stunting-related developmental care.<sup>[14,15]</sup>

## Clinical and health system implications

The findings highlight the clinical value of caregiver-focused educational interventions in improving maternal capacity to provide developmentally supportive care for children with stunting. Integrating developmental stimulation education into routine maternal and child health services, particularly through nurses and community health workers, may extend care beyond nutrition toward comprehensive developmental support. At the health system level, caregiver education represents a feasible and scalable strategy that can complement existing stunting programs and support early intervention efforts.

Beyond clinical implementation, these findings have broader implications for public health programmes and policy. Integrating caregiver education into community-based child health initiatives, including early childhood development programmes, may enhance coverage beyond health facilities. Strategies involving community health workers, home-based education, and integration

with existing child development services may improve accessibility and sustainability. Further operational research is needed to adapt and scale such interventions in resource-limited settings.

### Study limitations

This study has several limitations, including the quasi-experimental design without randomization and the use of self-reported measures, which may introduce residual confounding and social desirability bias. In addition, the study was conducted in a single community setting, potentially limiting the generalizability of the findings. Nevertheless, the longitudinal design, complete follow-up, and use of mixed-effects modelling strengthen confidence in the observed intervention effects.

### Conclusion

This study demonstrated that a developmental stimulation education program produced sustained improvements in maternal knowledge and caregiving practices among mothers of children with stunting over a six-month period. Integrating caregiver-focused developmental stimulation education into routine maternal and child health services may strengthen stunting management by enhancing the developmental environment of affected children. Further, it is important that early stimulation education of the mothers and caregivers be incorporated into nutritional rehabilitation programmes and become part of standard nutritional care protocols.

**Funding:** This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

**Author contributions:** N.I. conceptualized and designed the study, coordinated data collection, and contributed to manuscript drafting. A.O. provided methodological input, supervised the research process, and critically reviewed the manuscript for intellectual content. S.K. contributed to instrument development, data interpretation, and manuscript revision. P.A.W.S. performed data analysis, interpreted the statistical findings, and contributed substantially to manuscript writing and revision. All authors reviewed and approved the final version of the manuscript.

### Declaration of Generative AI and AI-assisted Technologies in the writing process

During the development of this manuscript, AI-assisted tools were utilized solely to support language refinement

and clarity. All generated content was carefully reviewed, revised, and validated by the authors, who assume full responsibility for the accuracy and integrity of the final manuscript.

**Conflicts of Interest:** The authors declare no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

**Ethics Approval:** The study was approved by the Ethics Committee of the Faculty of Health and Sciences, Universitas Muhammadiyah Gombong, Indonesia (Approval No. 248.6/II.3.AU/F/KEPK/VIII/2024). Written informed consent was obtained from all participants prior to study enrolment.

### References

1. Suryawan A, Jalaludin MY, Poh BK, Sanusi R, Tan VMH, Geurts JM, et al. Malnutrition in early life and its neurodevelopmental and cognitive consequences: a scoping review. *Nutr Res Rev*. 2022 Jun;35(1):136–49.
2. Sideropoulos V, Draper A, Munoz-Chereau B, Ang L, Dockrell JE. Childhood stunting and cognitive development: a meta-analysis. *J Glob Health*. 15:04257.
3. Upadhyay RP, Pathak BG, Raut SV, Kumar D, Singh D, Sudfeld CR, et al. Linear growth beyond 24 months and child neurodevelopment in low- and middle-income countries: a systematic review and meta-analysis. *BMC Pediatr*. 2024 Feb 8;24(1):101.
4. Gaidhane A, Telrandhe S, Holding P, Khatib MN, Patil M, Gaidhane S, et al. Impact of an integrated parenting and nutrition intervention on growth and development in stunted children at 24 months: evidence from the Stepping Stones programme in rural India. *BMJ Glob Health*. 2026, 11(1).
5. Serván-Mori E, Quezada-Sánchez AD, Fuentes-Rivera E, Pineda-Antunez C, Hernández-Chávez M del C, García-Martínez A, et al. Proximal determinants of suboptimal early child development during the first three years of life in socially deprived Mexican contexts. *PLOS ONE*. 2023 Nov 2;18(11):e0291300.
6. Crawford EJ, Crook EC, Waldbly L, Douglas P. New perspectives on responsive infant care: A qualitative study of the ways in which

- Neuroprotective Developmental Care (NDC) shapes mother-infant co-occupations. *J Occup Sci.* 2024 Apr 2;31(2):337–53.
7. Has EMM, Krisnana I, Efendi F. Enhancing Maternal Caregiving Capabilities Model to Prevent Childhood Stunting: A UNICEF-Inspired Model. *Sage Open Nurs.* 2024 Apr 1;10:23779608231226061.
8. Chapple A, Reynolds E, Mude A, Riaz W, Engle-Stone R. Is investing in food small and medium enterprises a viable path to improved nutrition in Sub-Saharan Africa? A multidisciplinary perspective. *Appl Econ Perspect Policy.* 2024;46(2):475–98.
9. Medise BE, Julia M, Devaera Y, Sitaresmi MN, Asmarinah, Widjaja NA, et al. Understanding the pubertal, psychosocial, and cognitive developmental trajectories of stunted and non-stunted adolescents: protocol of a multi-site Indonesian cohort study. *Front Pediatr.* 2024;12.
10. Ronaasen J, Wynne T, Biersteker L, Farrer W, Steenkamp L. The Opportunity for Nutrition Education to be a Mechanism for Climate Resilience: Experiences From a Facilitator-Led Programme in Diverse Contexts. *Australas J Early Child.* 2025 Dec 30;18369391251408649.
11. Bahari Gharehgoz A, Heidarabadi S, Alizadeh H, Asgari M. Effectiveness of Care for Child Development Program on the Sensitivity and Responsiveness Skills of Mothers. *Iran J Child Neurol.* 2022;16(1):51–63.
12. Yusriadi Y, Sugiharti S, Ginting YM, Sandra G, Zarina A. Preventing Stunting in Rural Indonesia: A Community-Based Perspective. *Afr J Food Agric Nutr Dev.* 2024.
13. Russell AL, Hentschel E, Fulcher I, Ravà MS, Abdulkarim G, Abdalla O, et al. Caregiver parenting practices, dietary diversity knowledge, and association with early childhood development outcomes among children aged 18-29 months in Zanzibar, Tanzania: a cross-sectional survey. *BMC Public Health.* 2022 Apr 15;22(1):762.
14. Black MM, Walker SP, Attanasio O, Rubio-Codina M, Meghir C, Hamadani JD, et al. Promoting Childhood Development Globally Through Caregiving Interventions. *Pediatrics.* 2023 May 1;151(Supplement 2):e2023060221B.
15. Do H, McCoy DC. The Relations Between Caregiver Education, Home Stimulation, and Children's Developmental Outcomes: Research in Majority World Countries. *Int J Early Child.* 2025 Apr 1;57(1):295–318.

## The missing evidence: anaemia, postpartum bleeding and maternal death

The WOMAN-2 study presents the missing evidence around anaemia and postpartum bleeding, advocating for decisive action to cut the prevalence and severity of anaemia in women and girls to reduce maternal deaths.

Maternal mortality remains one of the most pressing challenges in global health. Progress has slowed – particularly in sub-Saharan Africa and South Asia, where most maternal deaths occur.

Now, the WOMAN-2 study provides new evidence which suggests anaemia may be the hidden factor driving maternal deaths.

Bringing together new data and powerful personal accounts from women, the report calls for global action to reduce the prevalence and severity of anaemia in women and adolescent girls, with the ultimate aim of reducing maternal deaths.

<https://thebloodtrials.org/the-missing-evidence/>

# Holding the line: lived experiences of mental health, moral injury, and organizational support among public-health professionals in the Rohingya Camp, Bangladesh — a phenomenological study

Md. Ruhul Amin 

## Author Affiliation:

Department of Public Administration,  
Comilla University, Cumilla,  
Bangladesh

## Correspondence:

[rubel@cou.ac.bd](mailto:rubel@cou.ac.bd)

**Submitted:** January 2026

**Accepted:** March 2026

**Published:** May 2026

**Citation:** Amin. Holding the line: lived experiences of mental health, moral injury, and organizational support among public-health professionals in the Rohingya Camp, Bangladesh — a phenomenological study. *South Sudan Medical Journal*, 2026;19(2):119-125  
© 2026 The Author (s) **License:** This is an open access article under [CC BY-NC](https://creativecommons.org/licenses/by-nc/4.0/) DOI: <https://dx.doi.org/10.4314/ssmj.v19i2.9>

## ABSTRACT

**Introduction:** Public-health professionals in the Rohingya camps work amid chronic scarcity and ethical tension. This study examined their lived experiences of mental health, moral injury, and organizational support to identify practical, context-fit improvements.

**Method:** Qualitative hermeneutic phenomenology with fifteen semi-structured interviews across cadres. Data were audio-recorded, transcribed/translated, and analysed using a reflexive thematic analysis informed by a hybrid deductive–inductive approach, drawing on Job Demands–Resources, Conservation of Resources, moral-injury, and Perceived Organizational Support lenses. Trustworthiness was supported by analyst triangulation, an audit trail, and COREQ (Consolidated Criteria for Reporting Qualitative Research) guidance.

**Results:** Nine themes clustered into four domains: (1) demand–resource imbalance with spillover beyond shifts (sleep disruption, rumination) and in-flow micro-regulation; (2) moral injury under scarcity and constrained agency (stock-outs, early discharges, bureaucratic delays, indicator dominance); (3) organizational support as felt (reflective supervision, buddy rotations/micro-check-ins, protected micro-breaks, confidential on-roster counselling); and (4) policy–context fit (offline-first data entry, stable templates, locally empowered approvals).

**Conclusions:** Low-friction, context-fit methods can break loss spirals, reduce moral-injury assessments, and maintain safe, compassionate care, making staff well-being a design challenge.

**Keywords:** demands–resources, job, moral injury, phenomenology, public-health professionals, Rohingya.

## Introduction

Prolonged humanitarian crises in Cox's Bazar expose health workers to extreme stress, ethical challenges, and resource scarcity.<sup>[1,2]</sup> Studies show humanitarian and

health workers face high burnout and secondary trauma due to continuous exposure to intense suffering, often mirroring distress experienced by displaced populations.<sup>[3]</sup> Recent studies show persistent anxiety and depression among workers, highlighting the need for systemic, context-sensitive interventions beyond individual resilience training.<sup>[4]</sup> Within mental health discourse, moral injury denotes psychological harm from witnessing or engaging in actions that violate one's moral values. Moral injury describes ethical distress from dilemmas or betrayal, extending beyond burnout to explain suffering not rooted solely in exhaustion.<sup>[5]</sup>

Supportive workplaces reduce ethical harm through supervision, peer support, and accessible counselling, strengthening staff resilience and long-term sustainability.<sup>[6]</sup> In under-resourced camp settings, well-being policies rarely translate into protected time, adequate support, or safe reporting, exposing gaps between policy and practice.<sup>[7]</sup>

In the Rohingya response, demand surges, infrastructure disruptions, and complex referrals often weaken the implementation of well-intended policy frameworks.<sup>[8]</sup> Conceptual clarity matters because organizational support includes diverse elements whose effects vary across roles, tasks, and work contexts. This study used phenomenology, the JD-R (Job Demands–Resources), and moral-injury frameworks to explore how public health professionals interpret their daily routines, ethical tensions, and personal experiences of burden and coping.<sup>[9]</sup> JD-R explains how high emotional and cognitive demands, combined with limited resources and support, drive either strain or engagement. Moral-injury frameworks link value conflicts and perceived transgressions to lasting psychological harm, guiding organizations to ethically support affected individuals.<sup>[10]</sup> Together, these lenses allow the study to connect subjective experience with modifiable features of work design and organizational climate.

This study also examined how public health professionals in Rohingya camps experience and manage work-related strain, ethical tensions, and organizational support. The JD-R model links high demands—caseloads, trauma, time pressure—to engagement or strain.<sup>[11]</sup> COR (Conservation of Resources) showed resource loss fuels further depletion, while small, consistent gains help mitigate these effects. POS (Perceived Organizational Support) connects individual experiences with organizational climate, shaping whether employees perceive policies as meaningful, supportive, and accessible during high-demand periods.

This phenomenological study explored Rohingya camp

health workers' mental health challenges, moral injury, and perceptions of organizational support, providing evidence to inform staff well-being policies and to improve humanitarian organizational practices.

### Method

In this study, a qualitative, hermeneutic phenomenological design was used to elicit and interpret the lived meanings of public health professionals. Ethical approval was obtained (Approval Code: ERCP-2026-0413-06), which guaranteed voluntary participation, informed consent, confidentiality, and anonymity. The participants had the freedom to drop out at any time. The researchers respected the accepted ethical principles of conducting research on human subjects.

Reporting was done according to COREQ (Consolidated Criteria for Reporting Qualitative Research) guidelines where necessary. Purposeful, maximum-variation sampling was employed to reflect heterogeneity across roles [e.g., public-health nursing, EPI (Expanded Programme on Immunization-focused on vaccinating children), surveillance, M&E (Monitoring and Evaluation), MHPSS (Mental Health and Psychosocial Support)/health promotion] and across organization types (government, INGO/NGO/UN). Purposeful sampling prioritizes theoretical relevance and rich data over representativeness; information power guides sample size, with fewer interviews needed for credible, detailed insights.<sup>[12]</sup> Conducted fifteen one-to-one interviews, meeting adequacy criteria, aligning with evidence that thematic patterns often stabilize within the first dozen interviews. Inclusion required ≥6 months in the Rohingya response, consent capacity, and diverse roles, sexes, and organizations; data were analysed using reflexive, hermeneutic thematic analysis. Used a mixed deductive–inductive approach to conduct the study. Two analysts double-coded 20–25% of transcripts, resolving discrepancies collaboratively. The remaining transcripts were single-coded and peer-reviewed for complex sections. Trustworthiness was ensured through triangulation, audit trails, reflexivity, COREQ-aligned reporting, and detailed contextual descriptions to support transferability.<sup>[13]</sup>

### Results

#### Participant characteristics

Fifteen public-health professionals participated, spanning public-health nursing, EPI, surveillance, M&E, and MHPSS/health-promotion roles across government,

**Table 1. Demographic characteristics of the respondents**

| <b>Id</b> | <b>Age</b> | <b>Sex</b> | <b>Cadre/Role</b>         | <b>Org Type</b> | <b>Duty Station (Code)</b>       | <b>Time in current role</b> | <b>Typical schedule</b>               |
|-----------|------------|------------|---------------------------|-----------------|----------------------------------|-----------------------------|---------------------------------------|
| 1         | 29         | Female     | PHN (Public Health Nurse) | Govt            | Camp 1E Health Post              | 2.5 yrs                     | 6 days/week; rotating day/evening     |
| 2         | 34         | Male       | EPI Focal Point           | Govt            | Camp 3 Clinic                    | 4 yrs                       | 5.5 days/week; campaign OT            |
| 3         | 31         | Female     | Surveillance Officer      | INGO            | Camp 5 Primary Health Care       | 3 yrs                       | 6 days/week; on-call outbreaks        |
| 4         | 27         | Female     | MHPSS                     | NGO             | Camp 7 Community Centre          | 1.5 yrs                     | 5 days/week                           |
| 5         | 36         | Male       | M&E Officer (Health)      | UN              | Camp 9 PHC                       | 5 yrs                       | 5 days/week; field visits 2x/week     |
| 6         | 30         | Female     | PHN (Nurse)               | INGO            | Camp 10 PHC                      | 2 yrs                       | 6 days/week; rostered weekends        |
| 7         | 33         | Male       | Surveillance Officer      | Govt            | Camp 12 Health Post              | 3.5 yrs                     | 6 days/week; night alerts possible    |
| 8         | 28         | Female     | MHPSS                     | UN              | Camp 14 Women-F Centre           | 2 yrs                       | 5.5 days/week; outreach shifts        |
| 9         | 35         | Male       | EPI Focal Point           | NGO             | Camp 16 EPI Centre               | 4.5 yrs                     | 6 days/week; campaign nights          |
| 10        | 32         | Female     | M&E Officer (Health)      | INGO            | Camp 18 Field Office             | 3 yrs                       | 5 days/week; monitoring missions      |
| 11        | 29         | Male       | Surveillance Officer      | UN              | Camp 20 Health Post              | 2 yrs                       | 6 days/week                           |
| 12        | 26         | Female     | PHN (Nurse)               | NGO             | Camp 22 PHC                      | 1.2 yrs                     | 6 days/week; alternating weekends     |
| 13        | 38         | Male       | EPI Focal Point           | Govt            | Camp 24 EPI Hub                  | 6 yrs                       | 5.5 days/week; campaign surge         |
| 14        | 30         | Female     | MHPSS                     | INGO            | Camp 27 Community Hub            | 2.8 yrs                     | 5 days/week; evening youth sessions   |
| 15        | 37         | Male       | M&E Officer (Health)      | Govt            | <i>Bhasan Char</i> Health Sector | 4 yrs                       | 5 days/week; monthly island rotations |

**Table 2. Thematic analysis**

| Participant ID | Theme                                        | Code                               | Excerpt (abridged)                                                   |
|----------------|----------------------------------------------|------------------------------------|----------------------------------------------------------------------|
| P07            | Chronic vigilance & emotional carryover      | Sleep disturbance & hypervigilance | "I wake at 3 a.m. thinking about the patient I sent home..."         |
| P12            |                                              | Cognitive carryover                | "Even at home the ward is in my head—voices, forms, queues."         |
| P03            | Momentary regulation in high-tempo care      | Micro-resets between cases         | "Three slow breaths and a sip of water—one minute to reset."         |
| P10            |                                              | Peer micro-check-ins               | "A two-minute 'you, okay?' from my buddy helps..."                   |
| P05            | Operational pressures & sense-making         | Queue-induced triage tension       | "When the waiting area swells...I start fast-tracking..."            |
| P09            |                                              | Documentation pressure             | "Midday the registers win—I'm split between form and patient."       |
| P01            | Ethical friction at the interface            | Resource-constrained refusals      | "'Today we cannot provide this' sits heavy—did I fight hard enough?" |
| P06            |                                              | Early discharge for capacity       | "Discharging early to free a bed felt like betrayal."                |
| P11            | Systemic bottlenecks & bureaucratic drag     | Protocol vs urgency                | "We needed an alert but waited for authorizations."                  |
| P02            |                                              | Procurement/approval delays        | "Stuck between patients and purchase approvals."                     |
| P13            | Metrication & narrative loss                 | Data–human story dissonance        | "The dashboard wants numbers; my shift is stories."                  |
| P04            |                                              | Supportive supervision             | "'How are you coping?' and rotation... I actually sleep."            |
| P08            | Scaffolded support & containment             | Protected micro-breaks             | "A break is real when someone covers me... 20 minutes."              |
| P14            |                                              | Unclear escalation ladder          | "Without a clear referral ladder, every refusal becomes a fight..."  |
| P15            | Policy–context fit & implementation fidelity | Policy–context misfit              | "Power cuts make uploads fail... re-enter late at night—exhausting." |

INGO/NGO, and UN facilities in the Rohingya camps (Cox's Bazar, Bangladesh) – Table 1.

### Thematic Analysis

Nine interlocking themes described how workers experience (1) mental health in daily camp work, (2) job demands and moral injury, and (3) organizational enablers/barriers (Table 2).

### Chronic vigilance and emotional carryover

Participants reported lingering hyperarousal beyond shifts, marked by disrupted sleep, intrusive recollections of ethical dilemmas, somatic tension, and intensified distress following surge periods or complex clinical decisions.

*"Sleep is never deep after a heavy clinic. I wake at 3 a.m. thinking about the patient I sent home—did I miss something?"* (P07)

*"Even at home the ward is in my head—voices, forms, the queue moving too fast. I slow my breathing, but scenes keep replaying."* (P12)

These accounts reflect a demand–resource imbalance (JD-R) with inadequate recovery opportunities, producing work–home spillover. The pattern aligns with resource-loss spirals (COR)—loss of sleep/time/psychological safety begets further loss. Comparable elevations in distress among humanitarian/health workers are reported in Bangladesh and allied settings.<sup>[14]</sup>

### Momentary regulation in high-tempo care

Staff narrated on-shift micro-regulation—brief breathing routines, one-minute pauses between patient clusters, micro-breaks with water/stretching—and boundary rituals (shower/journaling) to contain spillover. Peer micro-check-ins ("you, okay?") and task-swaps ("I'll finish the register if you handle the referral") were repeatedly described as small, usable buffers.

*"Before the next cluster I do three slow breaths, sip water—one minute to reset my head."* (P03)

*"A two-minute 'you, okay?' from my buddy helps more than a long meeting. It tells me I'm not alone."* (P10)

These micro-resources help interrupt loss spirals (COR) and partially rebalance the JD-R equation in real time. Their salience as felt supports dovetails with evidence that perceived organizational support (POS) via supervisor care, peer architecture, and protected recovery moments—attenuates strain and improves role sustainability.<sup>[15]</sup> In the Rohingya response, peer-based coping is consistent

with regional reports of socially embedded support in constrained health systems.

### Operational pressures shaping sense-making

Care delivery faced patient surges, documentation demands, and system failures, causing rapid triage, physical strain, and divided attention.

*"When the waiting area swells, I start fast-tracking, but my body is tense—like I'm missing something while keeping the line moving."* (P05)

*"Midday the registers win. I'm split between the form and the patient—both feel urgent."* (P09)

Dual clinical-reporting demands amplify cognitive load, worsening perceived quality and post-shift carryover—again a JD-R configuration of high demands with thin resources.<sup>[16]</sup> When reporting templates change mid-surge or systems lack offline resilience, workers experience "policy as friction," a mechanism linked to moral-distress appraisals in crisis care. Frontline accounts from Bangladesh similarly show how operational constraints interact with ethics and workload to shape mental-health outcomes.<sup>[17]</sup>

### Policy context fit & implementation fidelity

Participants contrasted policies "on paper" with whether they are workable under outages, surges, and staffing constraints.

*"The policy sounds good, but with power cuts the uploads fail, and we re-enter late at night—exhausting."* (P15)

*"Changing the form mid-campaign created chaos. Keep it stable and offline-first; then policy helps, not hurts."* (P08)

Poor implementation fit amplifies cognitive load (JD-R), drains scarce resources (COR), and fuels ethical strain when compliance competes with care. Guidance that ignores field constraints risks epistemic and practical misalignment in humanitarian settings. Participatory, co-designed adjustments stable templates, offline-first entry, scheduled micro-breaks with cover—are supported by implementation literature in mental-health systems redesign.<sup>[18]</sup>

### Discussion

This phenomenological study illuminates how public-health professionals in the Rohingya camps experience a persistent demand–resource imbalance that carries beyond the working shift (Themes 1–3), a pattern of moral injury

under scarcity and constraints (Themes 4–6), and a gap between policies on paper and supports as felt (Themes 7–9). Workers experienced disrupted sleep and rumination from ethical challenges, relying on peer support and in-the-moment coping, while systemic pressures worsened distress; structured support, micro-breaks, buddy systems, clear escalation, and confidential counselling proved protective, addressing daily mental-health experiences and organizational interventions. They also align with the study's integrated theoretical lens (phenomenology, JD-R, COR, moral injury, and POS).

Firstly, the work–home spillover and “always-on” vigilance map onto JD-R predictions that chronic emotional/cognitive demands with thin recovery resources generate strain and reduced engagement. Participants' descriptions of broken sleep, somatic tension, and intrusive replay also exemplify COR “loss spirals,” in which depletion of time, sleep, and psychological safety makes subsequent strain more likely. Comparable elevations in humanitarian/health worker distress have been reported in Bangladesh and allied crisis contexts, supporting external validity.<sup>[19]</sup>

Secondly, the data differentiate moral injury from burnout, showing guilt, shame, and self-doubt stem from morally injurious events, such as care denial or procedural delays, aligning with research linking value-violation distress to constrained agency. Findings show operational issues—surge changes, online-only systems, unclear escalation—amplify moral residue among refugee-camp public health workers.

Thirdly, policy-context mismatch hinders implementation; effective execution demands participatory design, offline workflows, stable templates, and empowered local decision-making. Findings indicate that seemingly minor design decisions may produce disproportionate influence on perceived quality, moral stress, and recovery.

### Conclusion

This study shows Rohingya camp health workers face chronic demand–resource imbalance and moral injury, but practical buffers—reflective supervision, buddy rotations, micro-breaks, clear referrals, and confidential counselling—help mitigate stress and support well-being. Implementation fidelity—stable templates, offline-first entry, and local approvals—is crucial to ensuring effective, low-cost support, reducing moral injury, and maintaining safe, humane care. Future research should test these adjustments through implementation evaluations and mixed-methods designs across cadres and settings.

**Conflict of Interest:** None

**Funding Sources:** None.

### References

1. Chowdhury SA, McHale T, Green L, Mishori R, Pan C, Fredricks I. Health professionals' perspectives on the impact of COVID-19 on sexual and gender-based violence (SGBV) and SGBV services in Rohingya refugee communities in Bangladesh. *BMC health services research*. 2022 Jun 4;22(1):743. DOI: <https://doi.org/10.1186/s12913-022-08122-y>
2. Sheather J, Apunyo R, DuBois M, Khondaker R, Noman A, Sadique S, McGowan CR. Ethical guidance or epistemological injustice? The quality and usefulness of ethical guidance for humanitarian workers and agencies. *BMJ global health*. 2022 Mar 1;7(3):e007707. DOI: <https://doi.org/10.1136/bmjgh-2021-007707>
3. Burton CW, Jenkins DK, Chan GK, Zellner KL, Zalta AK. A mixed methods study of moral distress among frontline nurses during the COVID-19 pandemic. *Psychological Trauma: Theory, Research, Practice, and Policy*. 2024 May;16(4):568. DOI: <https://doi.org/10.1037/tra0001493>
4. Conway P, Redmond T, Lundrigan S, Davy D, Bailey S, Lee P. Protecting the protectors: moral injury, coping styles, and mental health of UK police officers and staff investigating child sexual abuse material. *Depression and Anxiety*. 2024;2024(1):1854312. DOI: <https://doi.org/10.1155/da/1854312>
5. Agboh HN, Adjei GA, Okai GA. Psychological stressors among faith-based health workers during Marburg virus outbreak in Ghana: A qualitative study. *PLOS Mental Health*. 2025 May 29;2(5):e0000042. DOI: <https://doi.org/10.1371/journal.pmen.0000042>
6. Harris H, Clarkin C, Rovet J, Crawford A, Johnson A, Kirvan A, Gruszecki S, Wang S, Soklaridis S. Meaningful engagement through critical reflexivity: Engaging people with lived experience in continuing mental health professional development. *Health Expectations*. 2023 Oct;26(5):1793–8. DOI: <https://doi.org/10.1111/hex.13798>

7. Forrester-Bowling T, Lucas JJ, Brown AD, Bennetts S, Carolin R, Hayward J, Scott D, Peeters A, McLure J. Adapting Group Model Building for Mental Healthcare: A Participatory Co-Design Approach. *International Journal of Mental Health Nursing*. 2025 Feb;34(1):e13451. DOI: <https://doi.org/10.1111/inm.13451>
8. Parvin T, Rosenbaum S, Ozen S, Ewagata L, Ventevogel P. Mental health and perceived social support of humanitarian workers in Bangladesh during the COVID-19 pandemic. *Intervention Journal of Mental Health and Psychosocial Support in Conflict Affected Areas*. 2022 Jul 1;20(2):151-60. DOI: [https://doi.org/10.4103/intv.intv\\_18\\_21](https://doi.org/10.4103/intv.intv_18_21)
9. Häger M, Boman E, Forsman AK. Meeting the Needs of Older Adults with Mental Ill-Health in Non-Psychiatric Care Settings: Self-Rated Confidence in Helping and its Co-Variates within a Multiprofessional Study Sample. *Gerontology and Geriatric Medicine*. 2023 Jun;9:23337214231179819. DOI: <https://doi.org/10.1177/23337214231179819>
10. McPhee J, Warner T, Cruwys T, Happell B, Scholz B. 'They don't really know why they're here' mental health professionals' perspectives of consumer representatives. *International Journal of Mental Health Nursing*. 2023 Jun;32(3):819-28. DOI: <https://doi.org/10.1111/inm.13124>
11. Bakker AB, Demerouti E, Sanz-vergel A. Job Demands-Resources Theory: Ten Years Later. 10, 25-53. DOI: <https://doi.org/10.1146/annurev-orgpsych-120920-053933>
12. Malterud K, Siersma VD, Guassora AD. Sample size in qualitative interview studies: guided by information power. *Qualitative health research*. 2016 Nov;26(13):1753-60. DOI: <https://doi.org/10.1177/1049732315617444>
13. Nowell LS, Norris JM, White DE, Moules NJ. Thematic analysis: Striving to meet the trustworthiness criteria. *International journal of qualitative methods*. 2017 Sep 28;16(1):1609406917733847. DOI: <https://doi.org/10.1177/1609406917733847>
14. D'Alessandro AM, Ritchie K, McCabe RE, Lanius RA, Heber A, Smith P, Malain A, Schielke H, O'Connor C, Hosseiny F, Rodrigues S. Healthcare workers and COVID-19-related moral injury: an interpersonally-focused approach informed by PTSD. *Frontiers in psychiatry*. 2022 Feb 14;12:784523. DOI: <https://doi.org/10.3389/fpsyt.2021.784523>
15. Happell B, Warner T, Waks S, O'Donovan A, Manning F, Doody R, Greaney S, Goodwin J, Hals E, Griffin M, Scholz B. Something special, something unique: Perspectives of experts by experience in mental health nursing education on their contribution. *Journal of Psychiatric and Mental Health Nursing*. 2022 Apr;29(2):346-58. DOI: <https://doi.org/10.1111/jpm.12773>
16. Järvinen M, Kessing ML. Self-casting and alter-casting: Healthcare professionals' boundary work in response to peer workers. *Current Sociology*. 2023 May;71(3):414-31. DOI: <https://doi.org/10.1177/00113921211048532>
17. Wright M, Proeve M, Everitt CM, Bartsch DR. An Evaluation of co-Delivered Foundational BPD Training With Health Service Staff. *Personality and Mental Health*. 2025 May;19(2):e70016. DOI: <https://doi.org/10.1002/pmh.70016>
18. Horgan A, Manning F, Donovan MO, Doody R, Savage E, Bradley SK, Dorrity C, O'Sullivan H, Goodwin J, Greaney S, Biering P. Expert by experience involvement in mental health nursing education: the co-production of standards between experts by experience and academics in mental health nursing. *Journal of Psychiatric and Mental Health Nursing*. 2020 Oct;27(5):553-62. DOI: <https://doi.org/10.1111/jpm.12605>
19. Chatwiriyaiphong R, Fernandez R, Bosworth R, Kinghorn G, Moxham L. Conceptualisation of Mental Health Recovery by Health Professionals and Students in Southeast Asia: A Qualitative Systematic Review and Meta-Aggregation. *Journal of Psychiatric and Mental Health Nursing*. 2025 Jun;32(3):783-99. DOI: <https://doi.org/10.1111/jpm.13158>

# Endoscopic endonasal repair of post-traumatic sphenoid sinus cerebrospinal fluid leak with encephalocele

Justin Rubena<sup>1</sup>, Khalid Coco<sup>2</sup>, Francis Swaka<sup>3</sup>, Othwonh Twong<sup>3</sup>

### Author Affiliation:

1. Department of Ear, Nose and Throat, University of Juba and Juba Teaching Hospital, South Sudan
2. School Of Medicine, Ahfad University for women, Omdurman and Royal international Hospital, Khartoum, Sudan
3. Juba Teaching Hospital, Juba, South Sudan

### Correspondence:

Justin Rubena

[lugela4@yahoo.com](mailto:lugela4@yahoo.com)

**Submitted:** November 2025

**Accepted:** March 2026

**Published:** May 2026

**Citation:** Rubena et al Endoscopic endonasal repair of post-traumatic sphenoid sinus cerebrospinal fluid leak with encephalocele. South Sudan Medical Journal, 2026;19(2):126-129 © 2026 The Author (s) **License:** This is an open access article under [CC BY-NC](https://creativecommons.org/licenses/by-nc/4.0/) DOI: <https://dx.doi.org/10.4314/ssmj.v19i2.10>

### ABSTRACT

While cerebrospinal fluid (CSF) rhinorrhoea due to a road traffic accident is uncommon, its occurrence is clinically important because of the serious complications involving the central nervous system, such as meningitis. Often, the management of a post-traumatic CSF leak requires a multidisciplinary strategic approach. We report a case of post-traumatic sphenoid-sinus CSF leak with meningocele successfully repaired using a transnasal endoscopic multilayer technique in a setting with scarce resources. The initial neurosurgical intervention consisted of decompression of a massive pneumocephalus, followed by a definitive endoscopic skull-base repair. This procedure resulted in the cessation of the CSF leak and complete recovery. This case report highlights what can be achieved with a multi-disciplinary approach in a resource-constrained setting, such as South Sudan.

**Key words:** cerebrospinal rhinorrhoea, sphenoid sinus fracture, meningocele, endoscopic endonasal repair, resource-limited

### Introduction

Traumatic cerebrospinal fluid (CSF) leak is a recognized complication of head injury. It occurs in 1–3 % of closed-head traumas and 10–30 % of basilar-skull fractures.<sup>[1]</sup> Traumatic CSF rhinorrhoea accounts for about 90% of all CSF leaks, with 80% caused by motor-vehicle accidents or falls.<sup>[2]</sup> Accurate identification of the defect site and the presence of associated meningo-encephalocele are crucial before surgery. Localization can be aided by imaging and intra-operative fluorescein, though the sphenoid sinus remains a particularly challenging site. Persistent leaks lasting two to three weeks or more typically require surgical intervention, most effectively managed through an endoscopic endonasal approach.<sup>[3,4]</sup> We present a case of post-traumatic sphenoid-sinus CSF leak with encephalocele, managed collaboratively by ENT and Neurosurgery teams in Juba, South Sudan, demonstrating that complex skull-base repairs are achievable even in resource-limited environments.

### Case Report

On 14th June 2025, a 37-year-old male sustained a severe head injury following a fall and presented three weeks later with a persistent clear nasal discharge




Figure 1. CT scan sagittal view of massive pneumocephalus (Credit: Dr Justin Rubena Luamaya)




Figure 2. CT scan showing meningocele (hernia of the meninges and brain tissue through the lateral sphenoid sinus wall defect from the middle cranial fossa). (Credit: Dr Justin Rubena Luamaya)

from the left nostril and a constant severe headache. He was assessed clinically by a Consultant Neurosurgeon followed by CT imaging revealing a small fracture of the sphenoid bone with marked intracranial air collection - pneumocephalus (Figure 1). The CT scan in an axial view showed herniation of a meningocele from the left middle cranial fossa into the sphenoid sinus (Figure 2).

A burr hole was performed, with insertion of a drain to decompress the pneumocephalus, which led to gradual neurological stabilization and a reduction in headache. However, CSF rhinorrhoea persisted and one month later (24th July) rhinoendoscopy revealed, upon Valsalva manoeuvre, a gush of CSF from the left sphenoid sinus. CT imaging confirmed a posterolateral sphenoid-sinus defect with associated meningocele herniation, and endoscopic endonasal multilayer reconstruction was performed on the 2nd August.

A zero-degree scope (a rhinoscope with a camera allowing visualisation of the whole nasal cavity) was used to navigate through the nasal cavity alongside the medial attachment of the left superior turbinate to visualize the sphenoid opening. Blakesley forceps were used to do an uncinctomy (procedure to clear the uncinate complex) to expose the posterior rim of the maxillary ostium

(Figure 3). During this exposure of the mucosa, a branch of the sphenopalatine artery was severed, but bleeding was controlled using bipolar cautery. The anterior wall of the sphenoid was removed using a Kerrison Rongeurs instrument. This exposed the CSF leak site and the meningocele, which was seen protruding and pulsating from the lateral sphenoid wall.

Approximately 80 % of the anterior sphenoid-sinus wall was removed to expose the lateral aspect of the defect. A bipolar cautery was used to shrink the herniated meningocele and delineate the defect.

Multilayer reconstruction was performed as follows:

1. Underlay tragal cartilage for structural support.
2. Autologous abdominal fat to obliterate dead space and seal the defect (Figure 4).
3. Surgicel and Gelfoam for additional reinforcement.
4. A vascularized naso-septal flap was rotated over the defect (Figure 5).
5. Additional layers of Surgicel, bone wax, and tetracycline-soaked gauze were applied to reinforce the closure.




Figure 3. Removing the uncinate process to expose the posterior rim of the maxillary ostium and the medial pterygoid plate to access proximity to the lateral wall of the sphenoid. (Credit: Dr Justin Rubena Luamaya)



Figure 4. Horizontal incisional site (6cm) to harvest the abdominal fat graft after seven postoperative days. (Credit: Dr Justin Rubena Luamaya)



Figure 5. Placement of the vascularized naso-septal flap into the sphenoid defect. (Credit: Dr Justin Rubena Luamaya)

The bone wax and tetracycline gauze were removed four weeks post-operatively.

Post-operative recovery was uneventful. There was a complete cessation of CSF rhinorrhoea, without recurrence during follow-up, and no development of meningitis or pneumocephalus.

### Discussion

Traumatic CSF leaks of the sphenoid sinus are rare but potentially life-threatening because of the risks of central nervous system complications like meningitis, recurrent pneumocephalus, and neurological deficits.<sup>[5]</sup> Initial conservative management is the standard management in selected cases. However, persistent CSF leaks of more than 2-3 weeks require surgical intervention.<sup>[6]</sup> The endoscopic endonasal approach is superior to transcranial methods because it provides a direct view of the skull base and, furthermore, is associated with lower morbidity and higher success rates.<sup>[7]</sup> To ensure a watertight seal after reconstruction, a multilayer technique using autologous cartilage, fat, and vascularized mucosal flaps is preferred. This approach also minimizes recurrence.<sup>[8,9,10]</sup>

In our case, the initial neurosurgical intervention involved burr-hole decompression with a drain. This resolved the pneumocephalus and stabilized the patient for safe endoscopic repair. This staged, collaborative strategy demonstrates that even in resource-limited environments, advanced skull-base surgery can be successfully achieved with appropriate expertise and teamwork.

### Conclusion

Persistent post-traumatic CSF rhinorrhoea of the sphenoid sinus requires timely multidisciplinary management. The combination of neurosurgical decompression and endoscopic multilayer closure provides an effective, safe, and feasible solution, even in low-resourced African contexts. This case emphasizes the importance of enhancing ENT and neurosurgical collaboration and infrastructure in sub-Saharan Africa to manage complex skull base pathologies and prevent severe complications, such as meningitis and recurrent pneumocephalus.

**Authors' contributions:** RJ performed the endoscopic endonasal repair and drafted the manuscript. KC performed the initial neurosurgical evaluation and burr-hole decompression, and contributed to multidisciplinary planning. Others assisted during the surgical procedures.

**Acknowledgment:** The authors thank the staff of Juba

Teaching Hospital for their assistance and the patient and family for their cooperation and consent to publication.

## References

1. Ommaya AK et al. Cerebrospinal fluid rhinorrhea: Clinical observations and management. *Neurosurgery*. 1983;12(3):332–339.
2. Schlosser RJ, Bolger WE. Nasal CSF leaks: Critical review and surgical considerations. *Laryngoscope*. 2004;114(2):255–265.
3. Shetty PG et al. Evaluation of CSF leaks: High-resolution CT and MR cisternography correlation. *AJNR Am J Neuroradiology*. 1998; 19:633–639.
4. Hegazy HM, Carrau RL et al. Endoscopic repair of CSF rhinorrhea: A ten-year experience. *Laryngoscope*. 2000; 110:1166–1172.
5. Bernal-Sprekelsen M et al. Endoscopic management of CSF leaks. *Rhinology*. 2008; 46:92–96.
6. Tabae A et al. Endoscopic management of spontaneous CSF rhinorrhea: Predictors of success. *Otolaryngology Head Neck Surg*. 2009; 141:689–694.
7. Hadad G et al. A novel reconstructive technique after endoscopic expanded endonasal approaches: The vascularized Naso septal flap. *Laryngoscope*. 2006; 116:1882–1886.
8. Wormald PJ. *Endoscopic sinus surgery*. Stuttgart, New York: Thieme. 2013;5:28-44
9. Shimizu A, Makihara S, Imoto R, Hirashita K, Ando M. Endoscopic transpterygoid repair of sphenoid sinus meningocele: A comprehensive case report and literature review. *Auris Nasus Larynx* 2024; 51(5):875-879.
10. Alam S, Ahmed N, Joy SS, Manzoor MT, Rahad-Ul-Islam M, Rahman A, Chaurasia B. A rare case of cerebrospinal fluid rhinorrhea via Sternberg canal: a case report with endoscopic endonasal repair. *Annals of Medicine and Surgery*. 2025 Oct 1;87(10):6779-6784

## End TB

The World Health Organization (WHO) is urging countries to accelerate action to end tuberculosis (TB) and expand access to lifesaving services by using new innovations such as diagnostic tests that can be used near the point-of-care and tongue swabs that can help detect the disease faster reaching more people.

‘The new guidelines on tests for TB that can be used near the point-of-care, issued by WHO, mark another step towards faster detection and treatment of one of the world’s deadliest infectious diseases. These portable, simple-to-use tests bring TB diagnosis closer to where people routinely seek care. Available at less than half the cost of many existing molecular diagnostics, they can help countries expand access to testing. The tests can operate on battery power and deliver results in less than one hour, allowing patients to start treatment sooner...’

Guidelines: <https://www.who.int/teams/global-programme-on-tuberculosis-and-lung-health/diagnosis-treatment-npoc-tongue-swabs-and-sputum-pooling-for-tb>

Extract from [HIFA - Healthcare Information For All](#).

# A rare pathological feature of portal vein thrombosis complicated by portal hypertension and haematemesis

Hamidine Illa<sup>1</sup>, Ganiou Tidjani Kabirou<sup>2</sup>, Amadou Magagi Mahamane Ibrahim<sup>3</sup>, Maman Brah Moustapha<sup>4</sup>, Soumaila Abdou<sup>5</sup>, Habou Garba Abdoul Malik<sup>6</sup>, Adamou Harissou<sup>3</sup>

### Author Affiliation:

1. Department of Hepato-Gastroenterology, Zinder National Hospital, Faculty of Health Sciences, André Salifou University of Zinder, Niger, Clinical Research and Health System Laboratory
2. Department of Medical Imaging, Zinder National Hospital, Faculty of Health Sciences, André Salifou University of Zinder, Niger, Clinical Research and Health System Laboratory
3. Department of General Surgery, Zinder National Hospital, Faculty of Health Sciences, André Salifou University of Zinder, Niger, Clinical Research and Health System Laboratory
4. Department of Onco-Haematology, Zinder National Hospital, Faculty of Health Sciences, André Salifou University of Zinder, Niger, Clinical Research and Health System Laboratory
5. Department of Internal Medicine, Zinder National Hospital
6. Department of Hepato-Gastroenterology, Zinder National Hospital

### ABSTRACT

Portal cavernoma is a rare cause of upper gastrointestinal bleeding resulting from complications of portal hypertension. We report the case of a 30-year-old woman admitted with haematemesis and splenomegaly. Imaging revealed a portal cavernoma associated with oesophageal varices. The patient received a blood transfusion, fluid resuscitation, and on discharge was put on propranolol for secondary prevention. Endoscopic ligation could not be performed due to service limitations. Diagnosis of portal cavernoma relies on imaging. Its management is complex in low-resource settings such as Niger.

**Keywords:** haematemesis, cavernoma, portal hypertension, Zinder, Niger.

### Introduction

Portal cavernoma, or cavernous transformation of the portal vein, is a rare vascular anomaly characterized by the development of a network of tortuous venous collaterals that bypass an obstructed portal vein.<sup>[1,2]</sup> It causes non-cirrhotic portal hypertension.<sup>[3]</sup> The exact incidence is unknown, but increasing use of Doppler ultrasound and CT imaging has improved recognition.<sup>[4]</sup> The condition may remain asymptomatic for years and often presents through complications such as upper gastrointestinal bleeding caused by rupture of oesophageal varices.<sup>[4,5]</sup> Clinically, the picture is dominated by splenomegaly, hypersplenism, and signs of portal hypertension.<sup>[5]</sup> The diagnosis rests on imaging—mainly Doppler ultrasound and CT scan—which demonstrate the absence of flow in the main portal vein and the presence of multiple serpiginous collateral vessels.<sup>[1]</sup> We report a case of portal cavernoma discovered following haematemesis in a young woman with no previous medical history and managed in a resource-limited environment.

### Case Report

A 30-year-old woman with no history of liver disease, surgery, or thrombophilia presented to the Hepatology and Gastroenterology Department of Zinder National Hospital with haematemesis associated with dizziness and fever. She reported no use of non-steroidal anti-inflammatory drugs, no chronic abdominal pain, and no known peptic ulcer disease.

**Correspondence:**

Hamidine Illa

[ihamidine@gmail.com](mailto:ihamidine@gmail.com)**Submitted:** November 2025**Accepted:** March 2026**Published:** May 2026

**Citation:** Illa et al. A rare pathological feature of portal vein thrombosis complicated by portal hypertension and haematemesis. South Sudan Medical Journal, 2026;19(2):130-134  
© 2026 The Author(s) License: This is an open access article under [CC BY-NC](#) DOI: <https://dx.doi.org/10.4314/ssmj.v19i2.11>

At admission, she was pale, febrile (38.5°C), tachycardic (120 bpm), and hypotensive (80/60 mm Hg). Physical examination revealed stage III splenomegaly according to Hackett's classification and melaena on rectal examination. Laboratory investigations showed microcytic hypochromic anaemia (Hb = 5.9 g/dL) and thrombocytopenia (67,000/mm<sup>3</sup>). The thick blood film was positive for *Plasmodium falciparum*, while liver and renal function tests and coagulation studies were normal.

Initial management consisted of intravenous crystalloids with saline at 5ml/kg/hour, fresh blood transfusion at 756 ml to a transfusion target of 8g/dL, and 80 mg of omeprazole daily. Antimalarial therapy orally by artemether was initiated.

Upper gastrointestinal endoscopy was performed after stabilization, which revealed grade III oesophageal varices without red signs. The latter, e.g., cherry-red spots, would warn of imminent bleeding from weak areas. Abdominal ultrasound showed a serpiginous vascular formation in the hepatic hilum, suggestive of a portal cavernoma (Figure 1). Colour Doppler confirmed a chaotic flow within the portal venous network (Figures 2–3). A CT scan demonstrated a portal cavernoma without a focal hepatic lesion, associated with homogeneous splenomegaly (Figure 4). No biliary dilatation or mass was noted. A diagnosis of portal cavernoma complicated by variceal bleeding in a non-cirrhotic patient was made. Due to limited resources, endoscopic variceal ligation was unavailable. Anticoagulation was not indicated due to the chronic, organized nature of the thrombosis, and further bleeding might have been precipitated. Weekly follow-up over one month showed progress with cessation of bleeding and improved haemoglobin (10 g/dL). The patient received 40mg propranolol long-term, oral iron for three months, and six-monthly follow-up visits. She also received oral iron supplementation for three months.



Figure 1. Abdominal ultrasound scan showing a serpiginous appearance at the hepatic hilum, in favour of a portal cavernoma.




Figure 2. Doppler ultrasound illustrating a vascular malformation at the splenic vein level with heterogeneous flow, consistent with an ectopic vascular anastomosis.




Figure 3. Doppler ultrasound showing the portal vein with a serpiginous appearance indicative of a portal cavernoma.



Figure 4. Axial abdominal CT scan showing a portal cavernoma with associated homogeneous splenomegaly.

## Discussion

This is the first case of portal cavernoma reported at Zinder National Hospital. In Romania, Cazacu et al. found a prevalence of 14.8% among patients with portal thrombosis.<sup>[6]</sup> Despite its rarity, the condition is increasingly detected thanks to advanced imaging.

Our patient was 30 years old, within the range reported for non-cirrhotic portal vein thrombosis. In Senegal, Diallo et al. reported a mean age of 41 years (15–75 years), showing that the disease affects young adults. Although some series report a male predominance (M/F = 2.7: 1), gender differences may vary depending on underlying pro-thrombotic conditions such as sickle cell disease.<sup>[5]</sup>

The causes of portal vein thrombosis are often multifactorial. Devika et al. in India identified local inflammatory factors (pancreatitis, abdominal infections such as diverticulitis) and systemic thrombophilia, such as hyperhomocysteinemia.<sup>[7]</sup> Resseguier et al. in France found in 24 of 32 cases, 15 haematological disorders (10 myeloproliferative syndromes, 2 antiphospholipid syndromes, 1 thalassaemia major, 1 hyperhomocysteinemia and 1 factor II mutation).<sup>[8]</sup> In children, umbilical vein catheterization is the main cause.<sup>[9]</sup> In our case, no aetiology was apparent, i.e. idiopathic (Elkrief et al.<sup>[3]</sup>)

Upper gastrointestinal bleeding was the presenting symptom, as often described in the literature. Wei et al. highlighted variceal bleeding as the most common initial presentation.<sup>[10]</sup> Boccatonda et al. and Layton et al. also reported that variceal rupture is a typical manifestation of cavernous transformation.<sup>[1,2]</sup>

Splenomegaly is a classical sign of portal hypertension; its stage III presentation in our patient reflects advanced collateral circulation. Elkrief et al. emphasized that splenic hypertrophy is one of the most sensitive clinical indicators of chronic portal hypertension.<sup>[3]</sup> The presence of fever and malaria might have acted as a precipitating or aggravating factor for thrombosis.

Anaemia in our case was due to acute bleeding on chronic asymptomatic bleeding, while thrombocytopenia reflected hypersplenism secondary to portal hypertension. Cazacu et al. described similar haematological abnormalities, considering hypersplenism as a constant feature of cavernoma.<sup>[6]</sup> Normal liver function tests can help rule out cirrhosis.

Ultrasound scan remains the first-line diagnostic tool, especially in resource-limited settings. In our case, it showed serpiginous vascular structures in the

hepatic hilum. Colour Doppler revealed turbulent, multidirectional flow, characteristic of cavernous transformation. Marra et al. confirmed the high sensitivity of ultrasound for differentiating partial from complete portal thrombosis.<sup>[11]</sup> Kalra et al. emphasized that in chronic cases, the portal vein may disappear centrally and be replaced by a network of periportal collaterals.<sup>[12]</sup> A CT scan confirmed the diagnosis, showing multiple venous collaterals and splenomegaly without hepatic lesions. Mild biliary compression may occur, corresponding to portal biliopathy, a complication described in 5–30 % of cases.<sup>[13–14]</sup> Jha et al. reported that CT findings typically include non-enhancement of the portal vein, serpiginous collaterals, and absence of focal hepatic lesions in non-tumoral thrombosis.<sup>[15]</sup>

Endoscopy revealed grade III oesophageal varices. Boccataonda et al. also noted that large varices are common in cavernomas.<sup>[1]</sup> The absence of red signs suggests a lower immediate bleeding risk but requires secondary prophylaxis.

Management of portal cavernoma is challenging. Acute bleeding requires resuscitation, transfusion, and vasoactive or beta-blocker therapy after stabilization and correction of anaemia. We used propranolol for secondary prophylaxis. In chronic thrombosis, recanalization is often impossible, and the benefit of anticoagulation remains controversial. Boccataonda et al. argued against anticoagulation in fully organized thrombosis.<sup>[1]</sup> In well-equipped centres, endoscopic variceal ligation and trans-jugular intrahepatic portosystemic shunt (TIPS) are standard options for recurrent bleeding or refractory portal hypertension. However, these techniques were unavailable in our hospital, reflecting the therapeutic limitations in many African settings.

Prognosis depends on the control of variceal bleeding and the absence of complications such as portal biliopathy or intestinal ischaemia. Our patient showed favourable progress with improved haemoglobin and no recurrent bleeding after one month. Regular follow-up is essential to adjust beta-blocker dosage and consider further endoscopic evaluation.

## Conclusion

Portal cavernoma represents the chronic evolution of portal vein thrombosis, often of unknown aetiology. Its discovery is usually prompted by complications of portal hypertension, such as variceal bleeding. Biological abnormalities are nonspecific, but imaging and endoscopy are key to diagnosis. Management must be

multidisciplinary, combining resuscitation, beta-blockers, endoscopic treatment, and possibly TIPS or surgical shunts where available. In resource-limited environments, supportive care and close monitoring remain the cornerstone of management.

**Conflict of interest:** None

## References

1. Boccataonda A, Gentilini S, Zanata E, Simion C, Serra C, Simioni P, et al. Portal vein thrombosis: state-of-the-art review. *Journal of Clinical Medicine*. 2024; 13 (5):1517.
2. Layton BM, Lapsia SK. The portal vein: a comprehensive review. *Radiographics*. 2023; 43 (11):e230058.
3. Elkrief L, Hernandez-Gea V, Senzolo M, Albillos A, Baiges A, Berzigotti A, et al. Portal vein thrombosis: diagnosis, management, and endpoints for future clinical studies. *The Lancet Gastroenterology & Hepatology*. 2024; 9(9):859–883.
4. Ravololoniaina T, Rantoanina A, Andrianah PG, Randrianjanahary O, Ahmad A, Samison LH. Le cavernome portal: cause rare d'hémorragie digestive et d'ictère. *Journal Africain d'Hépatogastroentérologie*. 2014; 8(2):100–103.
5. Diallo S, Diagne CN, Bassène ML, Gueye MN, Fall MP, Thioubou MA, et al. Portal thrombosis: clinical, etiological and therapeutic aspects in the hepatogastroenterology department of Aristide Le Dantec Hospital, Dakar, Senegal. *Open Journal of Gastroenterology*. 2021; 11 (11):220–229.
6. Cazacu SM, Alexandru DO, Dumitrescu D, Vieru AM, Urhuf MC, Săndulescu LD. Prevalence and risk factors for portal cavernoma in adult patients with portal vein thrombosis. *Diagnostics*. 2024; 14(13):1445.
7. Bhat D. Clinical perspectives on cavernous transformation of the portal vein and hyperhomocysteinemia. *Journal of Clinical Perspectives*. 2024; 7(4):454–458.
8. Resseguier AS, André M, Orian Lazar EA, Bommelaer G, Tournilhac O, Delèvaux I, et al. Natural history of portal cavernoma unrelated to chronic liver disease: a monocentric retrospective study of 32 cases. *La Revue de Médecine Interne*. 2016; 37(6):394–398.

9. Basse I, Diagne Guèye NR, Obambi Diop DC, Diawara NN, Ba A, Seck N, et al. Portal cavernoma in children revealed by gastrointestinal bleeding: a case report. *Pan African Medical Journal*. 2016; 25:158. Available from: <http://www.panafrican-med-journal.com/content/article/25/158/full/>
10. Wei B, Huang Z, Tang C. Optimal treatment for patients with cavernous transformation of the portal vein. *Frontiers in Medicine*. 2022; 9:853138.
11. Marra P, Dulcetta L, Carbone FS, Muglia R, Muscogiuri G, Cheli M, et al. The role of imaging in portal vein thrombosis: from diagnosis to interventional radiological management. *Diagnostics*. 2022; 12(11):2628.
12. Kalra N, Shankar S, Khandelwal N. Imaging of portal cavernoma cholangiopathy. *Journal of Clinical and Experimental Hepatology*. 2014;4(Suppl 1):S44–S52.
13. Fatima-Zahra C, Berhili C, Borahma M. Portal cavernoma cholangiopathy: a case series with literature review. *IOSR Journal of Dental and Medical Sciences*. 2024; 23(8):56–60.
14. Kanan MF, Nagra R, Papantonatos C, et al. Successful treatment of portal cavernoma cholangiopathy with portal vein recanalization and transjugular intrahepatic portosystemic shunt. *Cureus*. 2025; 17(9):e92452.
15. Reena CJ, Satinderpal SK, Amit DK. Portal vein thrombosis: imaging the spectrum of disease with an emphasis on MRI features. *American Journal of Roentgenology*. 2018; 211:14–24.

## Advocacy Training for Community Health Workers

### A free digital advocacy training to elevate the voices of community health workers.

Community health workers (CHWs) are the most proximal and trusted care providers for millions worldwide. However, CHWs have historically been excluded from conversations and decision-making that influence their work and communities.

CHWs with advocacy training are **2-4x more likely** to engage in political, civic, and workplace advocacy. We worked with the **Community Health Impact Coalition** and an advisory group of CHWs from around the world to develop this training to equip CHWs with the skills to effectively share their personal stories and promote the health issues most important to their communities. All content is action-oriented and reflects the realities of CHWs' work.

The training is designed for individuals as well as group training and is flexible to fit existing training and work schedules. Videos, reading materials, and reflection exercises communicate the history and background of CHW programs, how to advocate for improved health care and working conditions, how to tell personal stories to advocate for change, and how to use technology tools to participate in global conversations.

The course is open-access and officially recognized as a **Digital Public Good**. This training includes four modules with nearly 40 videos, reflection exercises, Facilitator Guide, and Student Workbook. All materials are free for use under Creative Commons **CC-BY-NC-SA 4.0** license.

<https://digitalmedic.stanford.edu/advocacy-training-community-health-workers>

# Interpersonal and community-level factors influencing intimate partner violence in South Sudan: a literature review

Nyinypiu Tong Chol Adong

**Author Affiliation:**

School of Public Health, University of  
Juba, South Sudan

**Correspondence:**

[nyinypiu@gmail.com](mailto:nyinypiu@gmail.com)

**Submitted:** October 2025

**Accepted:** March 2026

**Published:** May 2026

**ABSTRACT**

The pervasive intimate partner violence continues to be perpetuated by entrenched gender norms, weak women's economic power, and post-war realities in South Sudan. In the face of global attention, South Sudan now considers gender-based violence not only as a health risk but also a human rights issue to safeguard. This study strived to identify and analyse interpersonal and community-level factors influencing intimate partner violence in South Sudan, from the existing literature. A systematic literature review was conducted using Heise's ecological framework, focusing on marital relations, household decision-making dynamics, and women's socio-economic status. The search engines utilised to obtain literature included PubMed, Medline, VU Library, Google Scholar, and the Ministry of Health's website. The research findings reveal that the genesis of intimate partner violence, in most cases, emanates from marital arrangements, including forced unions, polygamous setups, and bride price transactions. Importantly, women's exclusion from critical family decision-making processes exacerbates physical and sexual violence. Also, low educational attainment among women, as well as the levels of household poverty, unemployment, and peer influence, are strong predictors of violence in intimate relations. It is important to note that addressing intimate partner violence must be grounded in proper identification and understanding of the dynamics that perpetuate it. The interventions include a threshold for education above primary education, which is protective, women's empowerment, and poverty reduction at the community level.

**Keywords:** sexual violence, intimate partner violence, ecological model, marital dynamics, community factors, South Sudan

**Citation:** Adong. Interpersonal and community-level factors influencing intimate partner violence in South Sudan: a literature review. *South Sudan Medical Journal*, 2026;19(2):135-140 © 2026 The Author (s) **License:** This is an open access article under [CC BY-NC](#) DOI: <https://dx.doi.org/10.4314/ssmj.v19i2.12>

**Introduction**

In South Sudan, intimate partner violence (IPV) is rooted in social traditions, norms, beliefs, marital dynamics, and post-civil war realities.<sup>[1,2]</sup> The legal framework presents ambiguities that perpetuate such violence, although the national constitution proclaims gender equality.<sup>[3]</sup> The constitution is glaringly contravened by customary laws that "legitimise" male dominance and female subservience.<sup>[3]</sup>

At the core of IPV lies the institution of marriage, considering how the union is initiated and the prevailing social structure.<sup>[4-8]</sup> Marital violence often results from forced unions, polygamy, and bride price transactions.<sup>[6-8]</sup> These practices commodify women and affirm the men's entitlement over their spouses. With the judicial systems grounded in the long-standing customs, norms, and cultures, women often find it hard to file for divorce and exit from abusive relationships.<sup>[1]</sup> Importantly, women are excluded, or rarely consulted, in decision-making regarding family matters, which are the prerogative of men.<sup>[9]</sup> The already sore marital dynamics are compounded by economic violence, which subjects women to dependency and acquiescence to harmful practices.<sup>[10]</sup>

IPV is usually perpetuated by socialised negative notions that emanate from peer networks, mental health disorders, and militant masculinities emerging from the legacy of conflicts.<sup>[11-13]</sup> In the face of economic marginalisation of women, violence is strongly manifested in post-conflict situations as a means of reasserting male dominance and shifting gender roles.<sup>[13]</sup> At the community levels, IPV is shored up and enabled by shocking levels of household poverty, unemployment, and financial hardships.<sup>[10]</sup>

This study undertook a multi-layered inquiry into IPV, examining how relational and community-level factors converge to perpetuate abuse within marital relationships.<sup>[18]</sup> This includes reimagining marital relations, gender roles, and accountability based on the rule of law.

### Method

This research is a literature review that combines both peer-reviewed journals and grey literature. The search engines used to ascertain articles include PubMed, Medline, VU Library, Google Scholar, and the Ministry of Health websites to obtain relevant IPV literature in South Sudan and other relevant contexts. Articles published in the last ten years (2010 – 2019) in English were preferred as there is little change in attitudes, beliefs, and cultural norms related to IPV in South Sudan over this period.

The Heise's Integrated Ecological Framework was chosen for this review since it is more appropriate for the analysis of IPV in the context of South Sudan. However, the focus was on the analysis of interpersonal factors: marital conflict and types of marriages, women's decision-making status, educational level, community, and men's controlling behaviours. The remaining categories of the framework, such as the individual, society, and policies, were not within the scope of this study.<sup>[18]</sup>

## Results

### Interpersonal relations

The interpersonal factors include marital conflict and types of marriages, women's decision-making status, educational level, and men's controlling behaviours.

### Marital conflict and types of marriages

In South Sudan, marital conflict was found to be a major predictor of IPV.<sup>[4]</sup> Girls who were once raped were forced to marry the perpetrators as a way of resolving stigma issues associated with the survivor of violence. This was found to result in parental conflict between the couples with physical and emotional IPV.<sup>[5]</sup> Women married through abduction, payment of dowry, or married into polygamous relationships were more at risk of IPV.<sup>[6-8]</sup>

A study of gender norms in low- and middle-income countries (LMICs) found that IPV was associated twice as often with multiple sexual partners or extramarital sex compared to women with single sexual partners.<sup>[19]</sup> In Uganda, physical IPV was associated with extra-marital sex by women in intimate partner relationships, while sexual IPV was related to women whose husbands were jealous of seeing their wives talk with other men. The jealous men accused women of unfaithfulness and severely restricted them from contact with strangers and family members, resulting in emotional IPV.<sup>[20]</sup>

A study in LMICs found that physical IPV was 2.6 times more common where women had conflict in marital relationships compared to women who had agreeable relationships.<sup>[21]</sup> Women who offered sex to their husbands were less at risk of sexual violence compared to women who denied sex.<sup>[22]</sup> Additionally, in Ethiopia, physical IPV was more likely to affect women whose husbands had girlfriends and thought there was nothing they could do about it.<sup>[23]</sup> The odds of IPV were doubled for women who refused sex compared to those who complied.<sup>[7]</sup>

A study found that HIV infection was a source of marital conflict and increased IPV in northern Uganda, where there are South Sudanese refugees.<sup>[24]</sup> It was recognized that IPV was higher among couples infected with HIV compared to those who were not infected.<sup>[25]</sup>

### Women's decision-making status

In South Sudan, men were found to exclude women from important household decisions. Women were considered foreigners to their husbands' families.<sup>[9]</sup>

A multi-country study in conflict-affected settings, which included South Sudan, found that women were excluded from negotiating peace deals. Hence, peace accords failed to address violence against women and girls during the post-conflict period.<sup>[26]</sup> Women were not consulted or considered in decision-making that was related to politics or serious household issues.<sup>[27]</sup>

In Ethiopia, a study found a 50% reduction in IPV when women had equal decision-making power. There was also a 35% reduction in sexual IPV in women who equally shared in household decision-making processes with their husbands.<sup>[28]</sup>

A study in LMICs found that women in intimate partner relationships where men dominated decision-making were 8.2% more at risk of physical violence.<sup>[21]</sup> Additionally, joint spending of husbands' incomes was less associated with IPV.<sup>[21]</sup>

### **Educational level**

In South Sudan, 84% of women were found to be illiterate and vulnerable to IPV.<sup>[4,29]</sup> Poor education of women was associated with economic violence in intimate partner relationships, as they were forced to be financially dependent on men for survival.<sup>[30]</sup> Physical IPV was associated with gender inequality in less educated women compared to highly educated women.<sup>[31]</sup>

A multi-country cross-sectional study indicated that increased educational levels were associated with 9% decreased odds of physical IPV and 85% reduction in sexual IPV in the Democratic Republic of the Congo.<sup>[31]</sup> Research has shown that secondary or higher education for both partners was associated with reduced risks of IPV, while there was no protective effect of primary education against IPV.<sup>[25]</sup>

A combined study in Ethiopia and Tanzania found a two-thirds decrease in IPV when both partners had completed secondary education.<sup>[25]</sup> Another study found that the risks of IPV were twice as likely for uneducated or only primary-educated women.<sup>[32]</sup> IPV was less prevalent among women who had achieved higher educational levels.<sup>[21]</sup>

However, women who were more educated than their husbands were at higher risk of IPV compared to women who had less or equal educational levels as their husbands. Also, women in marriages where couples were equally highly educated experienced the lowest levels of IPV in Uganda.<sup>[21,33]</sup> The odds of IPV were three times higher for uneducated women compared to educated women in

intimate partner relationships.<sup>[21,34]</sup>

### **Men's controlling behaviours and mental disorders**

In South Sudan, women whose husbands exercised controlling attitudes were more likely to suffer both physical and emotional IPV.<sup>[5]</sup> Women whose husbands threatened their well-being were more at risk of IPV compared to women who were not afraid of their husbands.<sup>[35]</sup> There was evidence that men who exercised controlling behaviour over women were more likely to be violent compared to men who acknowledged gender equality.<sup>[4,25]</sup> IPV was also associated with women whose husbands were rude or hostile.<sup>[21,35]</sup> Men in rural areas who are more controlling and possessive of women compared to men in the towns were at higher risk of inflicting violence in intimate relationships.<sup>[4,9]</sup>

A multi-country study found that the odds of IPV were 3.5 times higher where husbands had a history of physical fights.<sup>[21]</sup> Emotionally stressed women were also more likely to experience IPV, and women with mental disorders had double the risk of IPV compared to women with healthy minds.<sup>[21]</sup>

### **Community level**

At the community level, factors that contribute to IPV include unemployment, poverty, and negative peer influence.

### **Poverty, socio-economic status, and unemployment**

South Sudan's families with low socio-economic status were more associated with IPV compared to affluent families.<sup>[10]</sup> Women of high socio-economic status were less at risk of IPV compared to women of poor backgrounds.<sup>[5]</sup> IPV was associated with a low Human Development Index (HDI), with South Sudan ranking 169th out of 188 countries worldwide.<sup>[4]</sup> Women were found to be the poorest and most vulnerable to economic IPV than men.<sup>[3,29]</sup> Families headed by women were the poorest compared to families headed by men.<sup>[4]</sup> A study in South Sudan found that unemployed poor women were at risk of economic violence compared to women who earned salaries.<sup>[4]</sup>

In Uganda, couples living in poverty were more at risk of IPV, which is associated with the perpetuation of social vices such as transactional sex and infectious diseases like HIV, sexually transmitted infections, and tuberculosis.<sup>[24]</sup> A multi-country study found that a lack of women's empowerment was associated with negative cultural norms

that denied women the right to own land and property.<sup>[25]</sup> Poverty made men lose control over their families, resort to drinking alcohol, and engage in physical IPV.<sup>[31]</sup>

A cross-national study on gender norms found that women dependent on informal jobs or farming were more likely to experience IPV compared to women who worked in formal jobs.<sup>[21]</sup> IPV was less likely when husbands were employed. Women of low socio-economic status were more vulnerable to IPV compared to women of high socio-economic status.<sup>[21]</sup> Sexual IPV was more likely when women were economically dependent on men compared to women who were financially independent. There was a 50% reduction in IPV where women owned businesses.<sup>[21]</sup>

### Negative peer influence

Multicountry studies have found that peer-group socialization is a major contributor to IPV.<sup>[25,36,37]</sup> Boys who socialized and oriented themselves negatively on how to handle women were more likely to inflict IPV compared to those who were not influenced through peer pressure.<sup>[21,38]</sup> Negative peer influence was associated with alcohol and drug abuse, resulting in IPV.<sup>[2]</sup> Intoxication with substances was found to trigger violence in the family, especially battery of women, compared to relationships in which couples were sober.<sup>[39,40]</sup>

## Discussion

### Interpersonal level

Marital conflicts should be addressed through couples' training and advocacy programmes aimed at raising awareness of human rights and women's rights. Skills that strengthen negotiation, conflict resolution, and leadership capabilities are important. Extreme patriarchal practices such as girl-abductions that are used to initiate marriages need to be addressed urgently, as are traditional practices such as polygamy that are rooted in gender inequality. Uganda's IPV prevention projects, such as Safe Homes and Respect for Everyone (SHARE) and Start Awareness, Support and Action (SASA), should also be adopted in South Sudan to confront household sexual violence and extra-marital sex or concurrent sex, which are associated with marital conflict and IPV. These strategies have proven to be effective in reducing IPV and concurrent sex associated with HIV infections.

Promoting girls' formal education is an important mechanism to ensure that more girls achieve secondary

education, thereby reducing the risk of IPV. Women should also be given capacity-building programmes through vocational training to acquire lifelong skills that enable them to achieve economic independence, thereby reducing IPV. Higher educational attainment is also associated with improved socio-economic status, thereby increasing employability.

Community mobilisation with group training and advocacy through video show projects such as "Through Our Eyes" (TOE) are directed at improving women's decision-making abilities. Equal gender decision-making processes are associated with reduced levels of IPV and improved gender relations. This also raises awareness of women's rights to participate in decisions about important family matters.

### Community level

The livelihood programmes, such as microfinance and cash transfers, coupled with skills training, are important interventions directed towards reducing women's poverty and vulnerability to IPV at community levels. Women who are economically empowered have increased ways of mitigating abusive relationships. Reduced household poverty levels also lead to satisfying reproductive and sexual lives, as women can access health services.

The livestock microfinance for peace in other sub-Saharan Africa is associated with decreased IPV and less poverty. These programmes should also be adopted in South Sudan where livestock (cattle and goats) is abundant, as a way of promoting peace, improving livelihood, and mitigating IPV.

Group training and advocacy for boys and men, engaging them in transforming negative attitudes towards women to reduce negative peer influence, is also a significant strategy for addressing IPV. The negative socialisation by men is seen to be associated with IPV at community levels. These training programmes should target gender norms and long-held traditional beliefs and practices that subjugate women.

## Conclusion

The evidence demonstrates IPV in South Sudan is deeply rooted in interpersonal, socio-economic, and community factors. Addressing IPV, therefore, requires multi-layered interventions. At the interpersonal level, couples' training, advocacy, and conflict resolution programmes can mitigate marital discord. Expanding girls' access to secondary education and vocational training enhances women's

autonomy and reduces vulnerability. At the community level, livelihood initiatives such as microfinance, cash transfers, and livestock-based peacebuilding can alleviate poverty and empower women economically. Finally, engaging men and boys in transformative gender training is essential to dismantling harmful norms and reducing peer-driven violence.

## References

1. Krug EG, Mercy JA, Dahlberg LL, Zwi AB. The world report on violence and health. *The lancet*. 2002 Oct 5;360(9339):1083-8.
2. Ndoromo O, Österman K, Björkqvist K. Low Intensity Intimate Partner Aggression as a Mediating Factor for the Intergenerational Transmission of Physical Punishment of Children in South Sudan. *Eurasian Journal of Medical Investigation*. 2018;2(4):183-9.
3. Mabor JM. Gender inequality, an examination of law and practice in South Sudan. *Afribery*. 2021.
4. Armah M, Amaah P MEJ. Health Cluster - Unity State, South Sudan. 2016.
5. Jansen HA, Garcia-Moreno C, Devries K, Heise L, Kiss L, Abramsky T, et al. What factors are associated with recent intimate partner violence? findings from the WHO multi country study on women's health and domestic violence. *BMC Public Health*. 2011;11(1).
6. UNICEF. Gender-Responsive Water, Sanitation and Hygiene: Key elements for effective WASH programming. *Transform Soc Norms Sex Reprod Heal*. 2017;(March).
7. Ndoromo O, Österman, Karin KB. Domestic Violence as a Risk Factor for Children Ending up Sleeping in the Streets of Post-War South Sudan. *Journal of Child & Adolescent Behavior*. 2017. 40. 41.
8. Ho U, Valladares E, Pen R, Lars A. Violence against pregnant women: prevalence and characteristics. A population-based study in Nicaragua. *BJOG an Int J Obstet Gynaecol*. 2005;112(September):1243-8.
9. Fulu E, Jewkes R, Roselli T, Garcia-Moreno C. Prevalence of and factors associated with male perpetration of intimate partner violence: Findings from the UN multi-country cross-sectional study on men and violence in Asia and the Pacific. *Lancet Glob Heal*. 2013;1(4): e187-207. 109X(13)70074-3 Available from: <http://dx.doi.org/10.1016/S2214>.
10. Abate BA, Wossen BA, Degfie TT. Determinants of intimate partner violence during pregnancy among married women in Abay Chomen district, Western Ethiopia: a community based cross sectional study. *BMC Womens Health*. 2016;1-8. Available from: <http://dx.doi.org/10.1186/s12905-016-0294-6>.
11. Salman SMA. The new state of South Sudan and the hydro-politics of the Nile Basi. *Water Int*. 2011;36(2):154-66.
12. Hill C. Associations between peer network gender norms and the perpetration of intimate partner violence among urban Tanzanian men: A multilevel analysis. *Matern Child Health J*. 2019;19(4):427-36.
13. Murphy M, Bingenheimer JB, Ovince J, Ellsberg M, Contreras-urbina. The effects of conflict and displacement on violence against adolescent girls in South Sudan: the case of adolescent girls in the Protection of Civilian sites in Juba. *Sex Reprod Heal Matters*. 2019;0397(May).
14. Partners H. User Guide: How to Support a Survivor of Gender-based Violence when there is no GBV actor in your area. 2012.
15. Gurman TA, Trappier RM, Acosta A, McCray PA, Cooper CM, Goodsmith L. "By seeing with our own eyes, it can remain in our mind": Qualitative evaluation findings suggest the ability of participatory video to reduce gender-based violence in conflict-affected. *June 2014Health Education Research* 29(4) doi: <https://doi.org/10.1093/her/cyu018>.
16. Orly Maya Stern. Gender Country Profile; International Law and Policy Institute. 2015.
17. Al-Tuwaijri S. Gender Based Violence in Fragile, Conflict, and Violence (FCV) Situations. *World Bank Gr*. 2018;1-8. 74. 75. 76. 77. 78.
18. Schwab-Reese LM, Renner LM. Screening, management, and treatment of intimate partner violence among women in low-resource settings. *Women's Health*. 2018;14. doi: <https://doi.org/10.1177/1745506518766709>.
19. García-Moreno C, Zimmerman C, Morris-

- Gehring A, Heise L, Amin A, Abrahams N, et al. Addressing violence against women: A call to action. *Lancet*. 2015;385(9978):1685–95.
20. García-Moreno C, Stöckl H. Protection of sexual and reproductive health rights: Addressing violence against women. *Int J Gynecol Obstet*. 2009;106(2):144–7.
21. Manjoo R, Mcraith C, Journal L. GBV and Justice in Conflict and post conflict areas. 44 *Cornell Int'l LJ*. 2012;11–32.
22. Semahegn A, Torpey K, Manu A, Assefa N, Tesfaye G, Ankomah A. Are interventions focused on gender-norms effective in preventing domestic violence against women in low and lower-middle income countries? A systematic review and meta-analysis. *Reprod Health* 16, 93 (2019). <https://doi.org/10.1186/s12978-019-0726-5>.
23. Lindsay Stark, Khudejha Asghar, Gary Yu, Caroline Bora, Asham Assazenew Baysa KLF. Prevalence and associated risk factors of violence against conflict – affected female adolescents: a multi – country, cross – sectional study. *J Glob Health*. 2017;7(1). 34–39.
24. West G. A mixed methods assessment of barriers to maternal, newborn and child health in. *Reprod Health*. 2017;1–13.
25. LOGICA. Gender and Conflict Note - South Sudan. 2012.
26. Gottschalk N. Uganda: early marriage as a form of sexual violence. *Forced Migr Rev*. 2016; 42. 43. 44. 45. 46. 47. 48. 49.
27. Annan J, Brier M. The risk of return: Intimate partner violence in Northern Uganda's armed conflict. *Soc Sci Med*. 2010; 70:152–9.
28. Abeya SG, Afework MF, Yalew AW. Intimate partner violence against women in western Ethiopia: prevalence, patterns, and associated factors. *BMC Public Health* 11, 913 (2011). <https://doi.org/10.1186/1471-2458-11-913>.
29. Eyobo MB, Bepo E, Lako RL, Wani G, Julla AI, Remijo CD, et al. Malaria indicator survey 2009, South Sudan: baseline results at household level. *Malar J*. 2014;13(1):45.
30. McCloskey LA, Hunter T. Determinants of Intimate Partner Violence in Sub-Saharan Africa: A Review of Prevention and Intervention Programs. *Partner Abuse*. 2016;
31. Tol WA, Stavrou V, Greene MC, Mergenthaler C, Van Ommeren M MC. Sexual and gender-based violence in areas of armed conflict: a systematic review of mental health and psychosocial support interventions. *Confl Health*. 2013;7(1).
32. Lien L, Hauff E, Martinez P, Eide AH, Swarts L, Ayazi T. Alcohol use in South Sudan in relation to social factors, mental distress and traumatic events. *BMC Public Health* 16, 937 (2016). <https://doi.org/10.1186/s12889-016-3605-9>
33. Ali AA, Yassin K, Omer R. Domestic violence against women in Eastern Sudan. *BMC Public Health* 14, 1136 (2014). <https://doi.org/10.1186/1471-2458-14-1136>
34. Murphy M, Bingenheimer JB, Ovince J, Ellsberg M, Contreras-urbina. The effects of conflict and displacement on violence against adolescent girls in South Sudan: the case of adolescent girls in the Protection of Civilian sites in Juba. *Sex Reprod Heal Matters*. 2019;0397(May).
35. Wachter K, Horn R, Puffer E. Drivers of intimate partner violence against women in three refugee Camps. *Violence Against Women*, 24(3), 286–306. <https://doi.org/10.1177/1077801216689163>
36. Nata Duvvury, Khalifa Elmusharaf, Stacey Scriver MC. Economic & Social Costs of VAWG in South Sudan. *Tech Rep*. 2019;(March).
37. Elli N Hayati, Ulf Högberg, Mohammad Hakimi, Mary C Ellsberg ME. Behind the silence of harmony: risk factors for physical and sexual violence among women in rural 35 Indonesia. *BMC Womens Health*. 2011;11(1):52. 50. 51. 52. 53. 54. 55. 56. 57. 58. 59. 60.
38. Jane Kani Edward. The SUDD Institute: Special Report. Vol. 14, Sapio. 2014. Available from: <http://ci.nii.ac.jp/naid/40004819541/en/>
39. Terry MS. Applying the Social Ecological Model to Violence against Women with Disabilities. *J Women's Heal Care*. 2014;3(6).
40. Yigzaw T, Berhane Y, Deyessa N, Kaba M. Perceptions and attitude towards violence against women by their spouses: A qualitative study in Northwest Ethiopia. *Glob Health Action*. 2010;24(1).

## Letter to the Editor

## RE: South Sudan Postgraduate Training Programme

12 February 2026

Dear Editor,

I agree with many of the concerns expressed by Dr E.A. Hakim, et al, regarding the need for a programme for postgraduate training in South Sudan (SSMJ Vol 19, No 1, February 2026). I feel it is important to point to other issues his letter exposes.

Firstly, most of the concerns expressed—high peripartum mortality, mortality in the first five years of life, all too common home deliveries— indicate the need to address medical issues outside of Juba. This is undermined by the misappropriation of funds supposed to flow through UNICEF to implementing partners at the county level. By all means, utilize professionals from outside countries to address the gaps in expertise in medicine in Juba, but from a practical standpoint, Link House will not make a significant impact on the issues expressed in his letter because most of those issues arise in the rural counties, not Juba.

Secondly, there is a great need for an element of humility at the Ministry of Health (MoH) in South Sudan. The vastly better outcomes in our East African neighbours reported by Dr. Hakim speak to the reality that the medical education systems in other countries in East Africa, including Uganda, Kenya, Tanzania, and Ethiopia, are more advanced. Their professionals, at all levels, are more highly qualified than ours. That is not surprising, given the reality that southern Sudan was engaged in a civil war with Khartoum while those other nations were finding their footing in government and education. It does our country no good to deny these facts. South Sudan needs help from our neighbours in order to improve the education of young medical professionals. During my years of association with the Rumbek Nurses Training School (RNTS), our goals of training nurses were undermined by restrictions imposed by the MoH in hiring instructors from outside South Sudan. The reality was there were not sufficiently trained mentors from South Sudan to accomplish the work, whereas numerous potential candidates were available and willing to come from Uganda and other countries.

Medical training is an apprenticeship based on science. To have well-trained doctors, nurses, clinical officers from South Sudan, we need experts in their fields to conduct the training. From my personal observations of being in Sudan/South Sudan for over 16 years, the best South Sudanese doctors at present were trained in Uganda or other East African countries. We must take advantage of the expertise in our neighbouring nations to provide mentorship for our young clinicians.

Thirdly, the desire to improve healthcare delivery nationwide points to the need to be able to hire the best qualified candidates for key positions in county hospitals and PHCC's. We are struggling to reopen a hospital that has been nonfunctional for over a decade. We are desperate for an excellent matron to oversee the process and an experienced laboratory technician to ensure high-quality and reliable test results. Numerous East Africans are available and desire to come, but the MoH will not allow those professionals to be employed because they are not South Sudanese. If the medical schools for Juba and Upper Nile University are ever to be comparable to those in Uganda and Kenya, it is essential to hire experienced clinicians and instructors who are best qualified to work with our medical colleagues and instruct the candidates, irrespective of their country of origin.

The healthcare of South Sudanese citizens is too important to be treated lightly. It is not sufficient to have the issues covered on paper. If we are to change the disastrous trends in South Sudanese healthcare where we are falling further behind our neighbours, we need to humble ourselves and address this glaring deficit.

J. Clarke McIntosh, MD

Medical Consultant, Akot Medical Centre

jclarkemcintosh@gmail.com

### Letter to the Editor

### Dr Eluzai A. Hakim Response

Dear Editor,

I refer to Dr McIntosh's letter, published in this issue in response to my article, "South Sudan Postgraduate Training Programme" in the SSMJ Vol 19, No 1, February 2026. Unlike Dr McIntosh, I contend that the Link House is a very important facility for supporting training in South Sudan especially in the initial stages of the evolution of the postgraduate programme to accommodate high-calibre South Sudanese specialists living abroad, and trainers from the developed world.

Dr McIntosh makes some very important points which require the attention of the South Sudan Ministry of Health (MOH), particularly regarding the development of a coherent National policy on a medical training programme. Recognising the lack of such a policy in 2008, I initiated the formation of a link between St. Mary's Hospital, Newport, Isle of Wight, and Juba Teaching Hospital (*The Juba Link*). The Link committee in the UK was made up of hospital consultants, senior nurses, a chief operating officer of the hospital trust, the manager of the St. Mary's Hospital Postgraduate Centre, a senior midwife, a hospital non-executive director and two interns. The two interns spent three months at Juba Hospital working alongside South Sudanese doctors for mutual experience. The corresponding Juba Committee was composed of a consultant surgeon, a paediatrician, a paediatrics registrar, and the hospital matron.

The aim of the Link was to harness the expertise of UK trainers to provide structured and regular on-site training for South Sudanese doctors in Juba to reduce costs. In addition, the link provided opportunities for South Sudanese doctors at various levels to visit UK hospitals for periods of six weeks to three months to experience different ways of clinical working in order to influence and, hopefully, improve healthcare delivery in South Sudan.

This scheme attracted much interest from Consultants in the Wessex Region of the United Kingdom, several of whom offered to travel to Juba at their own expense if accommodation was available. Encouraged by this enthusiasm by potential trainers, the Link committee launched a successful fundraising campaign on the Isle of



Wight which enabled the construction of a guest house called **the Link House** (see Figure) to accommodate visiting trainers. The MOH leased a plot of land on the premises of Juba Teaching Hospital where the *Juba Link* built a beautiful six ensuite-room guest house. Several trainers used the Link House on visits to Juba Hospital, saving the government appreciable amounts of money. A memorandum of understanding was signed between the MOH and the Juba Link to conduct trainer visits to Juba. Had it not been for the political upheaval in July 2016, the programme would have blossomed. The Link House was subsequently used for non-trainers as accommodation for hospital medical officers free of charge. This led to its rapid decline structurally without any maintenance! The Link House is available to support future trainers but needs renovation. I have taken steps to reactivate the Link with Juba Hospital through a London Hospital to support the postgraduate programme which I am determined to support.

To realise this objective the MOH must dedicate expertise and resources to developing a robust structured training programme as this cannot be solely achieved through voluntary effort. The Juba Link has ceased to exist through retirements of the trustees and lack of financial support.

Finally, the Consultants currently working at Juba Hospital cannot be bystanders but must bring pressure to bear on the MOH to develop this programme.

Dr Eluzai Hakim, FRCP Edin, FRCP

[eluzaihakim@doctors.org.uk](mailto:eluzaihakim@doctors.org.uk)

## News from the Gordon Memorial College Trust Fund

At the last annual Committee meeting of the Gordon Memorial College Trust Fund (GMCTF) on 19th March 2026, applications for postgraduate grants were considered.

The results are presented below:

Total number of individual applications submitted: **142**

Total number of individual applications approved: **31** (10 female)

- Applicants with Sudanese nationality: **4** (2 female)
- Applicants with South Sudanese nationality: **27** (8 female)

Total amount awarded **\$101,100 USD**

- **\$12,200** to Sudanese individuals (\$5,000 to females)
- **\$88,900** to South Sudanese individuals (\$25,350 to females)

### Breakdown of 27 South Sudanese grant recipients

#### Current location:

- 20 – Ethiopia
- 4 – Uganda
- 2 – South Sudan
- 1 – Kenya

#### Institution of study:

- 5 - Addis Ababa University
- 4 - Woliata Sodo University
- 3 - Jimma University
- 3 - Yekatit 12 Hospital Medical College

- 2 - Hawassa University
- 3 - St. Paul's Hospital Millennium Medical College
- 3 - Makerere University
- 1 - Adama Hospital Medical College, Ethiopia
- 1 – COSECSA (Juba Military Referral Hospital)
- 1 - Kampala International University
- 1 - University of Nairobi

### Post-grad qualifications that will be gained include:

- Specialty Certificate - Internal Medicine
- Specialty Certificate - Obstetrics and Gynaecology
- Membership of COSECSA
- Master's degree in general surgery
- MD – Paediatrics
- MSc. Neurosurgery

Many applicants were either ineligible for the award of a grant or did not align well with GMCTF's published criteria on the charity website [www.gmctf.org](http://www.gmctf.org).

Potential applicants should closely follow the GMCTF guidance (including the information we require from referees) if they want to improve their chances of success. Recycling previous references is not acceptable. Those applying for extension of their grants must produce clearly written evidence from their Programme Directors and/or Deans of the relevant faculties that they have successfully completed the courses in the current year of study and have been deemed suitable to progress to the next level of the course.

**Eluzai Abe Hakim**, FRCP Edin, FRCP

Member, GMCTF Committee.

Every effort has been made to ensure that the information and the drug names and doses quoted in this Journal are correct. However readers are advised to check information and doses before making prescriptions. Unless otherwise stated the doses quoted are for adults.

# Is it Flu or Ebola?



## Flu (influenza)



The **flu** is a common contagious respiratory illness caused by flu viruses. The flu is different from a cold.


Flu can cause mild to severe illness, and complications can lead to death.

## Ebola



**Ebola** is a rare and deadly disease caused by infection with an Ebola virus. Sporadic outbreaks have occurred in some African countries since 1976.

## How Flu Germs Are Spread



The flu is spread mainly by droplets made when people who have flu cough, sneeze, or talk. Viruses can also spread on surfaces, but this is less common.

People with flu can spread the virus before and during their illness.

## How Ebola Germs are Spread



People get Ebola by direct contact with

- The body fluids of a person who is sick with or has died from Ebola.
- Objects contaminated with body fluids of a person sick with Ebola or who has died of Ebola.
- Infected fruit bats and primates (apes and monkeys)
- And, possibly from contact with semen from a man who has recovered from Ebola (for example, by having oral, vaginal, or anal sex)

## Who Gets The Flu?



Anyone can get the flu.

Some people—like very young children, older adults, and people with some health conditions—are at high risk of serious complications.

## Who Gets Ebola?



People most at risk of getting Ebola are

- People with a travel history to countries with widespread transmission or exposure to a person with Ebola.
- Healthcare providers taking care of patients with Ebola.
- Friends and family who have had unprotected direct contact with blood or body fluids of a person sick with Ebola.

## Signs and Symptoms of Flu



The signs and symptoms of flu usually develop within 2 days after exposure. Symptoms come on quickly and all at once.

## Signs and Symptoms of Ebola



The signs and symptoms of Ebola can appear 2 to 21 days after exposure. The average time is 8 to 10 days. Symptoms of Ebola develop over several days and become progressively more severe.

- ***People with Ebola cannot spread the virus until symptoms appear.***



- **Fever or feeling feverish**
- **Headache**
- **Muscle or body aches**
- **Feeling very tired (fatigue)**
- **Cough**
- **Sore throat**
- **Runny or stuffy nose**



- **Fever**
- **Severe headache**
- **Muscle pain**
- **Feeling very tired (fatigue)**
- **Vomiting and diarrhea develop after 3–6 days**
- **Weakness (can be severe)**
- **Stomach pain**
- **Unexplained bleeding or bruising**

For more information about the flu and Ebola, visit [www.cdc.gov/flu](http://www.cdc.gov/flu) and [www.cdc.gov/ebola](http://www.cdc.gov/ebola).

May 4, 2015

CS252294-1