

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/338384361>

# Expanding Point-of-Care Ultrasound Training in a Low- and Middle-Income Country: Experiences From a Collaborative Short-Training Workshop in Kenya

Article in *Family Medicine* · January 2020

DOI: 10.22454/FamMed.2020.986896

CITATIONS

5

READS

61

9 authors, including:



**Catherine Gathu**

Aga Khan University Hospital, Nairobi

2 PUBLICATIONS 20 CITATIONS

[SEE PROFILE](#)



**Akiruga James**

Moi University

9 PUBLICATIONS 7 CITATIONS

[SEE PROFILE](#)



**Kevin Bergman**

Contra Costa Regional Medical Center

5 PUBLICATIONS 28 CITATIONS

[SEE PROFILE](#)


Some of the authors of this publication are also working on these related projects:



Point of Care Ultrasound Project [View project](#)



Milo Bigpic Family Project [View project](#)



# Expanding Point-of-Care Ultrasound Training in a Low- and Middle-Income Country: Experiences From a Collaborative Short-Training Workshop in Kenya

Leanne Jones, MD, MPH; Catherine Gathu, MBChB; Daria Szkwarko, DO, MPH; Sam Mucheru, MBChB; Naushad Amin, MD; James A. Amisi, MBChB, MMED; Kevin Bergman, MD; Mena Ramos, MD; Neil Jayasekera, MD

**BACKGROUND AND OBJECTIVES:** In Kenya, little data exists on point-of-care ultrasound (POCUS) training and use for family medicine physicians. In 2017, a 3-day POCUS workshop assembled most of the family medicine physicians in Kenya. Through surveys, we assessed how this workshop could affect the level of POCUS use, skill, and confidence in family medicine practitioners in the long term.

**METHODS:** Structured surveys, distributed before, after, and 10 months post-workshop assessed demographics, POCUS use, barriers, comfort, and skills based on attendee self-assessment. We compared data from the preworkshop surveys to postsurveys and post-postsurveys to assess immediate and long-term differences. Wilcoxon signed-rank test was used to evaluate continuous data, and significance was based on a *P* value of <.05.

**RESULTS:** The proportion of participants who self-reported using POCUS increased significantly between presurvey and post-postsurvey (29.7% to 63.2%, *P*=.0161). Mean confidence scores increased significantly from presurvey to postsurvey and post-postsurvey. For all body systems, self-reported mean skill scores increased significantly from presurvey to postsurvey and post-postsurvey. Lack of access to machines and mentorship are substantial barriers to increasing POCUS use.

**CONCLUSIONS:** This study highlights the utility of one-time POCUS training in increasing long-term POCUS uptake by participants. While encouraging, our findings also show barriers to increasing POCUS use. These barriers must be addressed, potentially through intradepartmental and interorganizational exchanges of resources to ensure that future POCUS workshops are successful in supporting POCUS use in Kenya.

(*Fam Med.* 2020;52(1):38-42.)  
doi: 10.22454/FamMed.2020.382138

Access to diagnostic imaging is a challenge in health facilities of low- and middle-income countries (LMICs). A 2011 World Health Organization (WHO) survey reported that the density of computerized tomography (CT) scans was 2.06 per million people in LMICs compared to 44.31 in high-income countries.<sup>1</sup> As an alternative, point-of-care ultrasound (POCUS) use has recently increased in LMICs.<sup>2</sup>

POCUS is attractive for multiple reasons. Units are compact and operate with rechargeable batteries, minimal infrastructure and training are required, and diagnosis can be done solo at the bedside. The WHO stated that capital costs for the most advanced ultrasound machine were

---

From the University of Massachusetts Medical School, Department of Family Medicine and Community Health, Worcester, MA (Drs Jones and Szkwarko); Department of Family Medicine, Aga Khan University Hospital Nairobi, Kenya (Drs Gathu and Mucheru); Department of Family Medicine, Warren Alpert School of Medicine, Brown University, Providence, RI (Dr Szkwarko); Contra Costa Family Medicine Residency Program, an affiliated program of the University of California San Francisco Department of Family and Community Medicine (Drs Amin, Bergman, Ramos, and Jayasekera); and Moi University School of Medicine, Department of Family Medicine, Eldoret, Kenya (Dr Amisi).

10% of those for a CT machine.<sup>3</sup> Although quality evidence is lacking, a systematic review demonstrated that portable ultrasound may have a clinical impact in up to 70% of cases.<sup>2</sup>

In Kenya (an LMIC), family physicians are well situated to utilize POCUS in multiple settings yet lack formal training. In 2017, the Kenya Association of Family Physicians (KAFP) organized a 3-day training in collaboration with Moi, Aga Khan, Kenyatta, and Kabarak Universities, as well as the Contra Costa, University of Massachusetts, and Brown University family medicine (FM) residency programs. Contra Costa FM has been conducting POCUS workshops for over 5 years<sup>4</sup> with a course based on American College of Emergency Physician Guidelines. Our study aimed to assess the immediate and long-term (10-month) impact of an adapted version of this course on POCUS use, confidence, and skills for FM trainees and faculty in Kenya.

## Methods

In May 2017, we conducted a 3-day POCUS course for all FM physicians and trainees in Kenya, with residents (registrars) and faculty

(consultants) from four out of five FM residency programs attending. The course comprised lectures, large-group demonstrations, and hands-on scanning. See Figure 1 for course agenda.

We distributed pre- and post-structured surveys before and immediately after the workshop, and 10 months postworkshop (Appendix 1). Question types included multiple choice, open-ended, and Likert scale.

Surveys were anonymous but linked by a unique identifier. A single investigator (L.J.) entered all data into a Microsoft Excel (version 1901) spreadsheet and used Wilcoxon signed-rank test to evaluate continuous data. Significance was based on a *P* value of <.05. Statistical analysis was performed using Stata/IC 15.

The University of Massachusetts Institutional Review Board reviewed the project and determined that it was not human subjects research given that this was a program evaluation and survey data was anonymous.

## Results

Of 41 participants, 39 completed the surveys and were included in

pre/post analysis. Of these participants, 28 (72%) were male and 11 (28%) were female; 26 (67%) were registrars; 34 (87%) reported never having POCUS training previously. Twenty (51%) participants completed the 10-month post-survey, 19 (49%) of whom were matched to the original pre- and posttraining analysis and were included in post-post analysis (Table 1).

The three most cited barriers on presurvey were cost of machine, insufficient training, and lack of formal POCUS curriculum (Figure 2). The proportion of respondents who cited lack of training as a barrier to POCUS use decreased significantly from presurvey to postsurvey (47.37% to 5.13%, *P*=.00000) and from presurvey to post-postsurvey (47.37% to 15.79%; *P*=.0198). The proportion of participants who self-reported using POCUS increased significantly between presurvey and post-postsurvey (29.7% to 63.2%; *P*=.0161).

Participants rated their confidence in using POCUS in practice, teaching POCUS, and sharing POCUS findings with patients. For all three questions, mean confidence score increased significantly from presurvey to postsurvey, and from presurvey to

Figure 1: POCUS Workshop Agenda

<b>Friday, May 12<sup>th</sup></b>	<b>Saturday, May 13<sup>th</sup></b>	<b>Sunday, May 14<sup>th</sup></b>
1230-1300: Registration and Check-In 1300-1315: Introduction 1315-1330: Ultrasound Basics 1330-1400: FAST Exam 1400-1500: FAST and E-FAST lab 1500-1530: Chai 1530-1600: Pulse Health Care 1600-1625: Liver/Spleen/Gallbladder 1625-1650: Renal/Aorta 1650- 1800: Liver/Spleen/Gallbladder and Renal/Aorta lab 1800-1900: Dinner	0730-0830: Breakfast 0800-0830: Check-In 0830-0850: Cardiac Echo 0850-1000: Cardiac Echo lab 1000-1030: Chai 1030-1050: Pulmonary 1050-1110: DVT 1110-1210: Pulmonary and DVT lab 1210-1300: Pass the Pointer 1300-1400: Lunch 1400-1430: Soft tissue/Procedural 1430-1530: Soft tissue/Procedural lab 1530-1600: Chai 1600-1620: OB 1620-1640: Gynecology 1640-1730: OB/GYN lab 1730-2200: Dinner	0730-0830: Breakfast 0800-0830: Check-In 0830-0900: FASH exam 0900-1000: FASH lab 1000-1030: Chai 1030-1050: RUSH exam 1050-1150: RUSH lab 1150- 1250: OSCE 1250-1350: Advanced applications/Closing remarks 1330-1430: Survey completion and lunch

Abbreviations: POCUS, point-of-care ultrasound; FAST, focused assessment with sonography in trauma; E-FAST, extended FAST; DVT, deep venous thrombosis; OB, obstetric; GYN, gynecologic; FASH, focused assessment with sonography for HIV-associated tuberculosis; RUSH, rapid ultrasound for shock and hypotension; OSCE, objective structured clinical examination.

Table 1: Key Demographics of Survey Respondents

Characteristics	Pre/Post	Post-Post	P
Total Respondents	39	19	-
Age (mean, years)	35.35	35.45	.9633
Gender			
Male	28 (72%)	12 (63%)	.5046
Female	11 (28%)	7 (37%)	.5046
Clinical Practice			
Registrar	26 (67%)	15 (79%)	.3449
Consultant	11 (28%)	4 (21%)	.5670
Other	2 (5%)	0 (0%)	-
Training Program			
Aga Khan	9 (23%)	3 (16%)	.5369
Kabarak	6 (15%)	2 (11%)	.6775
Kenyatta	8 (21%)	6 (32%)	.3613
Moi	11 (28%)	8 (42%)	.2857
Past POCUS Training			
Yes	5 (13%)	-	-
No	34 (87%)	-	-
Level of Access to POCUS			
Have machine, can access	13 (33%)	7 (37%)	.7653
Have machine, not easily available	4 (10%)	4 (21%)	.2556
Machine available at other training sites	0 (0%)	1 (5%)	-
No access	17 (44%)	6 (32%)	.3856
Other	3 (8%)	1 (5%)	.6773
Frequency of POCUS Use in Practice			
Never	19 (49%)	6 (32%)	.2243
Daily	1 (3%)	0 (0%)	-
Few times a week	3 (8%)	9 (47%)	.0007
Few times a month	8 (21%)	3 (16%)	.6539
Few times a year	6 (15%)	1 (5%)	.2707

Abbreviations: Pre/Post, participants who completed presurvey and postsurvey; Post-Post, participants who completed presurvey, postsurvey, and post-postsurvey; P, P value; Registrar, Kenyan resident-equivalent; Consultant, Kenyan attending equivalent, POCUS, point-of-care ultrasound.

post-postsurvey, but decreased significantly from postsurvey to post-postsurvey (Table 2).

For all body systems, self-reported mean skill scores increased significantly from presurvey to postsurvey, and from presurvey to post-postsurvey. Self-reported mean skill scores decreased significantly across all body systems from postsurvey to post-postsurvey, except for obstetric and liver/spleen/gallbladder (Table 3).

## Discussion

Considering the potential of POCUS use by FM physicians in Kenya, it is necessary to determine whether training modalities such as short workshops are effective. Our findings show that self-reported use, skills, and confidence with POCUS increased significantly after a short workshop and that this increase, while declining slightly after 10 months, remained significantly elevated from preworkshop levels.

This highlights the utility of one-time, basic POCUS training in increasing long-term POCUS uptake by participants and confirms findings of previous studies.<sup>5,6</sup> The workshop played a key role in addressing the lack of training that was noted by most trainees as a barrier in the presurvey and in other studies.<sup>7</sup>

Our study was one of the first to assess perceived barriers to POCUS use in an LMIC 10 months after a POCUS workshop, allowing insight

Figure 2: Perceived Barriers to POCUS Use by Survey Type




Table 2: Comparing Mean Confidence Scores

	Comparing Pre and Post					Comparing Pre and Post-Post					Comparing Post and Post-Post				
	N	M Pre	M Post	M Diff	P	N	M Pre	M PP	M Diff	P	N	M Post	M PP	M Diff	P
Using POCUS in practice	23	1.78	3.65	1.87	.00	12	1.67	3.00	1.33	.00	19	3.53	3.00	-0.53	.01
Teaching POCUS	23	1.35	3.30	1.96	.00	12	1.17	2.83	1.67	.00	19	3.11	2.63	-0.47	.04
Sharing results with patients	23	1.78	3.57	1.78	.00	12	1.75	3.08	1.33	.00	19	3.42	3.00	-0.42	.03

Note: Mean confidence scores reflect the average response on a 5-point Likert scale with 1=extremely confident and 5=not at all confident. The scale was reversed in analysis, and here reflects 1=not at all confident and 5=extremely confident.

Abbreviations: POCUS, point-of-care ultrasound; Pre, presurvey; Post, postsurvey; Post-Post, 10-month postsurvey; N, number analyzed; M Pre, mean confidence score from presurvey; M Post, mean confidence score from postsurvey; M PP, mean confidence score from 10-month postsurvey; M Diff, mean difference between scores; P, P value.

into the reasons for the decline in skills and confidence over that period. Lack of ready access to a machine was the most commonly cited barrier to POCUS use. Ultrasound devices, while relatively inexpensive, remain cost prohibitive in LMICs.<sup>8,9</sup> Nongovernmental organizations are distributing POCUS machines to rural health care facilities across the country, but full coverage to all facilities is yet to be realized.

The other commonly cited barrier related to lack of continuous, onsite POCUS training. Although faculty attended the workshop, they were also in the learning stages. Collaborations among departments and other institutions both nationally and internationally could provide opportunities for such mentorship.

This study has several strengths. Our workshop marked the first time that most Kenyan FM physicians and trainees were gathered

together, and our survey results provide new insight into the state of the field of family medicine in Kenya. Additionally, our 10-month postsurvey assessed the impacts of the workshop over time, suggesting that although uptake remains increased compared to preworkshop levels, future workshops must facilitate machine donations and foster accessible teaching relationships to promote sustainability.

Table 3: Comparing Self-Reported Mean Skill Scores

System	Comparing Pre to Post					Comparing Pre to Post-Post					Comparing Post to Post-Post				
	N	M Pre	M Post	M Diff	P	N	M Pre	M PP	M Diff	P	N	M Post	M PP	M Diff	P
Cardiac	36	1.36	3.28	1.92	.00	19	1.37	2.68	1.32	.00	19	3.16	2.68	-0.47	.00
Ob	36	1.69	3.22	1.53	.00	19	1.63	2.89	1.26	.00	18	2.89	2.89	.00	1.00
Pulm	36	1.33	3.42	2.08	.00	19	1.32	2.37	1.05	.00	18	3.22	2.39	-0.83	.00
FAST	36	1.56	3.58	2.03	.00	18	1.44	2.89	1.44	.00	19	3.37	2.89	-0.47	.01
FASH	36	1.14	3.42	2.28	.00	18	1.17	2.39	1.22	.00	18	3.28	2.39	-0.89	.00
RUSH	34	1.24	3.47	2.24	.00	18	1.22	2.50	1.28	.00	18	3.39	2.50	-0.89	.00
DVT	37	1.54	3.65	2.11	.00	19	1.63	3.21	1.58	.00	19	3.58	3.21	-0.37	.03
L/S/G	37	1.51	3.51	2.00	.00	19	1.47	3.05	1.58	.00	19	3.37	3.05	-0.32	.22
Gyn	37	1.46	3.27	1.81	.00	19	1.47	2.79	1.32	.00	19	3.26	2.79	-0.47	.04
Renal	37	1.43	3.46	2.03	.00	19	1.42	2.84	1.42	.00	19	3.42	2.84	-0.58	.03
Soft tissue	36	1.39	3.42	2.03	.00	19	1.58	2.84	1.26	.00	18	3.28	2.61	-0.67	.00

Note: Self-reported mean skill scores reflect the average response on a 5-point Likert scale with 1=highly skilled and 5=not at all skilled. The scale was reversed in analysis, and here reflects 1=not at all skilled and 5=highly skilled.

Abbreviations: POCUS, point-of-care ultrasound; Pre, presurvey; Post, postsurvey; Post-Post, 10-month postsurvey; N, number analyzed; M Pre, mean skill score from presurvey; M Post, mean skill score from postsurvey; M PP, mean skill score from 10-month postsurvey; M Diff, mean difference between scores; P, P value; Ob, obstetric; FAST, focused assessment with sonography in trauma; FASH, focused assessment with sonography in HIV-related tuberculosis; RUSH, rapid ultrasound for shock and hypotension; DVT, deep venous thrombosis; L/S/G, liver/spleen/gallbladder; Gyn, gynecologic

**ACKNOWLEDGMENTS:** Study findings were presented at the STFM Annual Spring Conference, May 7, 2018, Washington, DC, as an oral presentation entitled “On The Same Wavelength: Assessing a Cross-Cultural POCUS Workshop in Kenya.”

**FINANCIAL SUPPORT:** The 2017 POCUS Workshop was partially funded by Pulse Healthcare Limited.

**CORRESPONDING AUTHOR:** Address correspondence to Dr Leanne Jones, University of Massachusetts Medical School, Department of Family Medicine and Community Health, 55 Lake Avenue North, Worcester, MA 01655. 774-442-6499. Fax: 774-441-6212. leannemasae@gmail.com.

## References

1. Velazquez-Berumen A. WHO Perspectives on Health Technologies: Medical Imaging. Presentation 2011 Annual Meeting of European Society of Radiology. Vienna, Austria; March 4, 2011.
2. Becker DM, Tafoya CA, Becker SL, Kruger GH, Tafoya MJ, Becker TK. The use of portable ultrasound devices in low- and middle-income countries: a systematic review of the literature. *Trop Med Int Health*. 2016;21(3):294-311.
3. Future use of new imaging technologies in developing countries. Report of a WHO Scientific Group. *World Health Organ Tech Rep Ser*. 1985;723:1-67.
4. Jayasekera NB. Point of Care Ultrasound (POCUS) at Contra Costa. <https://cchealth.org/residency/ghf/pocus.php#resources>. Accessed July 12, 2018.
5. Henwood PC, Rempell JS, Liteplo AS, et al. Point-of-care ultrasound use over six-month training period in Rwandan district hospitals. *Afr J Emerg Med*. 2013;3(4)(suppl):S5-S6.
6. Wanjiku GW, Bell G, Wachira B. Assessing a novel point-of-care ultrasound training program for rural healthcare providers in Kenya. *BMC Health Serv Res*. 2018;18(1):607.
7. Shah S, Bellows BA, Adedipe AA, Totten JE, Backlund BH, Sajed D. Perceived barriers in the use of ultrasound in developing countries. *Crit Ultrasound J*. 2015;7(1):28.
8. Sue K. The occasional ED ultrasound: focused assessment with sonography for trauma (FAST). *Can J Rural Med*. 2015;20(1):33-39.
9. Blois B. Office-based ultrasound screening for abdominal aortic aneurysm. *Can Fam Physician*. 2012;58(3):e172-e178.
10. Moore DE Jr, Green JS, Gallis HA. Achieving desired results and improved outcomes: integrating planning and assessment throughout learning activities. *J Contin Educ Health Prof*. 2009;29(1):1-15.