

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/274195478>

Characteristics of serum agglutinins in marine crab *Scylla serrata* Forskal and their interaction with various bacteria species

Article in *International Journal of Current Microbiology and Applied Sciences* - January 2013

CITATIONS

2

READS

71

4 authors, including:



Ogutu Philip Ariya

Masinde Muliro University of Science and Technology

14 PUBLICATIONS 17 CITATIONS

[SEE PROFILE](#)



Chrispinus Mulambalah

Moi University

32 PUBLICATIONS 148 CITATIONS

[SEE PROFILE](#)



Donald Siamba

Masinde Muliro University of Science and Technology

30 PUBLICATIONS 146 CITATIONS

[SEE PROFILE](#)


Some of the authors of this publication are also working on these related projects:



natural products reserach [View project](#)



Effects of Nematode Infection on Immune Responses to Killed Vaccine and Live Mycoplasma Capricolum Capripneumoniae antigenus in Goats [View project](#)



Original Research Article

Characteristics of serum agglutinins in marine crab *Scylla serrata* (Forsk.) and their interaction with various bacteria species

A.Ogutu Philip¹, P.Mullainadhan², S.Mulambalah Chrispinus³, N.Siamba Donald^{1*}

¹Masinde Muliro University of Science and Technology, Department of Biological Sciences, P.O. Box 190, Post Code 50100, Kakamega, Kenya

²University of Madras, Guindy Campus, Department of Zoology, Laboratory of Pathobiology, Chennai – 600 025, India

³Moi University, School of Medicine, Department of Medical Microbiology and Parasitology, P.O. Box 4606 Post Code 30100 Eldoret, Kenya

*Corresponding author

ABSTRACT

Keywords

Multiple-agglutinins; hemagglutination; bacterial agglutination; serum agglutinin; crustacean; divalent cation dependent; carbohydrates; glycoproteins; host defense.

Agglutinins or lectins are conventionally defined as proteins/glycoproteins of non-immune origin with a remarkable ability to specifically and reversibly interact with carbohydrate ligands. Although lectins are produced by a myriad of organisms, some of which are phylogenetically remote, and therefore exhibit distinct physicochemical and biochemical characteristics, lectins from different sources may essentially exhibit common biological activities. This study was therefore undertaken to preliminarily characterize the lectins in the serum of *S. serrata* as a prerequisite for isolation, purification and biological activity evaluation. Multiple agglutinins were detected in the serum of the marine crab *Scylla serrata*. The serum agglutinins agglutinated a wide range of mammalian erythrocytes (RBC). The hemagglutinating activity was high within a pH range of 7.0 to 8.5, and was dependent on divalent cations Ca^{2+} , Sr^{2+} & Mg^{2+} , with preference for Ca^{2+} . An agglutinating activity independent of Ca^{2+} was also demonstrated, indicating the presence of heterogenous agglutinins. The serum HA activity was inhibited by a wide range of related and unrelated carbohydrates, suggesting the presence of multiple agglutinins. The serum agglutinins reacted/interacted with a variety of bacteria including the *Vibrio* species known to be the most frequent opportunistic pathogens of aquatic crustaceans.

Introduction

Agglutinins or lectins are conventionally defined as proteins/glycoproteins of non-immune origin with a remarkable ability to specifically and reversibly interact with carbohydrate structures present on cell surfaces, extracellular matrices or secreted

glycoproteins (Goldstein *et al.*, 1980; Barondes, 1988; Sharon and Lis, 1995; Weis., 1997; Goldstein 2002). They have also been demonstrated to additionally interact or recognize non-carbohydrate ligands bearing an appropriate structural

determinants (Gabijs, 1997; Kawagishi *et al.*, 1994; Zhang *et al.*, 1995; Gokudan *et al.*, 1999; Murali *et al.*, 1994; Maheswari *et al.*, 2002). The lectins may possess mono-, di-, or multi-valent carbohydrate binding sites, whereas agglutinins possess di- or multi-valent sites. The latter structural characteristic is an essential prerequisite for agglutinin molecules to cause agglutination of cells displaying on their surfaces more than one saccharide of sufficient complementarity (Barondes, 1981).

Lectins are produced by a myriad of organisms, some of which are phylogenetically remote, and exhibit distinct physicochemical and biochemical characteristics although lectins from different sources may essentially exhibit common biological activities. Various studies with lectins at molecular levels are in progress all with ultimate goal of not only the identification and structural characterization of all of their natural ligands, and the elucidation of the nature of the relevant binding interactions at atomic resolution, but also a full understanding of the biological roles of these molecules. The latter must eventually be confirmed by natural or genetically manipulated mutations in the expression of the lectins and/or their ligands in intact animals.

The biological functions of agglutinins are attributed primarily to their ability to recognize and bind to specific carbohydrate structures (Sharon and Lis. 1989). Invertebrate agglutinins have been implicated in a variety of endogenous biological functions. These include feeding, larval settlement, fertilization, and diverse endogenous functions such as cell aggregation, embryonic development, metamorphosis, regeneration, wound

repair and transport of carbohydrates (Yeaton, 1981; Rögner and Uhlenbruck. 1984; Olafsen, 1986; Vasta, 1991). Based on these inherent properties, lectins are not only already being exploited in identifying cells with specified glycan signatures, but also in quantifying these carbohydrate motifs. New and efficient techniques for cell separation/enrichment based on the conjugation of lectins to magnetic particles and affinity columns have been developed (Galvez *et al.* 2002; Shinya *et al.* 2004; Bakalova & Ohba 2003; Putnam *et al.* 2003). Lectins are also major components of the innate immune system of several decapod crustaceans, including lobster, crabs, shrimps and freshwater crayfish (Bang, 1962; Cornick, and Stewart. 1968; Ratcliffe, *et al.* 1985; Richards, and. Renwranz. 1991; Mercy and Ravindranath 1994; Wilson, *et al.* 1999) where they defend these invertebrates from invading bacterial, fungal and viral pathogens (Stewart and Zwicker, 1972; Noga *et al.*, 1996).

Naturally occurring agglutinins in the serum of the marine crab *S. serrata* have been studied previously by different investigators, which mainly involved detection and isolation of single humoral agglutinins. In these independent studies, the isolated agglutinins were reported to express specificity for Neu5Gc (Mercy, and. Ravindranath. 1993) and sialoglycoconjugates such as fetuin (Chattopadhyay and Chatterjee, 1993) or BSM (Kongtawelert, 1998). Although the actual carbohydrate structures recognized by the two agglutinins reactive with sialoglycoconjugates remain unresolved, these findings, from different laboratories suggest the possible occurrence of at least three agglutinins in the serum of *S. serrata*. The role of Neu5Gc-specific agglutinin in hemocoelic clearance of

foreign materials has been investigated by Mercy and Ravindranath (1994), but the involvement of two other agglutinins in immuno-defense functions in *S. serrata* has not been examined. This study was therefore undertaken to preliminarily characterize the lectins in the serum of *S. serrata* as a prerequisite for isolation, purification and pharmacological studies.

Materials and Methods

Acquisition and Preparation of Reagents and buffers

Carbohydrates (mono-, di-, tri- & polysaccharides) were products of s.d. Fine, BDH, CDH & Qualigens (Bombay, India) or Sigma (St. Louis, Missouri, USA). Glycoproteins (bovine submaxillary mucin (BSM- Type II), asialo-mucin, fetuin, asialo-fetuin, porcine thyroglobulin, porcine stomach mucin (PSM- Type III) and ovalbumin), Ethylene glycol-bis (β -aminoethyl ether) N, N, N¹, N¹ - tetraacetic acid (EGTA) and bacterial species (*Vibrio anguillarum*, *V.alginolyticus*, *V.parahaemolyticus*, *V.fluvialis*, *V.vulnificus*, *V.mimicus*, *Escherichia coli*, *Pseudomonas sp* *Bacillus subtilis* and *Aerobacter aerogenes*- Type I) were purchased from Sigma All other chemicals and reagents used were of the highest analytical grade and were supplied by local agencies. The chemicals were used to prepare tris-buffered saline specified in table 1.

All these buffer solutions contained 0.02% NaN₃, and the osmolarity were determined using a cryoscopic osmometer (osmomat 030, gonotec, Germany) and adjusted if necessary by either dissolving a few crystals of NaCl or diluting with distilled water. All buffers were stored at 10⁰C, and used within a week. Other solutions used

were saline (0.9% NaCl & 0.02% NaN₃) and Alsever's solution (Garvey et al. 1979). Alsever's solution was autoclaved at 15 psi for 15 min, cooled to room temperature (RT = 26 \pm 2⁰C), then 50 mg streptomycin were dissolved in the solution and stored at 10⁰C until use.

Crabs and preparation of serum

Agile, uninjured, non-autotomized and intermoult male marine crabs (*Scylla serrata*) weighing 200-300 g were purchased from a local fish market in Chennai- India. The crabs were maintained in plastic tanks containing filtered sea water (salinity; 30‰) and fed *ad libitum* with the surface clam *Donax sp.* The sea water was aerated continuously and changed daily. The tip of the dactylus region of the walking leg was wiped with ethanol-soaked cotton, dried with tissue paper and cut with a sharp pair of scissors. The exuding hemolymph was collected in tubes held on ice and allowed to clot at room temperature (RT). The clot was vigorously disrupted using a glass rod, and centrifuged (450 x g, 10 min, RT). The resulting clear supernatant (serum) was used immediately.

Preparation of erythrocyte (RBC) suspension

Human A, B, O, sheep, goat, ox, buffalo, horse, rabbit, rat and mouse blood samples were obtained by venous or cardiac puncture where applicable. All the blood samples were collected in sterile Alsever's solution (Garvey et al. 1979) containing 10 μ g/ml of streptomycin, stored at 10⁰C and used within five days. The Alsever's solution was decanted and replaced daily. RBC in the blood samples were washed thrice with 0.9% saline and once with TBS-IV (containing 10 mM CaCl₂) by

centrifugation (400 x g, 5 min, RT). Unless otherwise specified, the washed RBC pellet was finally re-suspended in 5 ml TBS-IV (containing 10 mM CaCl₂) as 1.5% (v/v).

De-O-acetylation of bovine submaxillary mucin (BSM)

De-O-acetylation of BSM was performed following the procedure of Sarris & Palade (1979). Briefly, 5 mg of BSM were dissolved in 375 µl of TBS-IV (containing 10 mM CaCl₂) mixed with 125 µl of 0.4 N NaOH (10 mg BSM.ml⁻¹) and incubated on ice for 45 minutes. The solution was brought to room temperature and then neutralized by adjusting the pH to 7.2 using 0.1 N HCl. Finally, the solution was dialysed (MW exclusion limit < 3000-5000 Da) extensively against the same buffer and the dialysate represented de-O-acetylated BSM.

Hemagglutination (HA) assays

The hemagglutination assays were performed in V-bottomed microtiter plates (Greiner, Nürtingen, Germany) by serial dilution of 25 µl serum samples with an equal volume of appropriate TBS. 25 µl of RBC suspension was added to each well and incubated for 1 hr at RT. The HA titers were recorded as the reciprocal of the highest dilution of the samples causing complete agglutination of RBC (Garvey *et al.* 1979). Controls for all assays consisted of the substitution of the sample by TBS. Each experiment was performed in duplicate for at least three times using samples from different preparations, and the hemagglutination activities were analyzed based on the median titer values. The assay was repeated with TBS- IV containing 10 mM CaCl₂ at different pH (ranging between 6.5 and 8.5).

Divalent cation dependency and EDTA/EGTA sensitivity

Serum samples (each 250 µl) were dialysed (MW exclusion limit < 10,000 Da) extensively against cation-free TBS (TBS -I) at 15°C to examine cation dependency, or against TBS-VI containing 50 mM EDTA to test its sensitivity to EDTA. The samples dialysed against TBS-VI were subsequently re-equilibrated by dialysis against TBS-I. All the resulting dialysates were centrifuged (400 x g, 5 min, RT) and the hemagglutinating activity in the supernatant was determined using human B RBC in the presence of buffer that did (TBS-IV) or did not (TBS-I) contain 10 mM CaCl₂ or SrCl₂ or MgCl₂. EDTA sensitivity was also analysed by two-fold serial dilution of 25 µl serum sample in TBS-VI containing 25 or 50 mM EDTA and mixing with an equal volume of human B RBC in the corresponding buffer. For control, the EDTA-containing buffers were substituted by TBS-IV (containing 10 mM CaCl₂). The effect of EGTA was similarly analysed by serial dilution of 25 µl serum samples in TBS containing 25 or 50 mM EGTA (TBS-VII). For controls, the EGTA containing buffers were substituted by TBS-IV containing 10 mM CaCl₂.

Effect of different concentrations of divalent cations on HA activity of the serum

HA activity was assayed as described above. The effect of different concentrations of divalent cations was assayed by serially diluting the serum sample and adding human B RBC suspensions containing different cations at various concentrations (TBS-IV and TBS-V). The assay was repeated using rabbit, mouse and rat RBC.

Hemagglutination - Inhibition (HAI) assays

Several simple sugars and their derivatives (dissolved in TBS-VIII), polysaccharides and glycoproteins (dissolved in TBS-IV containing 10 mM, CaCl₂, pH 7.2.) were tested for their ability to inhibit serum hemagglutinating activity. The pH of the test substances was adjusted, wherever necessary to 7.2 using small crystals of NaOH. Serum was diluted with TBS-V containing 10 mM CaCl₂, pH 7.2 or 50 mM CaCl₂ pH 8.0 to hemagglutination titer of 4 against human B RBC. 25 µl of the diluted serum was added to each well and an equal volume (25 µl) of the test substance was serially diluted with the serum in the titer plate. After 1 hr incubation at RT, 25 µl of human B RBC suspension (prepared in TBS-IV containing 10 or 50 mM CaCl₂) was added to each well and incubated further for 1 hr at RT. The inhibitory potency of the test substance is expressed as the minimal concentration of the substance causing inhibition of the hemagglutinating activity.

Bacterial agglutination assays

Cultures of eight bacterial species; namely *Vibrio anguillarum*, *V.alginolyticus*, *V.parahaemolyticus*, *V.fluvialis*, *V.vulnificus*, *V.mimicus*, *Escherichia coli* and *Pseudomonas sp.* isolated from sea water and harvested at their stationary phase were used. An aliquot (3 ml) of each bacterial culture in nutrient broth was washed thrice in 0.9% saline by centrifugation (4,000 x g, 20 min, 4°C). The pelleted bacteria were resuspended in 3 ml of saline, heat-killed (1hr, 60°C) and then washed once with TBS-IV (containing 10mM CaCl₂, pH 7.2) by centrifugation as described above. Other two bacterial species *Aerobacter*

aerogenes-Type I and *Bacillus subtilis* were suspended in appropriate volume of the same buffer. The optical density of all the bacterial suspensions were adjusted to 0.8 (=1 x 10⁷ cells. ml⁻¹) at 580 nm.

Bacterial agglutination assay was performed in slides. 50 µl of serum sample mixed with an equal volume of each bacterial suspension in the cavity of glass slide. These slides were kept in moist chamber and incubated overnight at RT. The agglutination of bacteria was examined under phase optics (400 x) in a Carl Zeiss (Axiolab) phase contrast microscope, and the reaction was scored qualitatively based on the size and the number of bacterial aggregates observed.

Result and Discussion

Hemagglutination profile of the serum

The serum agglutinated all the mammalian RBC types tested, except ox RBC. It gave moderate to high HA titers for buffalo, rat, human B, mouse and rabbit RBC (titers =16 to 128). Sheep, goat, horse, human O and A RBC were weakly agglutinated (titers = 2 to 8).

Effect of pH, divalent cation and EDTA/EGTA on serum HA activity

The HA titer of serum was 32 or 64 within a pH range of 7.0 to 8.5 and the activity decreased to a titer of 8 at pH 6.5 (Figure 1). The HA sample dialysed against TBS-I and then tested in the absence of divalent cations did not show any HA activity (Table 2). However, the HA activity was restored in the presence of CaCl₂, SrCl₂ or MgCl₂, but it was completely restored only with CaCl₂. Similar results were obtained with serum samples dialyzed against TBS-VI containing 50 mM EDTA. In serial

Table.1 Types of tris-buffered saline (TBS) used to characterize the serum agglutins from the marine crab *Scylla serrata*

Type	Specifications
TBS-1	50 mM Tris-HCl, 125 mM NaCl (pH 7.2, 300 mOsm)
TBS –II:	100 mM Tris- HCl, 230 mM NaCl (pH 7.2, 560 mOsm);
TBS – III	100 mM Tris HCl, 200 mM NaCl (pH 7.2, 500 mOsm);
TBS – IV	Several buffers were prepared consisting of 50 mM Tris-HCl, and different concentrations of NaCl, CaCl ₂ , or MgCl ₂ or SrCl ₂ at different pH values and all at 300 mOsm such as: 115 mM NaCl, 10 mM CaCl ₂ (pH 6.5, 7.0, 7.5, 8.0, 8.5); 90 mM NaCl, 25 mM CaCl ₂ or MgCl ₂ or SrCl ₂ (pH 7.2); 50 mM NaCl, 50 mM CaCl ₂ (pH 7.2 or 8.0); 50 mM NaCl, 50 mM MgCl ₂ or SrCl ₂ (pH 7.2); 115 mM NaCl, 10 mM CaCl ₂ or SrCl ₂ or MgCl ₂ (pH 7.2);
TBS – V	50 mM Tris-HCl, 10 mM NaCl, 100 mM CaCl ₂ or SrCl ₂ or MgCl ₂ (pH 7.2, 370 mOsm);
TBS VI	50 mM Tris-HCl, 115 mM NaCl, 25 mM EDTA or 50 mM Tris-HCl, 100 mM NaCl, 50 mM EDTA (pH 7.2, 300 mOsm)
TBS- VII	50 mM Tris-HCl, 115 mM NaCl, 25 mM EGTA or 50 mM Tris HCl, 100 mM NaCl, 50 mM EGTA (pH 7.2, 300 mOsm)
TBS- VIII	50 mM Tris-HCl, 30 mM NaCl, 10 mM CaCl ₂ (pH 7.2, 135 mOsm).

Table.2 Divalent cation dependency and EDTA sensitivity of hemagglutinating activity of *Scylla serrata* serum.

Treatment of serum	Divalent cations tested [@]	HA titer*
Untreated serum (before dialysis)	CaCl ₂	32
Serum dialysed against cation-free TBS (TBS-I)	None	0
	CaCl ₂	32
	SrCl ₂	16
	MgCl ₂	16
Serum dialysed against TBS + 50 mM EDTA (TBS-VI) and re-equilibrated by dialysis against TBS-I	None	0
	CaCl ₂	32
	SrCl ₂	16
	MgCl ₂	16
Control (untreated sample) held for 48h at 10 ⁰ C	CaCl ₂	32

*Data represent median values from three determinations with human B RBC using serum samples from different preparations.

Table.3 Hamagglutination-inhibition of *Scylla serrata* serum (hemagglutination titer = 4) by glycoproteins against human B RBC

Glycoproteins tested	Maximum Concentration tested (mg.ml ⁻¹)	Predominant terminal sugar residues	Minimum inhibitory concentration (mg.ml ⁻¹)*
Porcine thyroglobulin	10	Neu5Gc	1.250
Fetuin	10	Neu5Ac α(2-3) Gal	0.156
Asialo-fetuin	5	Gal	0.312
Bovine submaxillary mucin (BSM)	10	O-IN-AcNeu5Ac α(2-6) GalNAc	0.156
Asialo-BSM	5	GalNAc	0.312
De-0-acetylated BSM	5	N-AcNeu5Ac α(2-6) GalNAc	0.312
Porcine stomach mucin (PSM)	10	Neu5Gc	0.625
Ovalbumin	10	GlcNAc	2.500

*The assay was performed three times for each glycoprotein

Figure. 1 Effect of pH on hemagglutinating activity of the serum of *Scylla serrata*.





Figure.2 Effect of different concentrations of divalent cations on hemagglutinating activity of the serum of *Scylla serrata*.



dilution experiment, both 25 and 50 mM EDTA completely abolished the serum HA activity whereas a residual activity (titer = 4) was observed with EGTA tested at identical concentrations. When tested in the presence of other cations, it was observed that HA activity of serum against human B RBC increased steadily from a titer of 32 or 16 in the presence of 10 mM CaCl₂ or SrCl₂ to 128 or 64 at 50 mM CaCl₂ (Figure 2). However, further increase in the concentration of these two divalent cations up to 100 mM did not enhance the HA activity of the serum. On the other hand, the HA activity increased only marginally from 10 mM to 25 mM MgCl₂ but in the presence of 50 mM CaCl₂, HA of the serum was enhanced from a HA titer of 128 to 256 (rabbit RBC), 64 to 128 (mouse RBC) and 32 to 128 (rat RBC).

Hemagglutination - inhibition (HAI) assays

Among the 41 carbohydrates tested, 25 carbohydrates were found to inhibit serum

hemagglutination activity at concentrations ranging from 12.5 to 100 mM (Table 3). Only the acetylated hexosamines (GlcNAc, GalNAc and ManNAc) were inhibitory at 50 or 100 mM. Among the six different polysaccharides tested [i.e Laminarin (β 1-3 homopolymer of glucose), Mannan (β 1-6 homopolymer of mannose) Dextran T₇₀ (α 1-6,3,2 homo- polymer of glucose), Dextran T₅₀₀ (α 1-6,3,2 homo- polymer of glucose), Inulin (α 2-6 homopolymer of fructose) and Colominic acid (α 2-8 homopolymer of Neu5Ac)], only mannan and laminarin inhibited the HA activity at 0.25 and 0.50 mg.ml⁻¹, respectively. In contrast, all the eight glycoproteins tested were inhibitory at various concentrations in which fetuin and BSM were the most potent inhibitors (0.156 mg.ml⁻¹). The patterns of inhibition observed with various simple sugars and their derivatives, polysaccharides, and glycoproteins were essentially same in the assays performed with TBS-IV (containing 10 or 50 mM CaCl₂, pH 7.2 or 8.0)

Bacterial agglutination profile of the serum

All the ten bacterial species tested, comprising both Gram -ve and Gram +ve bacteria, were agglutinated by serum of *S. serrata*. The highest degree of agglutination was observed with *Vibrio fluvialis* whereas moderate agglutination was seen with *V.alginolyticus*, *V.vulnificus*, *V.anguillarum*, and *Bacillus subtilis*. Other bacterial species including *V.parahemolyticus*, *V.mimicus*, *E.coli*, *Pseudomonas sp.* and *A. aerogenes*, were weakly agglutinated by the crab serum.

The serum was found to agglutinate various mammalian erythrocytes with high reactivity against rabbit, mouse, human B and rat RBC. These results concur with those from the previous study (Mercy and Ravindranath. 1992). The serum agglutinating activity was high within a pH range of 7.0 to 8.5, and lacks specificity for human blood groups, a feature that appears to be common among crustaceans (Sarris and Palade, 1979). Since cross adsorption tests demonstrated the efficiency of human B RBC in adsorbing from serum all agglutinating activity for this and other RBC types (Mercy and Ravindranath. 1992), this RBC was used as a suitable indicator cell type for all subsequent experiments performed.

Several/Related studies on humoral agglutinins in crustaceans revealed that their activity is dependent on divalent cations, usually calcium ions, and sensitive to divalent cation chelators such as EDTA and EGTA (Smith and Chisholm. 1992; Marques and Barraco. 2000). However, few exceptions to these features have been reported (Imai *et al.*, 1994; Maheswari *et al.* 1997; Kondo *et al.*, 1992). Dialysis

experiments performed in the current study against cation-free TBS demonstrated that the HA activity in the crab serum is dependent on divalent cations such as Ca^{2+} , Sr^{2+} or Mg^{2+} since the activity was not detected in the absence of these divalent cations. Moreover, the activity was recovered upon addition of these cations, but complete recovery was achieved only with CaCl_2 , suggesting that the serum HA activity exhibits requirement for divalent cations with a preference for Ca^{2+} . Similar results were obtained from dialysis experiments performed with EDTA, thereby indicating that the serum HA activity is reversibly sensitive to EDTA, a divalent cation chelator. Furthermore, serial dilution experiments performed with both EDTA and EGTA indicated that the serum HA activity is sensitive to EDTA, whereas it is only partially sensitive to EGTA (Ca^{2+} chelator). This finding suggests the possible occurrence of an agglutinating activity independent of Ca^{2+} , thus indicating the presence of at least two agglutinins, which are heterogeneous with regard to Ca^{2+} requirement. The results obtained with various concentrations of different divalent cations, demonstrated that HA activity increased in the presence of especially CaCl_2 or SrCl_2 in a dose-dependent manner, with a maximal activity detectable at 50 mM a further indication of the presence of multiple agglutinins.

In hemagglutination-inhibition assays performed in this study, several diverse and unrelated monosaccharides and their derivatives as well as di- and oligosaccharides inhibited the serum agglutinating activity. In addition, all the three acetylated hexosamines (GlcNAc, GalNAc and ManNAc), but not their hexose and hexosamine the counterparts

tested, consistently inhibited the HA activity of crab serum, thereby suggesting affinity of the serum agglutinin(s) for acetylated aminosugars. It is also notable that the two sialic acids Neu5Ac and Neu5Gc tested were not inhibitory even upto concentrations of 40 and 20 mM, respectively. By contrast, Mercy & Ravindranath (Mercy, and. Ravindranath. 1992) demonstrated the inhibitory activity of Neu5Gc at 0.6 mM as well as inability of all the three acetylated hexosamines to inhibit the serum HA activity of *S. serrata*. The exact reason for these discrepancies between the two studies is not clear at present. Further studies with polysaccharides indicated that only laminarin and mannan inhibited the serum agglutinating activity, albeit weakly, while free glucose and mannose were non-inhibitory, indicating that the agglutinin molecules in crab serum tend to exhibit affinity for extended structures particularly for polysaccharides with β -linked hexoses.

Further inhibition tests were performed with glycoproteins comprising eight sialo- and asialo-glycoconjugates, whose oligosaccharide side chains, as reported earlier (Schauer, 1982; Tsuji and. Osawa. 1986; Corradi *et al.*, 1995), contain predominantly terminal Neu5Gc (thyroglobulin and PSM) and Neu5Ac (fetuin, BSM and de-O-acetylated BSM), galactose (asialo-fetuin) and GlcNAc (ovalbumin). In this inhibition assay, the serum agglutinin(s) appear to show higher affinity to sialoglycoproteins terminated with Neu5Ac (fetuin and BSM) or Neu5Gc (PSM and thyroglobulin) as well as desialylated glycoproteins, namely, asialofetuin and asialo-BSM terminated with Gal and GalNAc respectively. These observations suggest that the serum agglutinin(s) recognize two types of sialic acids Neu5Ac or Neu5Gc which is

apparent in inhibition studies with glycoproteins, but not with free sialic acids. Thus all the results obtained from the inhibition studies clearly indicate that the serum agglutinins of *S. serrata* interact with a wide range of carbohydrates including acetylated hexosamines, acetylated or non-acetylated sialic acids and several other carbohydrates, and therefore their specificity for a specific carbohydrate structure could not be ascertained. These findings however strongly suggest the natural occurrence of multiple agglutinins in the serum of this crab.

It is also interesting to note that serum agglutinated a variety of bacteria including Gram +ve and Gram -ve types and the species of *Vibrio* tested are known to be the most frequent opportunistic pathogens of aquatic crustaceans (Egidius, 1987). The ability of the serum of *S. serrata* to agglutinate the bacteria, particularly the potential pathogens implicates a possible involvement of the humoral agglutinins in host defense response. It is therefore concluded that heamolymph extract of the marine crab *Scylla serrata* exhibits properties indicative of multiple lectins and further studies are recommended to isolate and characterize the individual lectins.

Acknowledgement

We thank Dr. M.H. Ravindranath (Director, John Wayne Institute for Cancer Treatment and Research, Santa Monica, CA, USA) and Dr. G. Dayanithi (University of Montpellier II, Montpellier, France) for generous gifts of many fine chemicals and laboratory materials used in this study. This study was supported by the University Grants Commission (UGC), New Delhi, India, through the award of

Junior Research Fellowship (No. F. 19-1/2000 (SA-II) dated 12.04.2001).

References

- Bakalova, R., and Ohba, H. 2003. Purification of normal lymphocytes from leukemic T-cells by lectinaffinity adsorbents - correlation with lectin-cell binding. *Cancer. Lett.* 192(1):59-65.
- Bang, F.B., 1962. Serological aspects of immunity in invertebrates. *Nature*, 196:88-89.
- Barondes, S.H., 1981. Lectins: Their multiple endogenous cellular functions. *Ann. Rev. Biochem.* 50:207-231.
- Barondes, S.H., 1988. Bifunctional properties of lectins; lectins redefined, *Trends Biochem. Sci.* 13:480-482.
- Cassels, F.J., J.J. Marchalonis and Vasta, G.R. 1986. Heterogenous humoral and hemocyte-associated lectins with N-acetylaminosugar specificities from the blue crab, *Callinectes sapidus* Rathbun. *Comp. Biochem. Physiol.* 85B:23-30
- Chattopadhyay, T., and Chatterjee, B.P. 1993. A low-molecular weight lectin from the edible crab *Scylla serrata* hemolymph: purification and partial characterization. *Biochem. Arch.* 9:65-72.
- Cornick, J.W., and Stewart, S. 1968. Interaction of the pathogen *Gaffkya homari* with natural defense mechanisms of *Homarus americanus*. *J. Fish. Res. Bd. Can.* 25:695-709.
- Corradi Da Silva, M.L., H.J. Stubbs, T. Tamura and Rice, K.G. 1995. ¹H NMR characterization of a hen ovalbumin tyrosinamide N-linked oligosaccharide library. *Arch. Biochem. Biophys.* 318:465-475.
- Egidius, E., 1987. Vibriosis: pathogenicity and pathology. A review. *Aquaculture*, 67:15-28.
- Fragkiadakis, G. and Stratakis, E. 1995. Characterization of hemolymph lectins in the prawn *Parapenaeus longirostris*. *J. Invertebr. Pathol.* 65:111-117.
- Gabius, H.J., 1997. Animal lectins. *Eur. J. Biochem.* 243:543-576.
- Galvez F., S.D Reid., G Hawkings and Goss, G.G. 2002. Isolation and characterization of mitochondriarich cell types from the gill of freshwater rainbow trout. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 282(3): R658-668.
- Garvey, J.S., N.E. Cremer and Sussdorf, D.H. 1979. *Methods in Immunology* (3rd Edn), W.A. Benjamin Inc., Reading, Massachusetts, 545pp.
- Gokudan, S., T. Muta, R. Tsuda, K. Koori, T. Kawahara, N. Seki, Y. Mizunoe, S.N. Wai, S. Iwanaga and Kawabata, S. 1999. Horseshoe crab acetyl group-recognizing lectins involved in innate immunity are structurally related to fibrinogen. *Proc. Natl. Acad. Sci.* 96:10086-10091.
- Goldstein, I.J., 2002. Lectin structure-activity: the story is never over. *J Agric Food Chem* 50(22):6583-6585.
- Goldstein, I. J., R.C. Hughes, M. Monsigny, T. Osawa and Sharon, N. 1980. What should be called a lectin? *Nature.* 285:66.
- Imai, T., R. Goto, J. Kittaka and Kamiya, N. 1994. Lectins in the rock lobster *Jasus novaehollandiae* hemolymph. *Crustaceana.* 67:121-130.
- Kamiya, H., and Ogata, K. 1982. Hemagglutinins in the acorn barnacle *Balanus (Megabalanus) roseus*: purification and partial characterization. *Bull. Japan. Soc. Sci. Fish.* 48:1421-1425.
- Kawagishi, H., M. Yamawaki, S. Isobe, T. Usui, A. Kimura and Chiba, S. 1994. Two lectins from the marine sponge

- Halichondria okadai*. An N-acetyl-sugar-specific lectin (HOL-II) . J. Biol. Chem. 296:1375-1379.
- Kondo, M., H. Matsuyama and Yano, T. 1992. The opsonic effect of lectin on phagocytosis by hemocytes of kuruma prawn, *Penaeus Japonicus*. Gyobyu Kenkyu. 27:217-222.
- Kongtawelert, P., 1998. Isolation and characterization of lectin from Thai marine crab (*Scylla serrata*) with binding specificity to sialoglycoconjugates and its application. Mol. Mar. Biol. Biotechnol. 7:280-286.
- Maheswari, R., P. Mullainadhan and Arumugam, M. 1997. Characterization of a natural hemagglutinin with affinity for acetylated aminosugars in the serum of the marine prawn, *Penaeus, indicus* (H. Milne Edwards) Fish Shellfish Immunol. 7:17-28.
- Maheswari, R., P. Mullainadhan and Arumugam, M. 2002. Isolation and characterization of an acetyl group recognizing agglutinin from the serum of the Indian white shrimp *Fenneropenaeus indicus*. Arch. Biochem. Biophys. 402,65-76.
- Marques, M.R.F., and Barraco, M.A. 2000. Lectins as non-self recognition factors in crustaceans. Aquacult. 191: 23-44.
- Mercy, Sr. P.D. , and . Ravindranath, M.H. 1994. Hemolysis and clearance of erythrocytes in *Scylla serrata* are related to the agglutination by native sialic acid- specific lectin .Comp. Biochem. Physiol. 109A:1075-1083.
- Mercy, Sr. P.D., and Ravindranath, M.H. 1993. Purification and characterization of N-glycolyl-neuraminic acid-specific lectin from *Scylla serrata* .Eur. J. Biochem. 215:697-704.
- Mercy, Sr. P.D., and Ravindranath, M.H. 1992. An agglutinin with unique specificity for N-glycolyl sialic acid residues of thyroglobulin in the hemolymph of a marine crab *Scylla serrata* (Forsk.) . Experientia. 48:498-499.
- Miller, V.H., R.S. Ballback, G.B. Pauley and Krassner, S.M. 1972. A preliminary physicochemical characterization of an agglutinin found in the hemolymph of the crayfish *Procambarus clarkii*. J. Invertebr. Pathol. 19:83-93.
- Murali, S., P. Mullainadhan and Arumugam, M. 1994. A lipopolysaccharide-binding hemagglutinin with specificity for acetylated aminosugars in the serum of hermit crab *Diogenes affinis* (Henderson). J. Invertebr. Pathol. 64:221-227.
- Muramoto, K., K. Ogata and Kamiya, H. 1985. Comparison of the multiple agglutinins of the acorn barnacle *Megabalanus rosa*. Agric. Biol. Chem. 49:85-93.
- Olafsen, J.A., 1986. Invertebrate lectins: biochemical heterogeneity as a possible key to their biological function. In 'Immunity in invertebrates' (M. Brehelin, Ed.) pp. 94-111, Springer-Verlag, Berlin.
- Putnam, D.D., V. Namasivayam and Burns, M.A. 2003. Cell affinity separations using magnetically stabilized fluidized beds: erythrocyte subpopulation fractionation utilizing a lectin-magnetite support. Biotechnol. Bioeng. 81(6):650-665.
- Ratcliffe, N.A., A.F. Rowley, S.W. Fitzgerald and Rhodes, C.P. 1985. Invertebrate immunity: basic concepts and recent advances. Int. Rev. Cytol. 97:183-350.
- Richards, E.H., and Renwranz, L.R. 1991. Two lectins of the surface of *Helix pomatia* haemocytes: a Ca²⁺-

- dependent, GalNAc-specific lectin and a Ca^{2+} -independent, mannose 6-phosphate-specific lectin which recognizes activated homologous opsonins. *J. Comp. Physiol.* 161B: 43-54.
- Rögener, W., and Uhlenbruck, G. 1984. Invertebrate lectins: the biological role of a biological rule. *Dev. Comp. Immunol.* 3:159-164.
- Sarris, A.H., and Palade, G.E. 1979. The sialoglycoproteins of murine erythrocyte ghosts. *J.Biol. Chem.* 254:6724-6731.
- Schauer, R., 1982. Sialic acids: Chemistry, Metabolism and Function, Springer-Verlag, New York, 323pp.
- Sharon, N., and Lis, H. 1989. Lectins as cell recognition molecules. *Science.* 246: 227-234.
- Sharon, N., and Lis, H. 1995. Lectins - proteins with a sweet tooth: functions in cell recognition, *Essays Biochem.* 30:59-74.
- Shinya, M., A. Okamoto, H. Sago, M. Saito, Y. Akiyama, M. Kitagawa and Tanaka, T. 2004. Analysis of fetal DNA from maternal peripheral blood by lectin-polymerase chain reaction-single strand conformation polymorphism. *Congenit. Anom.* 44(3):142-146.
- Smith, V.J., and Chisholm, J.R.S. 1992. Non-cellular immunity in crustaceans. *Fish shellfish Immunol.* 2: 1-31.
- Tsuji, T., and Osawa, T. 1986. Carbohydrate structures of bovine submaxillary mucin. *Carbohydr. Res.* 151: 391-402
- Tyler, A., and Metz, C.B. 1945. Natural heteroagglutinins in the serum of the spiny lobster, *Panulirus interruptus*. I. Taxonomic range of activity, electrophoretic and immunizing properties. *J.Exp. Zool.* 100: 387-406.
- Tyler, A., and Scheer, B.T. 1945. Natural heteroagglutinins in the serum of the spiny lobster, *Panulirus interruptus*. II. Chemical and antigenic relation to blood proteins. *Biol. Bull.* 89:193-200.
- Vasta, G.R., and Cohen, E. 1984. Carbohydrate specificities of *Birgus latro* (coconut crab) serum lectins. *Dev. Comp. Immunol.* 8:197-202.
- Vasta, G.R., G.W. Warr and Machalonis, J.J. 1983. Serological characterization of humoral lectins from the fresh water prawn *Macrobrachium rosenbergii*. *Dev. Comp. Immunol.* 7:13-20.
- Vasta, G.R., 1991. The multiple biological roles of invertebrate lectins: their participation in nonself recognition mechanisms. In 'Phylogenesis of Immune Functions' (G.W. Warr & N. Cohen, Ed). pp. 73-101, CRC Press, Boca Raton, Florida.
- Weis, W.I., 1997. Cell-surface carbohydrates recognition by animal and viral lectins. *Curr. Opin. Struc. Biol.*, 7:624-630.
- Wilson, R., C. Chen and Ratcliffe, N.A. 1999. Innate immunity in insects: the role of multiple, endogenous serum lectins in the recognition of foreign invaders in the cockroach, *Blaberus discoidalis*. *J. Immunol.* 162:1590-1596.
- Yeaton, R.W., 1981b. Invertebrate lectins II; Diversity of specificity, biological synthesis and function in recognition. *Dev. Comp. Immunol.* 5: 535-545.
- Zhang, Q., M. Tector and Salter, R.D. 1995. Calnexin recognizes carbohydrate and protein determinants of class I major histocompatibility complex molecules. *J. Biol. Chem.* 270:3944-3948.