

Effects of Bipolar Relapse on Bipolar Patients in Moi Teaching and Referral Hospital



Joy Muhia, Lynn Waithera, Khira Geoffrey Omega, Beldine FKwamboka, Rogers Songole* and Edith Kwoba

Clinical Psychology, Senior Lecturer in Mental Health, MOI University, Kenya

Received: May 04, 2018; Published: May 10, 2018

*Corresponding author: Rogers Songole, Clinical Psychology, Senior Lecturer in Mental Health, MOI University, Kenya, Tel: +254721208676; Email: rogerssongole@gmail.com

Abstract

Background: Social support plays a crucial role in the management of mental health conditions such as bipolar. Objective: The purpose of this study is to determine the effects of bipolar relapse in patients at MTRH.

Methods: The study design that was used is a retrospective study which targeted bipolar inpatients at the mental unit in Moi Teaching and Referral Hospital. Patients' hospital records for the past 5 years were reviewed by use of a chart review.

Results: The findings revealed that majority of the patients agreed that relapse affected their work, relationship with family, friends and colleagues.

Conclusion: It was concluded that Bipolar has certain effects to the day to day life conditions. As revealed in the study, most of the patients said that it affected their work/colleagues, patient family, relationships and problems in law to some extent.

Recommendation: The study recommended that families with people suffering from bipolar disorder are urged to learn their patients and handle them with great care to avoid adverse effects in their daily living.

Introduction

Bipolar disorder statistics from the World Health Organization (WHO) indicate bipolar mood disorder is the 6th leading cause of disability in the world. Social support plays a crucial role in the management of mental health conditions. Bipolar affects the community at large because all need to make certain adjustments so as to help support and accommodate the bipolar patient. Bipolar patients need social support from family, colleagues and friends and this plays a crucial role. The support system also need to know the different roles they ought to play so as to ensure a good prognosis for the patient while at the same time manage to take care of themselves.

Problem Statement

Bipolar causes significant psychosocial morbidity, because it frequently affects patients' relationships with family members and colleagues at work (Vornik & Hirschfeld, 2005). It also leads to increased family burden due to the increased cost of care and need for monitoring and support. This research thus aims to identify the effects that lead to relapse in bipolar patients and inform management interventions that will help reduce the rate of relapse.

Justification

This study is of significance to health workers, informal caregivers, families, colleagues, relatives and friends of those with bipolar so as to create awareness about the effects of bipolar among all those affected. It will also be important to other research as an inclusion to current literature.

Research Objectives

A. Broad objective

To determine effects of bipolar relapse on the patient.

B. Specific objectives

- i. To determine the effects of bipolar on patients' work
- ii. To determine the effects of bipolar on patients' family
- iii. To determine the effects of Bipolar on school
- iv. To determine the effects of bipolar on patients' relationships

Literature Review

Three types of life events have been studied in relation to BMD: negative, social rhythm disrupting, and goal-attainment [1]. Negative life events, low social support and low self-esteem predict bipolar depression but not manic symptoms [2]. Mania has been found to be preceded by schedule-disrupting life events[3]. Current findings indicate that manic symptoms are predicted by goal-attainment life events [4]. Overall, studies of life events have found that bipolar individuals experience increased stressful events prior to first onset and recurrences of mood episodes. Moreover, most studies have found that negative life events precede the manic/hypomanic as well as the depressive episodes of bipolar individuals [5-6]. Social Rhythm Stability Hypothesis (SRS) is one of the hypotheses that have been studied in relation to BMD. According to SRS, the core problem in BMD is instability of regular daily patterns of activity. Ehlers and others coined a new term - Zeitstorer - to describe a person or a social demand that throws off the regularity of the daily pattern of activity.

Whenever you start a new relationship, get a new job, buy a new pet, or have a baby, your schedule is disrupted. If you have BMD, this will result in wild cycling until you get used to the new routine. It has been shown long ago that circadian rhythm disturbances are both causes and symptoms of BMD. During depressive episodes, the phase is advanced - you usually become more of a "lark", you wake up earlier and have a lesser total amount of activity per day. During manic episodes, one is more of an "owl", staying up late and increasing total daily activity [5]. Another important aspect of an individual's current environment that affects the course of BMD is supportive or non-supportive interpersonal relationships. Social support from family and friends can buffer against the deleterious effects of stress or directly enhance functioning among bipolar individuals, whereas high expressed emotion or EEQ which refers to high levels of criticism, hostility, and/or emotional over involvement from a caregiving relative (typically a parent or spouse) during or immediately following a patient's acute episode of illness, from family members can provide additional stress and worsen the course of BMD. Four cross-sectional and one retrospective study found that bipolar individuals experience less social support than various control groups and that low social support is associated with mood episode relapses. A prospective study also found that poor social support predicts greater relapses and longer time to recovery [6].

BMD has far reaching effects, both into the lives of patients and those around them. BMD affects work, school, relationships, physical health and many other aspects of everyday life. In fact, in the early 1990s, it was calculated that the loss of productivity due to BMD cost \$15.5 billion annually. The most severe effect of BMD is suicide. Unfortunately, 25% - 50% of people with BMD attempt suicide and 11% commit suicide [7]. Psychological effects of BMD in the patient are broken down by the type of episodes the person is experiencing. Symptoms during mania/ hypomania may include: auditory and visual hallucinations, delusions including delusions of grandeur and thoughts that objects are sending special messages,

intense anxiety, agitation, aggression, paranoia, obsessive worried thoughts and feelings like feeling the need to check something, feeling like life is spinning out of control, heightened mood, exaggerated optimism and self-confidence and racing thoughts like rapidly changing streams of thought and being easily distractible. Vast psychological effects are also seen during depressive episodes: prolonged sadness, feeling helpless, hopeless and worthless, feelings of guilt, pessimism, indifference, reoccurring thoughts of death and suicide, inability to concentrate, indecisiveness and inability to take pleasure in former interests [7].

The physical effects of BMD come from both the disorder itself as well as its indirect effects. For example, many people with BMD also develop substance abuse problems in an attempt to medicate the symptoms of BMD. The effects of BMD can also result in negative productivity - in depressive episodes, but productivity can increase during hypomanic episodes. This irregularity often leads to job loss and emotional instability results in relationship loss. Physical effects of BMD include: increased physical and mental activity and energy; hyperactivity, significant changes in appetite and sleep patterns, trouble breathing, racing speech, social withdrawal, loss of energy, persistent lethargy; aches and pains, unexplained crying spells, poor overall health, weight gain, blood pressure, heart problems and diabetes [7]. When BMD is not properly controlled with medical treatment, the family may experience: Emotional distress such as guilt, grief, and worry, disruption in regular routines, having to deal with bizarre or reckless behavior, financial stresses as a result of reduced income or spending sprees, strained marital or family relationships, changes in family roles, difficulty in maintaining relationships outside the family and health problems as a result of stress [8]. According to a research by (Chatterton, et.al, 2008), bipolar families used significantly more health care resources and incurred significantly greater costs, although nearly 90% of these resources and costs were used to treat conditions other than BMD. The impact of mental disorders on communities is large and manifold. There is the cost of providing care, the loss of productivity, and some legal problems (including violence) associated with some mental disorders, though violence is caused much more often by "normal" people than by individuals with mental disorders [9].

Research Methodology

Study Design

The study was a retrospective study. Charts of the patients admitted in the mental health unit from 1st January 2012 to 31st December 2016 were reviewed.

Study Area

The research was carried out at the Moi Teaching and Referral Hospital at the Mental Health Unit in Eldoret.

Background of Study

Areathe Moi Teaching and Referral hospital is a government hospital located in Western Kenya, Rift Valley Province UasinGishu County, Eldoret Town along Nandi Road. It was established in 1916

with a bed capacity of 60 to cater for the African health needs. It later served as a District Hospital before attaining referral status in 1998 [10]. MTRH is a teaching and referral hospital hosting the Moi University schools of Medicine, Nursing, Dentistry and Public Health. The hospital serves a catchment area of approximately 2 million people. The hospital has an 800-bed capacity and receives patients from Western Kenya, parts of Eastern Uganda, and Southern Sudan. The hospital's major departments are Mental Health, Pediatrics, Surgery, Obstetrics and Gynecology and Internal Medicine. AMPATH is a specialized departmental wing of Moi Teaching and Referral Hospital (MTRH) that deals with HIV/AIDS and other chronic diseases. The MTRH mental unit serves an inpatient population of approximately 60 patients. Common diagnosis of patients admitted in the mental unit include schizophrenia, BMD and substance related disorders

Sampling

A. Target Population

Our target population comprised of approximately 1140 inpatients managed for BMD at the mental health unit.

B. Eligibility

a. Inclusion Criteria

- i. Charts of patients managed for bipolar mood disorder
- ii. The patients who were seen in the duration of the past 5 years

b. Exclusion Criteria

- i. Files missing important information relevant to the study

Study Population

The study populations were in-patients who have been admitted in the mental unit as a result of BMD from 1st January 2012 to 31st December 2016. According to the medical records, there are at least nineteen BMD admissions monthly in the mental unit. Due to this the study population for the period of study of 5 years will be 1140 patients.

Sample Size

The study was a census therefore a sample size was not necessary. All files within the stipulated period was reviewed

Sampling Technique

A sampling technique was not required as the study is a census.

A. Data Collection

a. Data Collection Tools

This was done with the use of a data extraction sheet to collect relevant data.

b. Data collection techniques

Selection of charts was done by checking the mental health unit records office and screening if the charts fulfill the inclusion criteria required. Only charts that have fulfilled the inclusion criteria were assessed.

c. Data Analysis

The data collected were both quantitative and qualitative. The quantitative data were analyzed into percentages, pie charts, tables and graphs. The qualitative data was organized into themes that were further be linked to one another for a comprehensive explanation. The quantitative data were analyzed digitally with the help of Microsoft excel and Statistical Package for the Social Sciences (SPSS) data package.

d. Data Presentation

Tables, charts and graphs were used to present in the final report. Both hard copy and soft copy of the final report was availed to COBES committee, students and any authorized individual or institution.

e. Data Storage

All electronic information was stored in a password protected computer and all hard copies of patient information was kept in a locked cabinet that can only be accessed by agreed members of the research team.

f. Ethical Considerations

Ethical approval was sought from the Institutional Research and Ethics Committee (IREC) and from Moi Teaching and Referral Hospital before conducting the study. The patient's identity was kept confidential by the use of numbers and not names to prevent disclosure. The research was carried out under keen guidance and observation from the supervisors and other qualified personnel on the ground.

Findings

Bipolar disorder is believed to have some effects in the daily living activities of patients. The study assessed these effects and found the following:

A. Patient's work/colleagues

Relapse effects on patient's work/colleagues are presented in Table 1 The study found out that majority of the patients agreed that relapse affected their work and also with colleagues, while a small majority of 42.8% did not experience any effect on work or colleagues.

Table 1: Patient's work/colleagues.

Work/colleagues	Frequency	Percent
No	416	42.8
Yes	557	57.2
Total	973	100.0

B. Patients' family

Table 2: effect on patients' family.

Family	Frequency	Percent
No	141	14.5
Yes	832	85.5
Total	973	100.0

Relapse effects on patients' family are presented in Table 2. As depicted from the table majority of the patients 85.5% noted that the relapse had a major effect on their families, while 14.5% said that it didn't have any effect on their families.

C. Effect on School

Relapse effect on school of the patient is presented in Table 3 below. The findings revealed that majority of the respondents 69.4% noted that the bipolar relapse did not have any effect on their studies while 30.6% of the respondents noted that it had a direct effect in their education.

Table 3: Effect on school.

Effect on school	Frequency	Percent
No	675	69.4
Yes	298	30.6
Total	973	100.0

D. Effect on patient's friends and relationships

Table 4 Effect on Patient's friends and relationships. As revealed in the study majority of the respondents 78% agreed that the relapses had effects on their friends and relationships while 22% did not feel any effect.

Table 4: Effect on Patient's friends and relationships.

Patient's friends and relationships	Frequency	Percent
No	214	22.0
Yes	759	78.0
Total	973	100.0

E. Problems with the law

The findings revealed that majority 85.5% of the respondents did not have a problem with the law while 14.5% experienced problems with the law (Table 5).

Table 5: Problems with the law.

Problems with the law	Frequency	Percent
No	832	85.5
Yes	141	14.5
Total	973	100.0

F. Effect on Suicide

The study revealed that the patients 86.7% did not encounter any suicidal issues while 13.3% experienced such (Table 6).

Table 6: Effect on Suicide.

Suicide	Frequency	Percent
No	844	86.7
Yes	129	13.3
Total	973	100.0

Discussion

The study found out that majority of the patients agreed that relapse affected their work and also relationships with colleagues. Majority of the patients noted that the relapse had a major effect on their families, while a few said that it didn't have any effect on their families. Studies indicate that when BMD is not properly controlled with medical treatment, the family may experience: Emotional distress such as guilt, grief, and worry, disruption in regular routines, having to deal with bizarre or reckless behavior, financial stresses as a result of reduced income or spending sprees, strained marital or family relationships, changes in family roles, difficulty in maintaining relationships outside the family and health problems as a result of stress [8]. The findings revealed that majority of the respondents noted that the bipolar relapse did not have any effect on their studies while only a few of the respondents noted that it had a direct effect in their education.

This is because majority of them got their first symptoms after they had either dropped out or had completed their education. This is attributed by the high number of individuals who had primary and secondary education as their highest level of education. This reveals that there was a high dropout rate. This is in agreement with a study carried out by Holtzman, Lolich, Ketter, & Vazquez whereby 20% of individuals had not completed primary education. Majority of the respondents agreed that the relapses had effects on their friends and relationships while a few did not feel any effect. This is brought about by the patient withdrawing from their environment and the presence of stigma that is associated with mental health. According to Latalova, Ociskova, Prasko, & Dana Kamaradova, stigmatization of patients with mental disorder is often presented like a stereotypical anticipation of unpredictability, danger, irresponsibility, and uncontrollability. Consistent with these negative anticipations, the behavior of the environment to the patient changes and the general public as a group keeps a significant distance. The patient becomes isolated with his or her problems.

The findings revealed that majority of the respondents did not have a problem with the law while a few experienced problems with the law. The study revealed that the patients did not encounter any suicidal issues while a few experienced such. According to World Health Organization [9], the impact of mental disorders on communities is large and manifold. There is the cost of providing care, the loss of productivity [10], and some legal problems (including violence) associated with some mental disorders, though violence is caused much more often by "normal" people than by individuals with mental disorders [11]. Previous studies have shown that individuals currently undergoing a depressive episode of bipolar disorders tend to withdraw from daily activities, spend less time working and with friends and more time alone at home, and spend time in passive leisure activities rather than healthy controls Barge-Schaapveld et al. Patients with bipolar disorders are also reported to make more use of avoidance and support-seeking strategies when coping with the stresses of everyday life than do non-depressed individuals Coyne et al.

Episodes of depression continue to influence daily life activities even after recovery. Focus groups of individuals recovered from a depressive episode indicated that they actively avoided the demands of work and close relationships out of fear of relapse. Coyne and Calarco, a result that has been replicated in questionnaire studies by Coyne et al. and Kirk et al. Given the high risk of relapse and the often-devastating effects of full-blown episodes, it can be expected that individuals with a history of bipolar disorder also develop adaptive strategies involving coping and time use to regulate exposure and responses to daily stress [12].

Conclusion and Recommendations

Bipolar disorder was concluded to have certain effects on the day to day life conditions. As revealed in the study, most of the patients said that it affected their work/colleagues, patient family, relationships and problems in law to some extent. From the study findings, the study made a recommendation that families with people suffering from bipolar disorder are urged to learn their patients and handle them with great care to avoid adverse effects in their daily living.


References

- Miklowitz DJ, Johnson SL (2009) Social and Familial Factors in the Course of Bipolar Disorder. *Clin Psychol (New York)*, USA 16(2): 281-296.
- Johnson SL, Winett CA, Meyer B, Greenhouse WJ, Miller I (1999) National Center for Biotechnology Information, NCBI.
- Malkoff Schwartz S, Frank E, Anderson B, Sherrill JT, Lori S, et al (1998) The JAMA Network.
- Johnson SL, Sandrow D, Meyer B, Winters R, Miller I, et al. (2000) National Center for Biotechnology Information.
- Zivkovic B (2006) CIRCADIANA, Coloumbia.
- Alloy LB, Abramson LY, Urosevic S, Walshaw PD, Nusslock R, et al. (2006) The psychosocial context of bipolar disorder: Enviromental, cognitive and developmental risk factors. *Clinical Psychology Review* 25(8): 1043-1075.
- Tracy N (2012) Healthy Place, USA.
- (2008) Mood Disorders Association of BC Heretohelp. Coloumbia.
- (2001) The World health report: 2001, Mental health: new understanding, new hope. Geneva, Switzerland, World Health Organization, London.
- MTRH (2016) Hospital Background. Moi Teaching & Referral Hospital, USA.
- Aillon JL, Ndetei DM, Khasakhala L, Ngari WN, Achola HO, et al. (2014) Prevalence, types and comorbidity of mental disorders in a Kenyan primary health centre 49(8): 1257-1268.
- Gitlin MJ, Swendsen J, Heller TL, Hammen C (1995) Relapse and impairment in bipolar disorder. *AM J Psychiatry* 0152: 1635-1640.



This work is licensed under Creative Commons Attribution 4.0 License

Submission Link: <https://biomedres.us/submit-manuscript.php>



Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles

<https://biomedres.us/>