

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/355187001>

# Phytochemicals, polyphenols content, in vitro antioxidant and antibacterial activities of *Albizia coriaria* Welw ex. Oliver flowers

Article in *Medicine Research* · January 2023

DOI: 10.53388/TMR20220531001

CITATIONS

2

READS

122

3 authors:



**Timothy Omara**

University of Natural Resources and Life Sciences Vienna

97 PUBLICATIONS 582 CITATIONS

[SEE PROFILE](#)



**Ambrose Kiprof**

Moi University

60 PUBLICATIONS 493 CITATIONS

[SEE PROFILE](#)



**Viola Jepchumba Kosgei**

University of Lorraine

17 PUBLICATIONS 102 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Chemical composition of *Pinus caribaea* Morelet var. *hondurensis* needles [View project](#)



Hybridization of Polyester Composites using Natural Fibres [View project](#)

# Phytochemicals, polyphenols content, in vitro antioxidant and antibacterial activities of *Albizia coriaria* Welw ex. Oliver flowers

Timothy Omara<sup>1,2,3,4\*</sup>, Ambrose K. Kiprop<sup>1,2</sup>, Viola J. Kosgei<sup>1,2</sup>

<sup>1</sup>Department of Chemistry and Biochemistry, School of Sciences and Aerospace Studies, Moi University, P. O. Box 3900-30100, Eldoret, Kenya. <sup>2</sup>Africa Center of Excellence II in Phytochemicals, Textile and Renewable Energy (ACE II PTRE), Moi University, P. O. Box 3900-30100, Eldoret, Kenya. <sup>3</sup>Institute of Chemistry of Renewable Resources, Department of Chemistry, University of Natural Resources and Life Sciences Vienna, 3430, Tulln, Austria. <sup>4</sup>Testing Department, Uganda National Bureau of Standards, Bweyogerere Industrial and Business Park, P.O. Box 6329, Kampala, Uganda.

\*Corresponding to: Timothy Omara, Department of Chemistry and Biochemistry, School of Sciences and Aerospace Studies, Moi University, P. O. Box 3900-30100, Eldoret, Kenya. E-mail: [prof.timo2018@gmail.com](mailto:prof.timo2018@gmail.com); [prof.timo2018@mu.ac.ke](mailto:prof.timo2018@mu.ac.ke).

## Author contributions

Timothy Omara was responsible for sample collection, methodology, sample and data analyses, and original draft writing; Ambrose K. Kiprop and Viola J. Kosgei were responsible for reviewing, editing, and supervision.

## Competing interests

The authors declare no conflicts of interest.

## Acknowledgments

The authors are indebted to Laboratory technicians of Moi University (Paul Kipkorir, Peter Biwott and Florence A. Opondo) for their assistances during this research. This research was funded by the World Bank and Inter-University Council of East Africa (IUCEA) through the Africa Center of Excellence II in Phytochemicals, Textile and Renewable Energy (ACE II PTRE) hosted at Moi University, Kenya (Credit No. 5798-KE). The funders had no role in the design of the study; collection, analyses or interpretation of data; writing of the manuscript or the decision to publish the results.

## Abbreviations

*A. coriaria*, *Albizia coriaria* Welw ex. Oliver; TFC, total flavonoid content; DPPH, 2,2-diphenyl-1-(2,4,6-trinitrophenyl) hydrazyl; IC<sub>50</sub>, half-maximal inhibitory concentration; ZOI, zone of inhibition; TPC, total polyphenolic content.

## Peer review information

Traditional Medicine Research thanks Bo Hu, Virender Kumar and Everton da Silva Santos for their contribution to the peer review of this paper.

## Citation

Omara T, Kiprop AK, Kosgei VJ. Phytochemicals, polyphenols content, in vitro antioxidant and antibacterial activities of *Albizia coriaria* Welw ex. Oliver flowers. *Tradit Med Res.* 2023;8(2):7. doi: 10.53388/TMR20220531001.

Executive editor: Rui-Wang Zhao.


Received: 31 May 2022; Accepted: 20 July 2022; Available online: 27 July 2022.

© 2023 By Author(s). Published by TMR Publishing Group Limited. This is an open access article under the CC-BY license. (<https://creativecommons.org/licenses/by/4.0/>)

## Abstract

*Albizia coriaria* Welw ex. Oliver (hereafter *A. coriaria*) is a revered medicinal tree whose organs are used by indigenous African societies for managing various ailments. Herein, we assessed the secondary metabolites, total polyphenolics, free radical scavenging and antimicrobial activities of *A. coriaria* flowers. The dry powdered sample was extracted serially with ethyl acetate, ethanol and distilled water. Results of classical phytochemical screening showed that saponins, phenols, alkaloids, flavonoids, tannins and terpenes were the dominant phytochemicals. The highest total polyphenolic content ( $10.37 \pm 0.02$  mg QE/g DW and  $72.09 \pm 0.09$  mg GAE/g DW) and free radical scavenging potential (IC<sub>50</sub> =  $24.23 \pm 0.04$  mg/mL) were for the ethyl acetate extract. It also had a higher bacteriostatic activity than the ethanolic extract against *Pseudomonas aeruginosa*, *Salmonella typhi*, *Escherichia coli* and *Staphylococcus aureus* with diameters of inhibition ranging from  $7.00 \pm 0.00$  to  $22.00 \pm 1.73$  mm. The aqueous extract showed no appreciable antibacterial effect on the tested microorganisms. This is the first report which support the use of floral extracts of this species in the management of ailments in African traditional medicine.

**Keywords:** bacteriostatic; *Albizia coriaria*; DPPH assay; traditional medicine; oxidative stress



**Highlights**

For the first time, this study reports on the phytochemicals, polyphenols content, free-radical scavenging and antibacterial potential of *Albizia coriaria* floral extracts.

**Medical history of objective**

*Albizia coriaria* is a medicinal plant used across African communities for management of human and veterinary ailments. Its various parts are used for treating complex medical conditions such as ophidian bites, lameness (Butenge), meat allergy, barrenness (*Atengina ekolupana* in cows), bacterial and oxidative stress-mediated diseases.

**Background**

Challenges posed by multidrug-resistant pathogens, side effects and inaccessibility of conventional drugs have re-ignited interest in the utilization of natural products in primary health care [1]. As such, intensive research has focussed on the documentation of traditionally used natural products, their chemical composition, pharmacological activities and structure-bioactivity relationships. On the other hand, reactive oxygen and nitrogen species owing to their high reactivity induce oxidative stress, causing destructive and irreversible damages to cellular components including DNA, proteins, lipids and lipoproteins [2]. These events are known to trigger the onset of diseases and medical conditions such as malignancies, inflammation, reperfusion injury, cardiovascular dysfunction, cirrhosis, atherosclerosis, arthritis, diabetes and emphysema [3, 4]. Commercial antioxidants have been used to scavenge free radicals in the human body [5]. Plant materials possess a host of exogenous antioxidant complexes (inform of phytochemicals) that can boost endogenous antioxidant defences of the body [6]. This could explain the widespread use of herbal remedies in the treatment of oxidative stress-mediated complications [7, 8].

*Albizia coriaria* Welw ex. Oliver (*A. coriaria*) is a widely used medicinal tree in Africa. Its flowers, whole plant, seeds, roots, leaves, stem and root barks are employed by indigenous communities for prophylaxis and management of cancers, oral, digestive, skin, respiratory and venereal ailments [9]. Though the extracts from leaves, root and stem barks have been shown to exhibit some pharmacological actions supporting the traditional claims of using of *A. coriaria* [10–23], there is hardly any study pertaining to the phytochemicals and bioactivity of its flowers [9]. Herein, a study was carried out to establish the main secondary metabolites, total phenolic and flavonoid contents, free radical scavenging and antibacterial potential of floral extracts of *A. coriaria*.

**Methods**

Analytical grade reagents, ciprofloxacin and bacteria (*Escherichia coli* ATCC 25922, *Staphylococcus aureus* ATCC 25923, *Pseudomonas aeruginosa* ATCC 27853 and *Salmonella typhi* 14028) were those used in our previous study [23], and were generously supplied by Centrihex Limited and Kobian Kenya Limited, Nairobi, Kenya.

**Sampling and extraction procedure**

*A. coriaria* flowers (2 kg) were collected from a flowering tree near Kyabazinga Way, Jinja, Uganda (0°26'34.3"N 33°12'27.3"E) [23] on Monday 8<sup>th</sup> February 2021. The sample was authenticated and a specimen (voucher No. 50997) was deposited at Makerere University Herbarium, Kampala, Uganda. The analytical sample was cleaned using water and then dried under shade for 21 days. The sample was then pulverized and 250 g of the powder was extracted serially by maceration in 500 mL of ethyl acetate, ethanol and distilled water for 2 days with intermittent agitation. The floral extracts were thereafter filtered. The organic solvent extracts were rotary evaporated under

reduced pressure whereas the aqueous extract was freeze-dried [23].

**Classical screening for secondary metabolites in the floral extracts**

The crude extracts obtained were redissolved in ethyl acetate, water and ethanol and subjected to phytochemical screening [24], the results of which were expressed as abundant, moderate, traces or absent based on colour or foam intensities [23, 25, 26].

**Quantification of total polyphenols in *A. coriaria* floral extracts**

Spectrophotometric measurement of total phenolics followed the Folin-Ciocalteu method [27], as previously used with some modifications [23]. Briefly, 0.5 mL of 0.01 mg/mL of the extracts prepared using distilled in sample vials were added to 2.5 mL of Folin-Ciocalteu reagent and left for 7 minutes. Thereafter, 2.5 mL of 6% sodium carbonate was added to the vials and incubated in the dark. After 0.5 hours, the absorbances of the solutions were read at 725 nm on an ultraviolet-visible spectrophotometer (Beckham Coulter DU 720, Brea, CA, USA). The total phenolic contents (in mg gallic acid equivalents per gram of dry weight, mg GAE/g DW) were computed from a calibration curve prepared using absorbances of standard gallic acid solutions (10 to 80 mg/mL) [23].

Total flavonoids was established using the aluminum chloride method [28] with modifications [23, 29]. Methanolic solutions of the extracts (0.01 g in 5 mL) and standard quercetin solutions (5 to 100 ppm) were prepared and 0.6 mL of each were individually added to 0.6 mL of 2% methanolic aluminum chloride solution. After incubation for 60 minutes, the absorbances were read with methanol as a blank at 420 nm. The total flavonoid content (TFC) was computed from the quercetin calibration curve and reported in milligram quercetin equivalents per gram dry weight (mg QE/g DW) [23].

**Total in vitro antioxidant activity of the floral extracts**

Antioxidant potential of the floral extracts were determined using the 2,2-diphenyl-1-(2,4,6-trinitrophenyl) hydrazyl (DPPH) assay [30] with some modifications [23]. Pipetted 75.0 µL of 1.3 mg/mL methanolic DPPH solution was transferred to test tubes with 200 µL of 10 to 60 µg/mL methanolic solutions of the extracts and ascorbic acid standard. The absorbances of the set up at 517 nm were read after incubation for 0.5 hours. The half-maximal inhibitory concentration (IC<sub>50</sub>) was computed from a plot of the percentage inhibition and concentration [23, 30].

**Antimicrobial activity of *A. coriaria* floral extracts**

The crude extracts were assessed for their inhibitory activity on *Salmonella typhi*, *Escherichia coli*, *Staphylococcus aureus* and *Pseudomonas aeruginosa* using culture-based agar disc diffusion assay as reported by Omara et al. [23]. Test extracts at 500 µg/mL were prepared in dimethyl sulfoxide and used to saturate paper discs which were subsequently air-dried. Bacterial cultures (1 × 10<sup>8</sup> CFU/mL) were subcultured onto prepared Muller Hinton broth in petri dishes. The impregnated discs were transferred carefully onto each plate in triangular formation, along with dimethyl sulfoxide and ciprofloxacin serving as negative control and positive controls, respectively. The plates were incubated at 37 °C for 24 hours after which the zone of inhibition (ZOI) was measured [23]. The minimum inhibitory and minimum bactericidal concentrations were determined for bacteria with ZOI > 12 mm after antibacterial screening following procedures used by previous authors [23, 31].

**Statistical analysis**

All analytical works were replicated at least once, and the data obtained were captured in Microsoft excel. The means of quantitative data were analyzed using one-way ANOVA followed by Tukey's post hoc test at 99% confidence interval. Further, Pearson's bivariate correlation was performed to establish any correlations between total polyphenolic content (TPC and TFC) and the IC<sub>50</sub> values of the floral extracts. All analyses were executed using GraphPad Prism statistical software (v9.1.0, GraphPad Software, San Diego, CA, USA).

## Results

### Percentage yield of the floral extracts

The extraction yields of the extracts are shown in Figure 1. Ethyl acetate extract had the highest yield ( $5.00 \pm 0.06\%$ ) vis-à-vis ethanol extract ( $2.61 \pm 0.04\%$ ) and aqueous extract ( $1.11 \pm 0.06\%$ ). One-way ANOVA confirmed that the yields differed significantly for the three solvents ( $P < 0.01$ ).

### Secondary metabolites in the floral extracts

Phenols, alkaloids, flavonoids, terpenes, saponins and tannins were the main secondary metabolites detected (Table 1). However, volatile oils, cardiac glycosides, steroids and quinones were absent. Alkaloids and flavonoids were not detected in the aqueous extract. Similarly, terpenes and tannins were absent in the ethanolic extract.

### TPC of the floral extracts

The total phenolic content of the flower extracts was computed from gallic acid curve. The TPC followed the order: ethyl acetate ( $72.09 \pm 0.09$  mg GAE/g DW) > ethanol ( $21.30 \pm 0.26$  mg GAE/g DW) > aqueous ( $15.40 \pm 0.29$  mg GAE/g DW) (Table 2), with statistically significant differences as attested to by one-way ANOVA results ( $P < 0.01$ ). Similarly, the TFC of the extracts established from the standard quercetin curve followed the sequence: ethyl acetate extract ( $10.37 \pm 0.02$  mg QE/g DW) > ethanolic extract ( $6.49 \pm 0.10$  mg QE/g DW) > aqueous extract ( $3.31 \pm 0.05$  mg QE/g DW).

### Total in vitro antioxidant activity of *A. coriaria* floral extracts


In this study, ethyl acetate extract of *A. coriaria* flowers had the lowest  $IC_{50}$  value of  $24.23 \pm 0.04$  mg/mL, followed by ethanolic extract ( $IC_{50} = 40.37 \pm 0.19$  mg/mL) and then aqueous extract ( $IC_{50} = 90.79 \pm 0.16$  mg/mL) (Table 3). However, the  $IC_{50}$  value of the standard antioxidant (ascorbic acid,  $IC_{50} = 0.17 \pm 0.01$  mg/mL) was lower than that of all the extracts. Statistical evaluation using one-way ANOVA indicated that significant differences ( $P < 0.01$ ) existed between the  $IC_{50}$  values of the extracts.

### Correlation between TPC and half maximal inhibitory concentration of *A. coriaria* floral extracts

Correlative analysis between the TPC and TFC indicated a positive correlation while only negative correlations were observed between TPC and  $IC_{50}$  values, and TFC and  $IC_{50}$  values obtained in DPPH assays (Table 4).

### Antibacterial activity results

Only organic solvent extracts exerted bacteriostatic effect on the tested bacteria, with ethyl acetate extract having the highest ZOI of  $22.00 \pm 1.73$  mm (Table 5). In regards to MIC and MBC, *Escherichia coli* had the lowest MIC (250  $\mu$ g/mL) for the ethyl acetate extract compared to that of *Pseudomonas aeruginosa* (500  $\mu$ g/mL). The MBC of the ethyl acetate extract was 500  $\mu$ g/mL and greater than 500  $\mu$ g/mL against *Escherichia coli* and *Pseudomonas aeruginosa*.



**Figure 1** Percentage yield of different solvent extracts of *Albizia coriaria* Welw ex. Oliver flowers

**Table 1** Phytochemicals screened in the organic solvent and aqueous floral extracts of *A. coriaria* sampled from Jinja, Uganda

Metabolite	Solution <sup>1</sup>	Result <sup>2</sup>
Phenols	EtAC	++
	EtOH	+
	AQ	+
Alkaloids	EtAC	+++
	EtOH	++
	AQ	-
Cardiac glycosides	EtAC	-
	EtOH	-
	AQ	-
Flavonoids	EtAC	+
	EtOH	++
	AQ	-
Saponins	EtAC	+
	EtOH	++
	AQ	+
Quinones	EtAC	-
	EtOH	-
	AQ	-
Steroids	EtAC	-
	EtOH	-
	AQ	-
Volatile oils	EtAC	-
	EtOH	-
	AQ	-
Terpenes	EtAC	++
	EtOH	-
	AQ	+
Tannins	EtAC	+
	EtOH	-
	AQ	++

<sup>1</sup>Extracts: EtAC = ethyl acetate, EtOH = ethanol, AQ = aqueous.

<sup>2</sup>The symbols: +++, ++, +, and - denotes abundant, moderate, traces and absent, respectively. *A. coriaria*, *Albizia coriaria* Welw ex. Oliver.

**Table 2** TPC of extracts of *A. coriaria* flowers from Jinja, Uganda

Extract	Total phenolic content (mg GAE/g DW)	Total flavonoid content (mg QE/g DW)
Ethyl acetate	$72.09 \pm 0.09$	$10.37 \pm 0.02$
Ethanolic	$21.30 \pm 0.26$	$6.49 \pm 0.10$
Aqueous	$15.40 \pm 0.29$	$3.31 \pm 0.05$

Results are presented as means  $\pm$  standard deviations of triplicates. *A. coriaria*, *Albizia coriaria* Welw ex. Oliver; TPC, total polyphenolic content.

**Table 3** DPPH results for inhibition concentration at 50% of *A. coriaria* flowers

Extract/standard	$IC_{50}$ (mg/mL)
Ethyl acetate	$24.23 \pm 0.04$
Ethanolic	$40.37 \pm 0.19$
Aqueous	$90.79 \pm 0.16$
Ascorbic acid	$0.17 \pm 0.01$

Results are presented as means  $\pm$  standard deviations of triplicates. *A. coriaria*, *Albizia coriaria* Welw ex. Oliver; DPPH, 2,2-diphenyl-1-(2,4,6-trinitrophenyl) hydrazyl;  $IC_{50}$ , half-maximal inhibitory concentration.

**Table 4 Pearson's bivariate correlation coefficients between TPC and IC<sub>50</sub> values of *A. coriaria* floral extracts**

Parameter	Total flavonoid content	Antioxidant activity
Total phenolic content	R = 0.932, P = 0.118	R = -0.753, P = 0.457
Total flavonoid content	1.000	R = -0.941, P = 0.220

*A. coriaria*, *Albizia coriaria* Welw ex. Oliver; TPC, total polyphenolic content; IC<sub>50</sub>, half-maximal inhibitory concentration.

**Table 5 Mean diameters of inhibition by the floral extracts**

Test solution	<i>Escherichia coli</i> (mm) <sup>1</sup>	<i>Staphylococcus aureus</i> (mm)	<i>Pseudomonas aeruginosa</i> (mm)	<i>Salmonella typhi</i> (mm)
Ethyl acetate extract	22.00 ± 1.73	11.00 ± 2.65	19.00 ± 2.10	7.00 ± 0.00
Ethanol extract	8.00 ± 1.73	3.00 ± 1.00	9.00 ± 2.65	5.00 ± 0.02
Aqueous extract	0.00 ± 0.00	0.00 ± 0.00	0.00 ± 0.00	0.00 ± 0.00
Ciprofloxacin (positive control)	14.00 ± 2.10	12.00 ± 0.01	31.00 ± 0.11	20.00 ± 1.53

<sup>1</sup>Results are average ± standard error of three experiments. Dimethyl sulfoxide did not inhibit bacterial growth.

## Discussion

Extraction of phytochemicals from *A. coriaria* flowers was done by maceration and ethyl acetate had the highest yield (5.00 ± 0.06%) vis-à-vis ethanol (2.61 ± 0.04%) and distilled water (1.11 ± 0.06%) (Figure 1). The yields obtained were comparable to 2.98% to 5.98% for *A. coriaria* leaf extracts [23] but lower than 8.3% to 12.7%, 11.9% and 10.88% for organic solvent extracts of its stem bark [31, 32] and leaves [23]. Statistical evaluation results indicated that the yields differed significantly for the three solvents ( $P < 0.01$ ), implying that ethyl acetate was the best solvent for recovering bioactive compounds from the flowers. The low yield of aqueous extract is most likely due to the effect of serial extraction i.e., most bioactive compounds had already been extracted by the organic solvents. Nevertheless, the results attested to the fact that *A. coriaria* flowers possesses both polar and non-polar phytochemicals, which is concordant with reports on its leaves [23] and stem bark extracts [15, 31, 32]. On the other hand, phenols, alkaloids, flavonoids, terpenes, saponins and tannins were the main secondary metabolites detected (Table 1). These groups of secondary metabolites were previously detected in *A. coriaria* stem bark and leaves [14, 17, 20, 22, 23, 33, 34] and floral extracts of *Albizia lebbek* (L.) Benth [35–37].

In regards to polyphenols, previous studies [15, 23] reported that TPC of ethanolic extracts of *A. coriaria* leaves (67.04 to 101.72 mg GAE/g DW) and stem bark (28.37 mg CAE/g E) were averagely higher than those of their corresponding ethyl acetate extracts (10.93 to 60.69 mg GAE/g DW and 28.36 mg CAE/g E). A plausible explanation for the higher TPC of the ethyl acetate extract than the ethanolic extract in this study could be due to extraction of majority of the polyphenols in the flower powder being non-polar and thus were extracted more by ethyl acetate than ethanol and water. In the case of water, the lower TPC could be because most of the polar polyphenols were already extracted by ethanol which is relatively polar. In addition, water is also reported to extract even non-active compounds in plant matrices, which may not constitute the TPC [23, 38]. In the same way, the TFC of the extracts followed the sequence: ethyl acetate extract > ethanolic extract > aqueous extract (Table 2). The results indicated that *A. coriaria* flowers have more flavonoid aglicones [39]. Omara et al. [23] reported that ethanolic extracts of *A. coriaria* leaves housed the highest quantity of total flavonoids (8.63 to 13.23 mg QE/g DW) vis-à-vis the ethyl acetate extracts (0.55 to 9.66 mg QE/g DW) and aqueous extracts (2.74 to 3.36 mg QE/g DW). As highlighted for the TPC, lower TFC of the aqueous extracts is reasonably due to the initial extraction of most flavonoids by the organic solvents. It is worth noting the TFC of the extracts were lower than their TPCs, corroborating the fact that most flavonoids are in essence phenolics [40]. Generally, extraction of polyphenols from plant matrices are influenced by the nature of the polyphenols, the extraction solvents' polarity, time and temperature used [39]. This explains, in part, the observed disparities in the polyphenolic content of the floral extracts.

Plant extracts elicit antioxidant activity in vitro owing to their reducing, chelating or oxygen quenching activities [41]. Antioxidants are compounds with characteristic potential to delay or inhibit oxidative damage to other molecules. This is primarily through

trapping of free radicals. Phytochemicals such as polyphenols and flavonoids abundant in plants are known to scavenge free radicals, including hydroperoxides and peroxides [41]. In this study, ethyl acetate extract had the lowest IC<sub>50</sub> value, followed by ethanolic and aqueous extracts (Table 3). These differences in the antioxidant potential may be attributed to the variations in the bioactive compounds in the extracts owing to the differences in the extracting power of the solvents used. In comparison with previous reports, the antioxidant potential obtained herein surpassed those reported for organic extracts of *A. coriaria* stem bark (with half effective concentrations of 0.02298 mg/mL and 0.01839 mg/mL) [15] but comparatively lower than those of organic solvent and aqueous extracts of *A. coriaria* leaves (IC<sub>50</sub> = 18.65 to 29.66 mg/mL) [23]. The lower free radical scavenging activity (highest IC<sub>50</sub> value) of aqueous extracts is because water solvates bioactive compounds along with inactive phytochemicals in matrices which may solely act as matrix interferences [38].

Pearson's correlative analysis between the TPC and TFC indicated a positive correlation ( $R = 0.932, P = 0.118$ ). This is arguably owing to the chemistry that polyphenolic compounds are constituted by both flavonoid- and non-flavonoid compounds [23, 26, 42]. Further correlative analysis between the TPC and IC<sub>50</sub> values gave a negative association ( $R = -0.753, P = 0.457$ ), which was also observed between TFC and IC<sub>50</sub> values ( $R = -0.941, P = 0.220$ ). This result is concordant with a preceding report which observed poor and/or negative correlation between TPC or TFC and the corresponding free radical scavenging potential (EC<sub>50</sub> and IC<sub>50</sub> values) [15, 23]. Overall, poor and/or negative correlations between IC<sub>50</sub> values of the extracts, and their TPC or TFC explains the chemistry that reduction in the DPPH radical progresses with decrement in scavenging action of the extracts. This suggests that polyphenols in the extracts are responsible for their DPPH radical scavenging potential [26, 42, 43].

In the antibacterial assay, only organic solvent extracts exerted bacteriostatic effect on the pathogens (Table 5). Based on Clinical and Laboratory Standards Institute Interpretive Criteria [44], only *Escherichia coli* and *Staphylococcus aureus* were susceptible to the ethyl acetate extract as their ZOI diameters lied within the one for conventional antibacterial drugs such as doxycycline, ampicillin and tetracycline. The results of antibacterial screening are similar to a previous observation wherein the ethyl acetate extract of *A. coriaria* stem bark elicited a higher bacteriostatic action against pathogenic bacteria vis-à-vis the methanolic extract [18]. However, Omara et al. [23] and Schultz et al. [15] reported higher bacteriostatic potency of ethanol extracts of *A. coriaria* leaves and stem bark than the corresponding ethyl acetate extracts. Interestingly, the aqueous extract in this study was inactive on all the tested pathogens, corroborating previous observations using the aqueous stem bark and leaf extracts of *A. coriaria* [18, 20, 23]. The secondary metabolites detected in the organic solvent extracts of *A. coriaria* flowers could be implicated for their bacteriostatic activities. Phytochemicals such as tannins, alkaloids, flavonoids and phenols are known antibacterial agents, either directly or through microbial proteins precipitation, intercalating microbial DNA or targeted deactivation of nucleic acid enzymes [45–47]. The bacteriostatic activities were in

correlation with phytochemical richness of the extracts. For instance, the extract obtained from maceration using ethyl acetate housed the highest levels of alkaloids, phenols, flavonoids and terpenes vis-à-vis other extracts (Table 1).

The lowest MIC was observed for *Escherichia coli* (250 µg/mL) while the MIC for *Pseudomonas aeruginosa* was 500 µg/mL. A previous report [18] reported that ethyl acetate extract of *A. coriaria* stem bark had MIC of 125 mg/mL for *Escherichia coli* and 250 mg/mL for *Pseudomonas aeruginosa*. A team investigating ethanolic extracts of *A. coriaria* stem bark had MIC > 256 µg/mL for *Pseudomonas aeruginosa* [19] while the same extract and its ethyl acetate extract had MIC of 250 and 500 µg/mL against *Staphylococcus aureus* and *Escherichia coli* [15] which are comparable to the MIC values obtained in this study. Omara et al. recently reported MIC values of 62.5–250 µg/mL for ethanol leaf extracts of this species against *Pseudomonas aeruginosa* [23].

The MBC of the ethyl acetate extract was 500 µg/mL and greater than 500 µg/mL against *Escherichia coli* and *Pseudomonas aeruginosa*. These were higher than the MBC of 125–250 µg/mL reported for the ethanolic extract of the leaves of *A. coriaria* against *Pseudomonas aeruginosa* [23]. A high MBC of 125 and >500 mg/mL were indicated for ethyl acetate stem bark extract of this species against *Escherichia coli* and *Pseudomonas aeruginosa*, respectively [18]. Overall, such higher minimum inhibitory and bactericidal concentrations emphasize that larger quantities of the floral extracts are necessary to traditionally treat the bacterial infections. This also reflected in the posology of this species as prescribed by herbalists in Uganda [48, 49].

### Conclusion

The results obtained in this study demonstrated that *A. coriaria* flowers houses phytochemicals with free radical scavenging and bactericidal activities, which is in support of its traditional utilization in the treatment of ailments in Africa. In this context, further studies assessing the toxicity, structure and bioactivity of the phytochemicals are required.

### References

- Khan SA, Imtiaz MA, Sayeed MA, Shaikat AH, Hassan MM. Antimicrobial resistance pattern in domestic animal - wildlife - environmental niche via the food chain to humans with a Bangladesh perspective; a systematic review. *BMC Vet Res*. 2020;16:302. <https://doi.org/10.1186/s12917-020-02519-9>
- Weidinger A, Kozlov AV. Biological activities of reactive oxygen and nitrogen species: oxidative stress versus signal transduction. *Biomolecules*. 2015;5:472–484. <https://doi.org/10.3390/biom5020472>
- Obakiro SB, Kiprof A, Kigundu E, et al. Traditional medicinal uses, phytoconstituents, bioactivities, and toxicities of *Erythrina abyssinica* Lam. ex DC. (Fabaceae): a systematic review. *Evid Based Complement Alternat Med*. 2021;2021:5513484. <https://doi.org/10.1155/2021/5513484>
- García-Sánchez A, Miranda-Díaz AG, Cardona-Muñoz EG. The role of oxidative stress in physiopathology and pharmacological treatment with pro- and antioxidant properties in chronic diseases. *Oxid Med Cell Longev*. 2020;2020:2082145. <https://doi.org/10.1155/2020/2082145>
- Omara T, Kagoya S, Openy A, et al. Antivenin plants used for treatment of snakebites in Uganda: ethnobotanical reports and pharmacological evidences. *Trop Med Health*. 2020;48:6. <https://doi.org/10.1186/s41182-019-0187-0>
- Kasote DM, Katyare SS, Hegde MV, Bae H. Significance of antioxidant potential of plants and its relevance to therapeutic applications. *Int J Biol Sci*. 2015;11:982–991. <https://doi.org/10.7150%2Fijbs.12096>
- Omara T, Kiprof AK, Ramkat RC, et al. Medicinal plants used in traditional management of cancer in Uganda: ethnobotanical surveys, phytochemistry and anticancer studies. *Evid Based Complement Alternat Med*. 2020;2020:3529081. <https://doi.org/10.1155/2020/3529081>
- Xu DP, Li Y, Meng X, et al. Natural antioxidants in foods and medicinal plants: extraction, assessment and resources. *Int J Mol Sci*. 2017;18:96. <https://doi.org/10.3390/ijms18010096>
- Omara T, Kiprof AK, Kosgei VJ. *Albizia coriaria* Welw ex Oliver: a review of its ethnobotany, phytochemistry and ethnopharmacology. *Adv Tradit Med*. 2021. <https://doi.org/10.1007/s13596-021-00600-8>
- Noté OP, Mitaine-Offer AC, Miyamoto T, et al. Cytotoxic acacic acid glycosides from the roots of *Albizia coriaria*. *J Nat Prod*. 2009;72:1725–1730. <https://doi.org/10.1021/np900126r>
- Muthaura CN, Keriko JM, Mutai C, et al. Antiplasmodial potential of traditional phytotherapy of some remedies used in treatment of malaria in Meru-Tharaka Nithi County of Kenya. *J Ethnopharmacol*. 2015;175:315–323. <https://doi.org/10.1016/j.jep.2015.09.017>
- Kama-Kama F, Midiwo J, Nganga J, et al. Selected ethno-medicinal plants from Kenya with in vitro activity against major African livestock pathogens belonging to the “*Mycoplasma mycoides* cluster”. *J Ethnopharmacol*. 2016;192:524–534. <https://doi.org/10.1016/j.jep.2016.09.034>
- Johns T, Faubert GM, Kokwaro JO, Mahunnah RL, Kimanani EK. Anti-giardial activity of gastrointestinal remedies of the Luo of East Africa. *J Ethnopharmacol*. 1995;46:17–23. [https://doi.org/10.1016/0378-8741\(95\)01224-2](https://doi.org/10.1016/0378-8741(95)01224-2)
- Mengesha H, Birrie H, Gundersen G. The molluscicidal effect of the bark of “bitza” tree: a local discovery towards the control of schistosomiasis in the Abay river valley of Western Ethiopia. *Ethiop J Health Dev*. 1997;11:89–92. <https://www.ejhd.org/index.php/ejhd/article/view/1032/780>
- Omara T, Kiprof AK, Kosgei VJ. Two new pentacyclic triterpenoids, an alkaloid and a long-chain fatty acid from *Albizia coriaria* (Welw ex. Oliver). *French-Ukrain J Chem*. 2022;10(01):128–141. <https://doi.org/10.17721/fujcV10I1P128-141>
- Olila D, Ziraba BR, Kamoga D. Bio-prospective studies on medicinal plants used to manage poultry diseases in the Mount Elgon region of Uganda. *Res J Pharmacol*. 2007; 1: 56-60. <http://docsdrive.com/pdfs/medwelljournals/rjpharm/2007/56-60.pdf>
- Akanga JO. Screening of anti-diarrhoea medicinal plants for in vitro antimicrobial activity against clinical and environmental enteropathogens. MSc thesis, Jomo Kenyatta University of Agriculture and Technology, Nairobi. 2008. <http://ir.jkuat.ac.ke/handle/123456789/858>
- Byamukama R, Barbara G, Namukobe J, Heydenreich M, Kiremire BT. Bioactive compounds in the stem bark of *Albizia coriaria* (Welw. ex Oliver). *Int J Biol Chem Sci*. 2015;9:1013–1024. <https://doi.org/10.4314/IJBCS.V9I2.37>
- Schultz F, Anywar G, Tang H, et al. Targeting ESKAPE pathogens with anti-infective medicinal plants from the Greater Mpigi region in Uganda. *Sci Rep*. 2020;10:11935. <https://doi.org/10.1038/s41598-020-67572-8>
- Nalubega R, Kabasa JD, Olila D, Kateregga J. Antibacterial activity and phytochemical screening of eleven plants used as poultry ethnomedicines in Southern Uganda. *Agric J*. 2011;6:303–309. <https://doi.org/10.3923/aj.2011.303.309>
- Luvonga MW. Evaluation of antimicrobial activity of some plants used by traditional healers for treatment of microbial infections in Kakamega district: Kenya. MSc thesis, Kenyatta University, Nairobi. 2007. <https://ir-library.ku.ac.ke/handle/123456789/432?show=full>

22. India J. Efficacy of some medicinal plants used in various parts of Kenya in treating selected bacterial and fungal pathogens. MSc thesis, Kenyatta University, School of Pure and Applied Sciences, Nairobi. 2015. <https://ir-library.ku.ac.ke/handle/123456789/14965>
23. Omara T, Kiprop AK, Kosgei VJ. Intraspecific variation of phytochemicals, antioxidant, and antibacterial activities of different solvent extracts of *Albizia coriaria* leaves from some agroecological zones of Uganda. *Evid Based Complement Alternat Med.* 2021;2021:2335454. <https://doi.org/10.1155/2021/2335454>
24. Trease GE, Evans WC. Phenols and phenolic glycosides. In: *Textbook of Pharmacognosy*. London: Balliesse, Tindall and Co Publishers;1989:343–383. ISBN: 97807020135770702013579
25. Gul R, Jan SU, Faridullah S, Sherani S, Jahan N. Preliminary phytochemical screening, quantitative analysis of alkaloids, and antioxidant activity of crude plant extracts from *Ephedra intermedia* indigenous to Balochistan. *Sci World J.* 2017;2017:5873648. <https://doi.org/10.1155/2017/5873648>
26. Suleiman MHA, Ateeg AA. Antimicrobial and antioxidant activities of different extracts from different parts of *Zilla spinosa* (L.) Prantl. *Evid Based Complement Alternat Med.* 2020;2020:6690433. <https://doi.org/10.1155/2020/6690433>
27. Velioglu YS, Mazza G, Gao L, Oomah BD. Antioxidant activity and total phenolics in selected fruits, vegetables, and grain products. *J Agric Food Chem.* 1998;46:4113–4117. <https://doi.org/10.1021/jf9801973>
28. Pękal A, Pyrzynska K. Evaluation of aluminium complexation reaction for flavonoid content assay. *Food Anal Methods.* 2014;7:1776–1782. <https://doi.org/10.1007/s12161-014-9814-x>
29. Chandra S, Khan S, Avula B, et al. Assessment of total phenolic and flavonoid content, antioxidant properties, and yield of aeroponically and conventionally grown leafy vegetables and fruit crops: a comparative study. *Evid Based Complement Alternat Med.* 2014;2014:253875. <https://doi.org/10.1155/2014/253875>
30. Khiya Z, Oualcadi Y, Gamar A, Berrekhis F, Zair T, EL Hilali F. Correlation of total polyphenolic content with antioxidant activity of hydromethanolic extract and their fractions of the *Salvia officinalis* leaves from different regions of Morocco. *J Chem.* 2021;2021:8585313. <https://doi.org/10.1155/2021/8585313>
31. Ganza B. Isolation and characterization of the bioactive compounds in the stem bark of *Albizia coriaria*. MSc thesis, Makerere University, Kampala. 2014. <http://makir.mak.ac.ug/handle/10570/3351>
32. Owuor BO, Ochanda JO, Kokwaro JO, et al. In vitro antiparasmodial activity of selected Luo and Kuria medicinal plants. *J Ethnopharmacol.* 2012;144:779–781. <https://doi.org/10.1016/j.jep.2012.09.045>
33. Wanyama PAG, Kiremire BT, Murumu JES, Kamoga O. Textile dyeing and phytochemical characterization of crude plant extracts derived from selected dye-yielding plants in Uganda. *Int J Nat Prod Res.* 2011;1:26–31.
34. Langat PK. Ethnobotanical study, toxicity and phytochemical screening of selected medicinal plants of Tinderet district, Nandi county, Kenya. MSc thesis, University of Nairobi, Nairobi. 2013. <http://erepository.uonbi.ac.ke/handle/11295/61411>
35. Padamanabhan V, Ganapathy M, Evanjelene VK. Preliminary phytochemical and anti-bacterial studies on flowers and pods of *Albizia lebbeck* (Benth). *Int J Emerg Technol Adv Eng.* 2013;3:541–544. [https://ijetae.com/files/Volume3Issue9/IJETAE\\_0913\\_89.pdf](https://ijetae.com/files/Volume3Issue9/IJETAE_0913_89.pdf)
36. Jasiem TM, AL-Bazaz HK, Baderden S, Nasser NM. Isolation of flavonoids and pharmacognostical study of Iraqi *Albizia lebbeck* L. *J Pharmaceut Sci Res.* 2019;11:239–242. [https://www.researchgate.net/publication/331045831\\_Isolation\\_of\\_flavonoids\\_and\\_Pharmacognostical\\_study\\_of\\_Iraqi\\_Albizia\\_lebbeck\\_L](https://www.researchgate.net/publication/331045831_Isolation_of_flavonoids_and_Pharmacognostical_study_of_Iraqi_Albizia_lebbeck_L)
37. AL-Sa'ady AJR, Abbas LMR. The effect of aqueous *Albizia* flowers extract *Albizia lebbeck* on production of certain enzymes from *Salmonella typhi*. *J Biotechnol Res Center.* 2016;10:25–30. [https://www.researchgate.net/publication/318967760\\_The\\_Effect\\_of\\_Aqueous\\_Albizia\\_Flowers\\_Extract\\_Albizia\\_lebbeck\\_on\\_Production\\_of\\_Certain\\_Enzymes\\_from\\_Salmonella\\_typhi](https://www.researchgate.net/publication/318967760_The_Effect_of_Aqueous_Albizia_Flowers_Extract_Albizia_lebbeck_on_Production_of_Certain_Enzymes_from_Salmonella_typhi)
38. Altemimi A, Lakhssassi N, Baharlouei A, Watson DG, Lightfoot DA. Phytochemicals: extraction, isolation, and identification of bioactive compounds from plant extracts. *Plants.* 2017;6:42. <https://doi.org/10.3390/plants6040042>
39. Felhi S, Daoud A, Hajlaoui H, Mnafigui K, Gharsallah N, Kadri A. Solvent extraction effects on phytochemical constituents profiles, antioxidant and antimicrobial activities and functional group analysis of *Ecballium elaterium* seeds and peels fruits. *Food Sci Technol Int.* 2017;37:483–492. <https://doi.org/10.1590/1678-457X.23516>
40. Ahmed D, Khan MM, Saeed R. Comparative analysis of phenolics, flavonoids, and antioxidant and antibacterial potential of methanolic, hexanic and aqueous extracts from *Adiantum caudatum* leaves. *Antioxidants.* 2015;4:394–409. <https://doi.org/10.3390/antiox4020394>
41. Flieger J, Flieger M. The [DPPH/DPPH-H]-HPLC-DAD method on tracking the antioxidant activity of pure antioxidants and goutweed (*Aegopodium podagraria* L.) hydroalcoholic extracts. *Molecules.* 2020;25:6005. <https://doi.org/10.3390/molecules25246005>
42. Kim JS, Lee JH. Correlation between solid content and antioxidant activities in Umbelliferae salad plants. *Prev Nutr Food Sci.* 2020;25:84–92. <https://doi.org/10.3746/pnf.2020.25.1.84>
43. Zengin G, Mahomoodally MF, Paksoy MY, et al. Phytochemical characterization and bioactivities of five Apiaceae species: natural sources for novel ingredients. *Ind Crops Prod.* 2019;135:107–121. <https://doi.org/10.1016/j.indcrop.2019.04.033>
44. Clinical and Laboratory Standards Institute. *Performance Standards for Antimicrobial Susceptibility Testing.* 27<sup>th</sup> ed. CLSI supplement M100. Wayne, PA. 2017. <https://clsi.org>
45. Pradhan D, Panda P, Tripathy G. Wound healing activity of aqueous and methanolic bark extract of *Vernonia arborea* in Wistar rats. *Nat Prod Rad.* 2009;8:6–11. <http://14.139.47.15/bitstream/123456789/3765/1/NPR%208%281%29%206-11.pdf>
46. Daglia M. Polyphenols as antimicrobial agents. *Curr Opin Biotechnol.* 2012;23:174–181. <https://doi.org/10.1016/j.copbio.2011.08.007>
47. Yi ZB, Yan Y, Liang YZ, Zeng B. Evaluation of the antimicrobial mode of berberine by LC/ESI-MS combined with principal component analysis. *J Pharm Biomed Anal.* 2007;44:301–304. <https://doi.org/10.1016/j.jpba.2007.02.018>
48. Namukobe J, Kasenene JM, Kiremire BT, et al. Traditional plants used for medicinal purposes by local communities around the Northern sector of Kibale National Park, Uganda. *J Ethnopharmacol.* 2011;136:236–245. <https://doi.org/10.1016/j.jep.2011.04.044>
49. Asiimwe S, Kamatenesi-Mugisha M, Namutebi A, Borg-Karlsson AK, Musiimenta P. Ethnobotanical study of nutri-medicinal plants used for the management of HIV/AIDS opportunistic ailments among the local communities of western Uganda. *J Ethnopharmacol.* 2013;150:639–648. <https://doi.org/10.1016/j.jep.2013.09.017>