

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/360334171>

# Clausena anisata (Willd.) Hook.f. ex Benth. (Rutaceae): Ethnomedicinal Uses, Phytochemistry, Pharmacological Activities, Toxicity and Clinical Application

Article in *Journal of Traditional and Complementary Medicine* · July 2022

DOI: 10.53388/TMR20220417001

CITATIONS

0

READS

323

4 authors:



**Timothy Omara**

University of Natural Resources and Life Sciences Vienna

97 PUBLICATIONS 581 CITATIONS

[SEE PROFILE](#)



**Ambrose Kiprof**

Moi University

58 PUBLICATIONS 493 CITATIONS

[SEE PROFILE](#)



**Viola Jepchumba Kosgei**

University of Lorraine

17 PUBLICATIONS 102 CITATIONS

[SEE PROFILE](#)



**Sarah Kagoya**

Kyambogo University

22 PUBLICATIONS 246 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Aviation Noise and Air Pollution: Results of a Study at Entebbe International Airport, Uganda [View project](#)



Artisanal and small scale gold mining in Syanyonja, Busia gold district, Eastern Uganda: Impacts on the mining population and the environment [View project](#)

# *Clausena anisata* (Willd.) Hook.f. ex Benth. (Rutaceae): ethnomineral uses, phytochemistry, pharmacological activities, toxicity, and clinical application

Timothy Omara<sup>1,2,3\*</sup>, Ambrose K. Kiprop<sup>1,2</sup>, Viola J. Kosgei<sup>1,2</sup>, Sarah Kagoya<sup>4</sup>

<sup>1</sup>Department of Chemistry and Biochemistry, School of Sciences and Aerospace Studies, Moi University, Eldoret 3900-30100, Kenya. <sup>2</sup>Africa Centre of Excellence II in Phytochemicals, Textile and Renewable Energy (ACE II PTRE), Moi University, Eldoret 3900-30100, Kenya. <sup>3</sup>Institute of Chemistry of Renewable Resources, Department of Chemistry, University of Natural Resources and Life Sciences Vienna (BOKU), Tulln 3430, Austria. <sup>4</sup>Department of Chemistry, Faculty of Science, Kyambogo University, Kampala 1, Uganda.

\*Corresponding to: Timothy Omara, Department of Chemistry and Biochemistry, School of Sciences and Aerospace Studies, Moi University, Uasin Gishu County, Academic Highway, Eldoret 3900-30100, Kenya. E-mail: [prof.timo2018@gmail.com](mailto:prof.timo2018@gmail.com); [prof.timo2018@mu.ac.ke](mailto:prof.timo2018@mu.ac.ke).

## Author contributions

Timothy Omara was responsible for the original draft writing; Ambrose K. Kiprop, Viola J. Kosgei and Sarah Kagoya were responsible for reviewing, editing, and supervision; Timothy Omara was responsible for methodology.

## Competing interests

The authors declare no conflicts of interest.

## Acknowledgments

The authors are grateful to the World Bank and the Inter-University Council of East Africa (IUCEA) for the fellowship awarded to Timothy Omara through the Africa Centre of Excellence II in Phytochemicals, Textile and Renewable Energy (ACE II PTRE) at Moi University, Kenya which made this communication possible. We thank Editage Ltd. for editing the English text of a draft of this manuscript.

## Abbreviations

*C. anisata*, *Clausena anisata* (Willd.) Hook.f. ex Benth.; *V. cholerae*, *Vibrio cholerae*; *P. acnes*, *Propionibacterium acnes*; *C. maculatus*, *Callosobruchus maculatus*; *A. obtectus*, *Acanthoscelides obtectus*; *C. quinquefasciatus*, *Culex quinquefasciatus*; *A. aegypti*, *Aedes aegypti*; *A. stephensi*, *Anopheles stephensi*; *A. variegatum*, *Amblyomma variegatum*; BW, body weight; QE, quercetin equivalent; GAE, gallic acid equivalent; TPC, total phenolic content; TFC, total flavonoid content; Eos, essential oils; DPPH, 2-diphenyl-1-picrylhydrazyl; FRAP, ferric reducing power; ABTS, 3-ethylbenzthiazoline-6-sulfonic acid; EC<sub>50</sub>, half-effective concentration; BHT, butylated hydroxytoluene; IC<sub>50</sub>, inhibitory concentration; SC<sub>50</sub>, scavenging activity; LC<sub>50</sub>, lethal concentration; DCM, dichloromethane; sPLA<sub>2</sub>, secretory phospholipase A2; 15-LOX, lipoxygenase; HL-60, human leukaemic; ACE, angiotensin-converting enzyme; ZOI, zone of inhibition; COX, cyclooxygenase; SNPs, silver nanoparticles; LD<sub>50</sub>, lethal doses; AChE, acetylcholinesterase; IL, interleukin.

## Peer review information

*Traditional Medicine Research* thanks all anonymous reviewers for their contribution to the peer review of this paper.

## Citation

Omara T, Kiprop AK, Kosgei VJ, Kagoya S. *Clausena anisata* (Willd.) Hook.f. ex Benth. (Rutaceae): ethnomineral uses, phytochemistry, pharmacological activities, toxicity, and clinical application. *Tradit Med Res* 2022;7(6):51. doi: 10.53388/TMR20220417001.

Executive editor: Chun Ling.

Received: 17 April 2022; Accepted: 02 May 2022;


Available online: 15 July 2022.

© 2022 By Author(s). Published by TMR Publishing Group Limited. This is an open access article under the CC-BY license. (<http://creativecommons.org/licenses/by/4.0/>)

## Abstract

*Clausena anisata* (Willd.) Hook.f. ex Benth. is a plant extensively used in traditional medicine. Here, a synopsis of the research on various aspects of *Clausena anisata* (Willd.) Hook.f. ex Benth. is presented. An electronic literature review revealed that formulations containing *Clausena anisata* (Willd.) Hook.f. ex Benth. are used to manage and treat diabetes, eye problems, malaria, snake envenomation, malignancies, as well as venereal, gastrointestinal, reproductive, respiratory, dermatological, and odontological ailments. To date, 90 compounds have been isolated and characterized from extracts of *Clausena anisata* (Willd.) Hook.f. ex Benth. Crude extracts and isolated compounds from *Clausena anisata* (Willd.) Hook.f. ex Benth. possess anti-inflammatory, antimicrobial, antioxidant, antiplasmodial, antiproliferative, anti-human immunodeficiency virus-1, antimycobacterial, antihypertensive, and antidiabetic activities. These bioactivities can be attributed to alkaloids, coumarins, limonoids, and phenylpropanoids present in different parts of the plant. Although some studies have indicated moderate toxicity of the extracts, some of the dominant compounds in this species, such as estragole and carbazole alkaloids, are mutagenic or cytotoxic. A clinical trial utilizing a Ghanaian herbal formulation containing *Clausena anisata* (Willd.) Hook.f. ex Benth. was found to be effective in reducing pain associated with osteoarthritis. Research progress to date supports the traditional use of this species in herbal medicine. However, these studies do not explain the relationships between traditional uses, pharmacological activities, and mechanisms of action. Thus, further studies should be designed to understand the biochemical properties and physiological effects of *Clausena anisata* (Willd.) Hook.f. ex Benth. extracts, facilitating the development and utilization of this medicinal resource.

**Keywords:** carbazole alkaloid; coumarin; limonoid; phenylpropanoid; essential oil



**Highlights**

This report presents a holistic overview of the ethnobotany, phytochemistry, and ethnopharmacology of *Clausena anisata* (Willd.) Hook.f. ex Benth., the only Rutaceous member of the *Clausena* genus extensively used in African traditional medicine.

**Medical history of objective**

*Clausena anisata* is a plant grown for its medicinal use in Asia. This plant is also widely used in traditional medicine in Africa; its leaves are usually placed under the bed or carried along during war by the Maasai because of its odoriferous properties. Plant twigs are used to steam patients to chase/dispel or warn off vengeful spirits.

**Background**

*Clausena anisata* (Willd.) Hook.f. ex Benth. (*C. anisata*) is a member of the Rutaceae (*Citrus*) family [1]. The *Clausena* genus was named after the Norwegian clergyman, Peder Clausson Friis; the genus comprises at least twenty-five known botanical species in Southern Asia, Australia, Africa, and the Pacific islands; these species were described in 1768 by the Dutch botanist Nicolaas Laurens Burman [2–4]. The major botanical synonyms of *C. anisata* total 43, including *C. dunniana* and *C. willdenovii* Wt. & Arn., *C. abyssinica* (Engl.) Engl., *C. inaequalis* (D.C.) Benth, *C. dentata* (Willd.) M. Roem., and *Amyris anisata* Willd. [5–7].

*C. anisata* is a deciduous shrub, small tree, or small tree with usually imparipinnate but sometimes paripinnate compound leaves densely dotted with glands and have a strong aniseed-like scent when crushed (Figure 1) [8] ([https://commons.wikimedia.org/wiki/File:Clausena\\_anisata\\_1.jpg](https://commons.wikimedia.org/wiki/File:Clausena_anisata_1.jpg), This file is licensed under the Creative Commons Attribution-Share Alike 4.0 International license). Leaves are usually clustered at twig ends. The flowers are small and white, with orange-yellow stamens. The inflorescence forms a branched axillary spray [9]. The fruit is a berry, approximately 1.3 cm wide, while the seeds are oblong in shape [7]. The plant is popularly known as “horsewood” or “maggot killer”, chiefly for its fetid leaves, which, when bruised, give rise to “Perdepis” (horse urine in the Afrikaans dialect) [10, 11].

*C. anisata* is the only member of *Clausena* genus in tropical Africa [12, 13]. Its country-specific geographical range includes forests, riverine thickets, and bushvelds from Guinea to Sierra Leone eastwards to Ethiopia, Sudan, and southward to the Cape in South Africa, only avoiding the driest regions [14–19]. The species has also been recorded in tropical and Southeast Asia, growing in India, Nepal, and Sri Lanka, extending as far as Queensland in northeastern Australia and some Pacific islands (Figure 2) [3, 11, 17, 20–27]. It is grown for its medicinal flavored oil in Malaysia, Indonesia, Vietnam, Thailand, and the Philippines (in the Philippine local brandy Anisdos) [1, 3, 7, 20, 28]. In Kenya, plants are usually placed under the bed, used for sleeping, or carried in wars among the Maasai because of its odoriferous properties [29]. Some of the local names of *C. anisata* used across the indigenous communities are summarized in Table 1 [30–85]. Its native ranges are tropical (South Africa, India, and South Central China) [21].

As a revered medicinal plant, *C. anisata* has been widely researched on. Two review articles have been published on this species, focusing on its bioactivity and phytochemistry [14, 86]. Given the exponential development of research approaches, analytical instruments, and strides in research concepts, many new features of the species have been determined recently. Nevertheless, there has been no up-to-date review of all the relevant research aspects of *C. anisata*. Therefore, this review is a holistic synopsis of the ethnomedicinal uses, phytochemistry, and ethnopharmacology of *C. anisata*, highlighting recent developments regarding the species and directions for future research.

**Ethnomedicinal uses of *C. anisata***

*C. anisata* whole plants, stems, roots, leaves, flowers, twigs, stems, and root bark, singly or in combination, are utilized in African folk medicine for various human and animal ailments (Table 2) [87–123]. Targeted ailments include malaria, fever, colds, bacterial and fungal infections, venereal diseases (gonorrhoea and syphilis), cough (tuberculosis), malignancies (breast and prostate cancer), and metabolic disorders, such as diabetes mellitus and coronary and renal diseases. The ethnoveterinary uses of this species include the management of maggots (myiasis) in cattle, back leg (Abakorpa), anthrax (Abasenga), helminthiasis, and other endoparasites. Multiple therapeutic remedies involving *C. anisata* have also been reported. For example, decoctions of *C. anisata* leaves mixed with *Lippia javanica* and *Eucalyptus grandis* leaves are used for respiratory ailments in South Africa [35]. In Ethiopia, the leaves of *C. anisata*, *Solanecio gigas*, and *Justicia schimperiana* are pounded together and applied dermally to treat skin irritation [64]. In another area in Ethiopia, inhaling *C. anisata* root powder with the roots of *Croton macrostachyus* and *Capparis tomentosa* is used as a treatment for devil attacks [65]. In South West Nigeria, a mixture of *C. anisata*, *Afraegle paniculata*, and *Azadirachtha indica* is administered to treat gut disturbance, and a concoction of the same called “Agbo” is used among the Yoruba as an antimalarial medicine [75]. In Nigeria, the Yoruba use a decoction of *C. anisata*, *Ageratum conyzoides*, *Momordica charantia*, and *Uvaria chamae* leaves (one glass cup thrice daily until the symptoms subside) for cancer management [77]. A similar decoction involving the roots of *C. anisata*, *Garcinia kola*, *Astonia boonei*, *Culcasia scandens*, and *Alafia barteri* is used to treat breast pain in Nigeria [19]. In Rwanda, *C. anisata* leaves, aerial parts of *Polydora serratuloides*, and leaves of *Kalanchoe* species are used to prepare decoctions for asthma treatment [83]. In Tanzania, a decoction of the roots of this species, along with those of *Byrsocarpus boiunianus*, *Albizia anthlmitica*, and *Carpolobia goetzei*, are used to treat constipation in children [124].

The leaves (branches/aerial parts) and stem bark are the most used parts for preparing herbal remedies. Phytochemical, pharmacological, and toxicity studies have primarily investigated these aspects. This is



Figure 1 *Clausena anisata* tree. Source: Wikimedia Commons.




Figure 2 Geographical distribution of *C. anisata* (in blue) based on Table 1 and other reports. *C. anisata*, *Clausena anisata* (Willd.) Hook.f. ex Benth.

**Table 1 Local names of *C. anisata* used across indigenous communities**

Folk name (local language)	Country	References
Umsanga (isiZulu), Iperepesi, Umtuto (Xhosa), Perdepis (Afrikaans), Isifudu (Zulu), Isifuthu, Umnukambhiba, Umnukelambiba, Umsanka, Umwashampunzi	South Africa	[30–38]
Mtombombare, Mnyapala, Munyapala (Digo), Kumunyabubi, Kisimbari, Shihunya bukundu (Luhya), Mukibia, Chesamishiet, Chebunoiwo (Marakweta), Matathi, Mutathi (Kikuyu), Mjavikali (Swahili), Chepkolol (Tugen), Siska (Boran), Kathimi kapala (Giriam), Kithiw'a, Muthungwa (Kamba), Siunya (Luo), Olmatasia (Maasai), Mukithia (Meru), Arawithargi (Sanya), Kalagapala-uvumba (Duruma), Olmatasia	Kenya	[39–53]
Nawawayo (NgaKaramojong), Munawaidudu/Mufunya idudu (Lusoga), Omutanu	Uganda	[54–56]
Ayida (Asante-Twi), Saman ndobir, Samandua	Ghana	[9, 57]
Gbossouazohouin	Benin	[58, 59]
Olmaa'ii, Ulumaa'i, Hulimay (Afan oromo), Ulumaa/Limich, Limche (Afan Oromo/Amaric), Etunisha, Zumi	Ethiopia	[60–71]
Mkoma vikali (Kwere, Nyangalio), Mjavikali, Msongambwa (Luguru), Mtanwa (Zinza), Mjafari	Tanzania	[10, 44, 72, 73]
Mbiet ekpene (Ibibio), Atabari Obuko, Agbasa (Yoruba)	Nigeria	[6, 19, 74–79]
Eyra	Togo	[80, 81]
Muvengahonye (Shona)	Zimbabwe	[82]
Umuno (Kinyarwanda)	Rwanda	[83]
Kattukariveppila/Kariveppila, Suganthaveppu, Potti/Pothi (Tamil, Malayalam), Ana	India	[7, 84]
Nampi (Tagalog)	Philippines	[85]

*C. anisata*, *Clausena anisata* (Willd.) Hook.f. ex Benth.

**Table 2 Ethnobotanical uses of *C. anisata* reported by ethnobotanical studies**

No.	Ailment (s) treated/ethnoveterinary uses	Part (s) used	Method of preparation and administration	Country/region	Authors
1	Diabetes, analgesic	R, L, St	Cooked in food as a culinary recipe	South Africa, Tanzania	[10, 36, 87]
2	Malaria, fever, colds, coughs, stroke, <i>Portabarak</i>	L, R, Bk, RB, St	Decoction taken, roots chewed for colds & coughs, leaf infusion taken, or steam inhaled.	Kenya, Nigeria, South Africa, Togo, Benin, Ethiopia	[15, 36, 39–42, 44, 50, 58, 81, 88–93]
3	Heart, liver and kidney diseases, jaundice, constipation	R, L, B, St	Decoction taken	Kenya, Tanzania, Togo	[40, 48, 80, 81, 85, 94]
4	Convulsions, gonorrhoea, epilepsy, syphilis, fungal infections	L, SB, R	Decoction taken. For epilepsy, the steam of leaves or roots are inhaled.	Tanzania, Uganda, Kenya, Nigeria, Benin	[44, 54, 59, 72, 73, 95–101]
5	Measles, toothache (toothbrush), ear infections, fever, influenza, eye problems, evacuant, headache, rheumatism, respiratory ailments (ulceration of the lung, tuberculosis, bronchial problems, asthma), sore throat	SB, RZ, L, R, St, Tw	Decoction taken, used as a chewing stick or chewed, placed and held on the aching tooth. For ear problems, juice of leaves is used as ear drop. Twigs used as a tooth brush. Leaf decoction taken for asthma.	Nigeria, Uganda, South Africa, Kenya, Benin, West Africa, Ethiopia, Rwanda	[22, 34, 35, 44, 45, 49, 62, 63, 74, 78, 83, 102–111]
6	Hypotension, hypertension, as an emetic, blood cleanser, ascariasis	B, L, St	Decoction of bark or leaves and stems taken.	Kenya, Ethiopia, South Africa	[45, 68, 91, 112]

Table 2 Ethnobotanical uses of *C. anisata* reported by ethnobotanical studies (continued)

No.	Ailment (s) treated/ethnoveterinary uses	Part (s) used	Method of preparation and administration	Country/region	Authors
7	Snakebites, colic pain, worms (children), sweating, analgesic (breast pains), as a tonic ( <i>umuthi omhlophe</i> )	L, R	Boil with rock salt and drink 250 mL or smear powder on bite.	Uganda, South Nigeria	Kenya, Africa, [19, 36, 44, 54, 113, 114]
8	Breast and prostate cancer, cancer (unspecified), skin irritation, evil eyes, spirits, mental illness/madness	L, R, SB, Tw, Fr	Leaf powder in honey eaten. For irritation, leaves mixed with other leaves applied after pounding. Root powder inhaled for evil eyes. For spirits, patient is steamed using twigs steam. Fruit or leaves chewed for madness.	Ethiopia, Kenya, Uganda, Nigeria	[47, 56, 61, 64-67, 69, 77]
9	Menstrual disorders, ease child birth, after-birth problems (post-partum haemorrhage, delayed/protracted labour), and increase lactation	L, R	Pounded roots used in soup; woman drinks it before bedtime or roots are chewed.	Nigeria, Kenya	[19, 44, 115, 116]
10	Calf diarrhoea ( <i>Aremor</i> ), worm infection in cattle, repellent	R	Infusion given	Uganda, Kenya, Cameroon	[55, 117, 118]
11	Maggots (myiasis), back leg ( <i>Abakorpa</i> ), anthrax ( <i>Abasenga</i> ) and helminthiasis	L	Crushed & packed onto wounds to expel maggots, decoction administered nasally for back leg/orally for anthrax/helminthiasis.	Zimbabwe, South Africa, Ethiopia, Kenya	[70, 119-121]
12	Mosquito, house fly, bedbug and cockroach repellent	L, St	By hanging or smoking	South Africa, Ethiopia	[30, 122]
13	Insecticide against stored-grain pests	SB, L, F, WP	Decoction taken orally, powdered and licked; sliced bark chewed; cold infusion taken orally.	Benin, Cameroon, Kenya	[115, 123]

F, flowers; Fr, fruit; L, leaves; R, roots; RB, root bark; St, stem; SB, stem bark; Tw, twigs; WP, whole plant; RZ, rhizome; *C. anisata*, *Clausena anisata* (Willd.) Hook.f. ex Benth.

explained by the high yield associated with them because of their high potential to accumulate, concentrate, and store therapeutic phytochemicals responsible for the treatment of various ailments [125]. The main methods of herbal preparation and administration are decoctions, cold infusions, pounding dried samples into powder and then licking, preparing poultices that are applied topically, squeezing fresh samples, and mixing with bathing water or direct chewing of the different parts (Table 2). In some instances, as is the case in the Xhosa of South Africa, the plant organ is cooked in food as a condiment [36]. Herbal remedies are rarely prepared by steam inhalation. For example, *C. anisata* twigs are used to steam patients to chase away revengeful spirits [17, 56].

#### Phytochemical profile of *C. anisata*

Phytochemical screening of extracts of *C. anisata* leaves, stem bark, root bark, stem, and roots indicated the presence of therapeutic secondary metabolites, including flavonoids, saponins, coumarins, phenols, alkaloids, phytosterols, tannins, phenols, triterpenes, steroids, cardiac glycosides, proanthocyanins, terpenoids, and saponosides [6, 9, 31, 60, 75, 82, 126-132].

Some studies have quantified the secondary metabolites in *C. anisata*. A multiflora quantitative study in Ghana indicated that methanolic leaf extracts of this species had  $0.87 \pm 0.43$  g/L of gallic acid,  $0.41 \pm 0.20$  g/L of vanillic acid,  $0.99 \pm 0.47$  g/L of syringic acid,  $0.74 \pm 0$  g/L of 2, 5-dihydroxy benzoic acid,  $0.61 \pm 0.30$  g/L of caffeic acid,  $0.55 \pm 0.27$  g/L of rosmarinic acid, and  $0.60 \pm 0.34$  g/L of *p-coumaric* acid as the major phenolic compounds [133]. Working with *C. anisata* acetonic leaf extracts, the total phenolic content (TPC) and total flavonoid content (TFC) were reported to be  $109.63 \pm 7.62$

mg gallic acid equivalent (GAE)/g and  $159.01 \pm 1.88$  mg quercetin equivalent (QE)/g, respectively [134]. In Zimbabwe, a TPC of  $0.11 \pm 0.004$  mg tannic acid equivalent per 100 mg of leaves was reported [82]. Another team reported that the TPC, TFC, and proanthocyanidin content of *C. anisata* leaves were  $29.50 \pm 1.26$  mg GAE/g,  $0.28 \pm 0.03$  mg CTE/g, and  $1.96 \pm 0.23$  mg CTE/g, respectively [135].

A report from South Africa indicated TPC values of  $31.30 \pm 0.05$  and  $28.10 \pm 0.99$  mg GAE/g, TFC values of  $11.70 \pm 0.17$  and  $7.60 \pm 0.20$  mg CE/g, and total iridoids values of  $3,019.60 \pm 63.35$  and  $3,264.70 \pm 96.40$  µg harpagoside equivalents per gram for stored and fresh leaves and twigs of *C. anisata* extracted with 50% methanol, respectively [136]. In the same study, the corresponding values of condensed tannins were  $1.39 \pm 0.03$  and  $1.33 \pm 0.02\%$  in dry matter, respectively. The extracts had no free gallic acid, while only the stored samples had gallotannins quantified to be  $68.45 \pm 5.14$  GAE/g [136]. Kadiri et al. indicated that *C. anisata* roots had  $0.32 \pm 0.03$ ,  $0.72 \pm 0.03$ ,  $3.79 \pm 0.26$ ,  $1.86 \pm 0.03$ , and  $0.05 \pm 0.01$  mg/g of tannins, saponins, alkaloids, flavonoids, and phenols, respectively [19]. Another investigation recorded  $6.69 \pm 0.54$  mg GAE/g and  $7.45 \pm 0.64$  mg GAE/g as the TPC values of ethanolic extracts of *C. anisata* leaves and stems, respectively; the corresponding TFC values were  $2.46 \pm 0.08$  mg QE/g and  $3.93 \pm 0.122$  mg QE/g [128]. The polyphenol content of hydroalcoholic leaf and stem extracts of *C. anisata* was 62.8% and 36.8%, respectively, according to Agbor et al. [126].

To date, 90 compounds have been isolated and characterized from extracts of *C. anisata*. Alkaloids, coumarins (sometimes substituted), limonoids, terpenoids, and phenylpropanoids are the major groups of compounds that have been characterized using chromatographic and spectroscopic techniques such as gas chromatography-mass



spectrometry, ultraviolet-visible, fourier-transform infrared, mass spectrometry, nuclear magnetic resonance spectroscopy, and complementary techniques such as heteronuclear multiple bond correlation spectroscopy and distortionless enhancement by polarization transfer spectroscopy.

### Alkaloids

The characteristic class of phytochemicals in rutaceous plants (of which *C. anisata* is a member) is carbazole alkaloids with complex structural diversity. They frequently occur in tandem with coumarins and phenylpropanoids (clausamines) and are classified as carbazole, lactone carbazole, and pyranocarbazole alkaloids. These alkaloids are responsible for the poor odor and nasal membrane irritation of this species [137]. In total, 35 alkaloids (1–35) have been reported in *C. anisata* (Table 3) [138–147]. Three novel and naturally isolated lactonic carbazole alkaloids (clausamines: A, B, and C) were identified in acetonic extracts of *C. anisata* branches.

### Coumarins

Coumarins are a characteristic class of heterocyclic compounds in

which the benzene ring is fused to a pyrone ring. In *C. anisata*, furanocoumarins, geranyl coumarins, and furanocoumarin lactones have been isolated (Table 4) [148–154]. A total of 38 coumarins (36–73) have been identified in this species.

### Tetranortriterpenes (limonoids)

Limonoids are chemically characterized by variations in the furanolactone core structure, a prototypical structure made of four six-membered rings with an adjoining furan ring. Six tetranortriterpenoids (74–79) were isolated from *C. anisata* stem bark and roots; these included limonin (74), zapoterin (75), clausenolide (76), clausenolide-1-ethyl ether (77), clausenarin (11 $\beta$ -hydroxydeacetylnomilin) (78), and 1-*O*-methylclausenolide (79) [142, 155].

### Other compounds

Other compounds not categorized into the preceding chemical groups have been isolated and characterized from *C. anisata* (Table 5) [156, 157]. Some of these compounds include chlorophyll derivatives (80, 81), peptide derivatives (82–84), and sterols (85, 86).

**Table 3 Alkaloids isolated from *C. anisata***

Compound	Part (s) used	References
Clausamine A (1)	Branches	[138, 139]
Clausamine B (2)	Branches, stem	[138, 140]
Clausamine C (3)	Branches, stem	[138–140]
Ekeberginine (4)	Branches, roots, stem bark	[138, 139, 141–143]
Methyl carbazole-3-carboxylate (5)	Branches	[138, 139]
Clausine E (6)	Stem, branches	[139]
Clausine F (7)	Stem, branches	[139, 140]
<i>O</i> -demethylmurrayanine (8)	Branches, stem bark, roots	[138, 139, 143]
Clausamine D (9)	Stem, branches	[139, 140]
Clausamine E (10)		
Clausamine F (11)	Branches	[139]
Clausamine G (12)		
Clausenol (13)		
Clausenine (14)	Stem bark	[144]
1-Methyl-3, 4-dimethoxy-2-quinolone (15)		[143]
3-Formyl-1-hydroxycarbazole (16)	Stem bark, roots	[143, 145]
Heptaphylline (17)		[141–143]
Girinimbine (18)	Roots, stem bark	[140, 143]
3-Methylcarbazole (19)	Stem bark, roots	[140, 143]
Mukonal (20)		
Glycosinine (21)		
Mukonidine (22)	Stem	[140]
Furanoclausamine A (23)		
Furanoclausamine B (24)		
Clausamine H (25)	Stem bark	[142]
Murrayamine-A (26)	Roots, stem bark	[141, 142]
Oxypeucedanin (27)	Not specified	[146]
<i>N</i> -methylswietenidine-B (28)	Stem bark, roots	[143]
Clausenamide (29)		
Mupamine (30)	Stem bark, roots	[143, 147]
Clausanitin (31)	Stem bark, roots	[143]
Atanisatin (32)		
Clausine B (33)	Leaves	[127]
Clausenocoumarin (34)	Leaves	[60]
Heptaphylline (35)	Roots	[60]

*C. anisata*, *Clausena anisata* (Willd.) Hook.f. ex Benth.

Table 4 Coumarins isolated from *C. anisata*

Compound	Part (s) used	References
Chalepin (36)	Roots, root bark, leaves	[60, 142, 146, 148, 149]
Osthol (37)	Root bark, roots	[25, 145, 148]
3-(1, 1-dimethyl allyl) xanthyletin (38)	Stem bark, root bark	[142, 148]
Xanthoxyletin (39)	Root bark, stem bark, whole plant	[142, 148, 150]
Imperatorin (40)	Aerial parts, roots, stem bark, leaves, whole plant	[60, 142, 145, 150–152]
Xanthotoxol (41)	Aerial parts, roots	[145, 151]
Lansamide-I (42)		
Indicolactone (43)		
Anisolactone (44)	Aerial parts	[151]
2',3'-Epoxyanisolactone (45)		
Heliettin (46)	Stem bark, roots, whole plant	[150, 153]
Gravelliferone methyl ether (47)	Stem bark, whole plant	[142, 150]
Swietenocoumarin I (48)	Stem bark, roots, whole plant	[153]
Triphasiol (49)		
Capnolactone (50)	Leaves, whole plant	[150, 152]
Anisocoumarin A (51)		
Anisocoumarin B (52)		
Anisocoumarin C (53)	Roots, stem bark, leaves, whole plant	[145, 150, 152, 153]
Anisocoumarin D (54)		
Anisocoumarin E (55)		
Anisocoumarin F (56)		
Anisocoumarin G (57)	Leaves, whole plant	[150, 152]
Anisocoumarin H (58)		
Anisocoumarin I (59)		
Umbelliferone (60)	Leaves	
Isoponcimar (61)	Leaves	[152]
Anisocoumarin J (62)	Leaves	
Unidentified compound with chromene skeleton (63)	Roots	[60]
Gravelliferone (64)	Stem bark, whole plant	[142]
5,7-Dimethoxy-8-(3-O-methylbut-2-O-enyl) coumarin (65)	Stem bark	[142, 150]
Excavatin D (66)		
7-Methoxy-6 (2'-oxo-3'-methyl butyl) coumarin (67)		
(R)-(+)-6-(20-hydroxy-30-methyl-30-butenyl)-7-methoxycoumarin (68)	Leaves	[142]
7-((E)-7-hydroxy-3,7-dimethylocta-2,5-dienyloxy)-coumarin (69)		
Phellopterin (70)	Stem bark	
Bergapten (71)	Leaves	
Isooxypeucedanin (72)	Leaves	
Seselin or 2',2'-dimethylpyranocoumarin (73)	Leaves	[154]

*C. anisata*, *Clausena anisata* (Willd.) Hook.f. ex Benth.

Table 5 Other compounds isolated from *C. anisata* extracts

Compound	Part (s) used	References
13 <sup>2</sup> (R)-hydroxyphytyton a (80)	Leaves	[142]
13 <sup>2</sup> (S)-phytyton a (81)		
N-benzoyl-L-phenyl alaninyl-N-benzoyl-L-phenyl alaninate (82)		
Aurantiamide acetate (83)	Roots, stem bark	[141, 143]
N-benzoyl-2-hydroxy-2-(4-methoxyphenyl) ethylamine (84)		
Sitosterol (85)		
Stigmasterol (86)		
Quercetin 3,4-dimethyl ether (87)		
(S/R)-3-(1,1,3-trimethyl-2-butenyl)-5-(2,3-dihydroxy-3-methylbutane)-7-methyl-2H-1-benzopyran-2-one (88)	Leaves	[127]
1,2-Benzenedicarboxylic acid, mono (2-ethylhexyl) ester (89)	Leaves	[156]
<i>Trans</i> -4-hydroxy-N-methylproline (90)	Leaves, stem	[157]

*C. anisata*, *Clausena anisata* (Willd.) Hook.f. ex Benth.

#### Compounds from essential oils (Eos) and volatile fractions of *C. anisata*

In addition to the isolated compounds, more than 150 phytoconstituents (91–340) have been identified in Eos and volatile fractions of *C. anisata* leaves, fruits (fruit pericarp), and seed in Supplementary Table S1. The dominant components of the Eos include (*E*)-anethole, limonene,  $\gamma$ -terpinene, sabinene, estragole or methyl chavicol, sabinene, germacrene D, and Z ( $\beta$ )-ocimene [158–162]. An overview of the composition of Eos from *C. anisata* growing in different parts of the world indicates that they have at least six chemovarieties: (i) estragole (methyl chavicol), (ii) ocimenone, (iii) (*E*)-anethole, (iv) (*E*)-foeniculin, (v) sabinene, and (vi)  $\beta$ -pinene [118]. These disparities in the constitution of Eos are explained by differences in climatic conditions, seasons, soil composition, and the intraspecific chemical polymorphism and extraction technique(s) used in Eos extraction [157, 163–166]. Overall, the Eos of *C. anisata* are rich in phenylpropanoids.

#### Pharmacological activities of *C. anisata* Eos, extracts, and isolated compounds

The pharmacological potential of Eos, crude extracts, and therapeutically active compounds (Figure 3) from *C. anisata* was evaluated. The reported biological activities include antioxidant, antimicrobial (antiviral, anti-human immunodeficiency virus-1, antibacterial, antifungal, and antidiarrheal), anti-inflammatory, anticancer (antitumor/antiproliferative), antidiabetic, antimycobacterial, immunomodulatory, analgesic, antipyretic, antiplasmodial, anticonvulsant, and anthelmintic activities. In addition, the antifeedant, repellent, and pesticidal activities of this species have been explored extensively.

#### Antioxidant/antiradical activity

The total in vitro antioxidant activity of Eos and extracts of *C. anisata* was assessed using multiple radical generating systems such as 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical scavenging, nitric oxide, ferric reducing power (FRAP), and 2,2'-azino-bis(3-ethylbenzthiazoline-6-sulfonic acid (ABTS) assays. These methods have been used in tandem in most studies to provide elaborate insights into the antioxidant potential of the species. In Zimbabwe, methanolic *C. anisata* leaf extracts afforded 80.90%  $\pm$  0.78% DPPH inhibition [82].

Another study reported that the Eos of *C. anisata* had a half-effective concentration (EC<sub>50</sub>) of 6.53 mg/L against 524  $\mu$ g/L for butylated hydroxytoluene (BHT) in a  $\beta$ -carotene-linoleate system [167]. The authors reported an EC<sub>50</sub> (10 days) of 1.77 mg/L against 156  $\mu$ g/L for BHT and 2.66 mg/L and 150  $\mu$ g/L for FRAP and DPPH assays, respectively [167]. The Eos of another collection of *C. anisata* had half inhibitory concentration (IC<sub>50</sub>) of 3.40 g/L found to be close and comparable to that of BHT (IC<sub>50</sub> = 42  $\mu$ g/L), thymol (IC<sub>50</sub> = 333

$\mu$ g/L), and carvacrol (IC<sub>50</sub> = 666  $\mu$ g/L) [26]. A multiflora screening study in Ghana recorded a radical scavenging activity of 14.50  $\pm$  0.06% for *C. anisata* methanolic leaf extracts in a DPPH assay [131].


The scavenging activities of the three solvent extracts of *C. anisata* were evaluated using DPPH, nitric oxide, FRAP, and ABTS assays in Supplementary Table S2 [31]. Acetonic extracts have been reported to exhibit strong antioxidant activities in vitro, which could explain the widespread medicinal use of *C. anisata* leaves and bark in the management of oxidative stress-induced diseases. Investigations of the antiradical activities of *E*-ocimenone rich-Eos of *C. anisata* leaves have indicated that it had lower radical scavenging activity (SC<sub>50</sub> = 5.10 g/L), which was less than that of the standard compound (BHT, SC<sub>50</sub> = 0.01 g/L) [168]. Investigating the effect of storage on the antioxidant potential of fresh and stored *C. anisata* leaves and twigs, Amoo et al. recorded EC<sub>50</sub> of 26.80  $\pm$  2.06 and 33.20  $\pm$  3.89  $\mu$ g/mL, respectively [136].

Similarly, Arsia, et al. reported maximum DPPH scavenging activities of 74.1%, 74%, 70.8%, 67.1%, and 55.1% for ethyl acetate/chloroform, acetonic, aqueous, hexane, and alcoholic leaf extracts of *C. anisata*, respectively [129]. Silver nanoparticles synthesized from ethanolic extracts of *C. anisata* leaf extracts had DPPH scavenging activity of 71.6% at 500  $\mu$ g/mL, comparable to 77.38% inhibition of the standard ascorbic acid at the same concentration [169]. The IC<sub>50</sub> value (200  $\mu$ g/mL) of the Silver nanoparticles was equal to that of ascorbic acid. Recently, Pavela et al., using DPPH and ABTS assays, reported that the antioxidant activity of *C. anisata* Eos as measured by IC<sub>50</sub> values were 1.81  $\pm$  0.01 mg/mL and 1.30  $\pm$  0.02 mg/mL [118]. The Trolox equivalent antioxidant concentration, as assessed using FRAP, was 151  $\pm$  10.20  $\mu$ mol TE/g. The high antioxidant potential obtained in the DPPH and ABTS assays was attributed to the abundance of anethole in the Eos, as previously reported by Senatore et al. [170]. The low reducing power in the FRAP assay was speculated to be elicited by the synergistic action of its other constituents, such as *p*-cymene, terpinene, and (*E*)-methyl isoeugenol [118]. Recently, the ethanolic extract of *C. anisata* leaves and stems was shown to possess strong antioxidant activity, with an IC<sub>50</sub> of 34.46  $\mu$ g/mL that the species requires a comprehensive evaluation of its antioxidant potential [157].

#### Antimicrobial activity

**Eos.** Employing the hole-plate diffusion method, volatile oils (1:1, 1:2, 1:5, and 1:10 dilutions) of *C. anisata* leaves exhibited dose-dependent inhibitory activity against *Aeromonas hydrophila* (zone of inhibition (ZOI) = 7.80–8.90 mm), *Alcaligenes faecalis* (ZOI = 10.20–12.70 mm), *Bacillus subtilis* (ZOI = 6.20–13.90 mm), *Beneckea natriegens* (ZOI = 14.30–19.70 mm), *Brochothrix thermosphacta* (ZOI = 5.40–6.90 mm), *Citrobacter freundii* (ZOI = 7.60–8.40 mm), *Clostridium sporogenes* (ZOI = 7.0–9.60 mm), *Enterobacter aerogenes* (ZOI = 7.90–9.70 mm), *Enterococcus faecalis* (ZOI = 12.20–15.80 mm), *Erwinia carotovora* (ZOI = 7.30–8.70 mm),





**Figure 3** Chemical structures of some compounds isolated from *C. anisata* that exhibited biological activities. *C. anisata*, *Clausena anisata* (Willd.) Hook.f. ex Benth.

Flavobacterium suaveolens (ZOI = 8.90–16.30 mm), *Lactobacillus plantarum* (8.30–8.90 mm), *Leuconostoc cremoris* (ZOI = 9.40–10.10 mm), *Klebsiella pneumoniae* (ZOI = 6.30–7.70 mm), *Micrococcus luteus* (ZOI = 7.0–7.70 mm), *Proteus vulgaris* (ZOI = 5.90–6.60 mm), *Pseudomonas aeruginosa* (ZOI = 5.70–6.50 mm), *Salmonella pullorum* (ZOI = 6.60–7.50 mm), *Staphylococcus aureus* (ZOI = 8.50–9.90 mm), *Serratia marcescens* (ZOI = 10.0–12.70 mm), and *Yersinia enterocolitica* (ZOI = 6.0–7.80 mm) [162]. The oils had no activity against *Escherichia coli*, *Acinetobacter calcoaceticus*, *Brevibacterium linens*, or *Moraxella* species. Fungal species *Geotrichum candidum* (85%–98%), *Aspergillus parasiticus* (44%–84%), *Candida albicans* (26%–72%), *Penicillium citrinum* (57%–71%), and *Alternaria alternata* (42%–69%) similarly showed significant sensitivity to the volatile oils, with *Aspergillus flavus* (28%–47%), *Aspergillus niger* (19%–56%), *Aspergillus ochraceus* (25%–57%), *Chaetomium globosum* (28%–48%), and *Fusarium culmorum* (15%–67%) inhibition at 1, 2, 5, and 10  $\mu\text{L}/\text{mL}$  of the Eos [162]. Yaouba et al. determined the antifungal activities of *C. anisata* leaf Eos against *Aspergillus flavus*, *A. niger*, *Aspergillus parasiticus*, and *Fusarium moniliforme* [168]. The growth of the fungi was completely inhibited when assessed at 4, 5, 5, and 5 mg/mL of the oils.

The Eos of *C. anisata* leaves were found to have Minimum inhibitory concentration (MIC) values of 62.50  $\mu\text{g}/\text{mL}$  and 125  $\mu\text{g}/\text{mL}$  each, respectively, against *Salmonella typhimurium* (*S. typhi*) and *Micrococcus luteus*, *Proteus mirabilis*, *Staphylococcus aureus* (*S. aureus*), and *Pseudomonas aeruginosa* (*P. aeruginosa*) [84]. *Bacillus subtilis*, *Escherichia coli* (*E. coli*), *Klebsiella pneumoniae* (*K. pneumoniae*), and *Proteus vulgaris* had MIC values of 250  $\mu\text{g}/\text{mL}$  each, while beta-hemolytic *Streptococcus pyogenes* had an MIC of 500  $\mu\text{g}/\text{mL}$ . A comparative assessment of the antimicrobial potential of the leaf Eos of three chemo-varieties of *C. anisata* was undertaken. Estragole-rich Eos of *C. anisata* had the highest antimicrobial activity against *Shigella* species and *Escherichia coli*, with a ZOI of 17 mm and 16 mm, respectively [171]. The oils produced lower bioactivities against *Salmonella typhi* (ZOI = 13 mm), *Staphylococcus aureus*, and *Proteus species* (ZOI = 11 mm in each case). In contrast, *Pseudomonas aeruginosa* and all fungi had a ZOI less than or equal to 9 mm. Another chemo-variety (trans-anethole-rich) of oil showed the highest antimicrobial activity against *Shigella* species (ZOI = 12 mm), *Salmonella typhimurium* and *Escherichia coli* (ZOI = 11 mm in each case). The other microorganisms evaluated had a ZOI of less than or equal to 10 mm. The third chemo-type (fenculin-rich) Eos were inactive against all tested pathogenic microorganisms with ZOI less than 10 mm [171]. A similar small ZOI ( $6 \pm 0$  mm at 10  $\mu\text{L}$ ) was later recorded for *C. anisata* leaf Eos against *Escherichia coli* by Sessou et al. [172]. The Eos constituent's linalool,  $\alpha$ -terpinene,  $\gamma$ -terpinene, cis-sabinene hydrate, and sabinene from *C. anisata* have been previously shown to be responsible for the antibacterial and antifungal activities of Eos in this species [162].

**Extracts.** Ndukwe et al. reported that extracts of *C. anisata* twigs and stems (used as chewing sticks in Nigeria) were inactive against *Staphylococcus aureus*, *Bacillus subtilis*, *Escherichia coli*, and *Pseudomonas aeruginosa* [78]. A similar observation was made by Tafadzwa in which methanolic extracts of *C. anisata* tuber extracts had a ZOI of  $1.25 \pm 0.50$  mm against *Pseudomonas aeruginosa* while *Staphylococcus aureus*, *Streptococcus* Group A, and *Escherichia coli* were not inhibited [82]. This corroborated the findings of Kemoli et al., who found that aqueous extracts of *C. anisata* stem and bark had no antibacterial activity against oral pathogens, such as *Streptococcus mutans*, *Streptococcus sobrinus* (clinical strains), *Porphyromonas gingivalis*, *Prevotella melaninogenica*, and *Bacteroides oralis* [53]. While verifying the same traditional claim in the Ethiopian context, *C. anisata* dichloromethane/methanol and aqueous extracts were reported to possess only moderate bioactivity against *Bacillus cereus*, *Staphylococcus aureus*, *Enterococcus faecalis*, *Streptococcus mutans*, *Klebsiella pneumoniae*, *Escherichia coli*, *Candida albicans*, *Candidaneoformans*, and *Lactobacillus acidophilus* with MIC values between 3 to 8 mg/mL [71]. A bioautographic investigation of selected Southern African plant species used to traditionally manage helminth infections indicated that *C. anisata* exhibited excellent

antifungal activities with MIC of 0.02 mg/mL and therapeutic index of 2.65 against *Aspergillus fumigatus* [173]. It also had the lowest cytotoxicity, with a median lethal concentration ( $\text{LC}_{50}$ ) of 0.17 mg/mL.

The antimicrobial activity of ethanol leaf extract of *C. anisata* was assessed, using agar well diffusion and micro-dilution methods, against clinical isolates of *Candida albicans*, gram-positive bacteria (*Bacillus subtilis*, *Staphylococcus aureus*, *Enterococcus faecalis*, and *Bacillus thuringiensis*), and gram-negative bacteria *Pseudomonas aeruginosa* and *Proteus vulgaris* [9]. The extracts had MIC values in the range of 0.50 to 7 mg/mL against gram-positive bacteria, 2.50 to 1 mg/mL against gram-negative bacteria, and 5.50 mg/mL against *C. albicans*. Upon fractionation, the methanolic fraction gave the highest bioactivity, with MICs ranging from 0.60 to 5 mg/mL and 1 to 3 mg/mL for gram-positive and gram-negative bacteria, respectively. The chloroform fraction had higher MIC values spanning from 3 to 7.50 mg/mL and 2 to 6.50 mg/mL for gram-positive and gram-negative bacteria, respectively. The last fraction, obtained from petroleum ether, had MICs spanning from 4.50 to 8 mg/mL for gram-positive and gram-negative bacteria. However, Arsia et al. indicated that ethyl acetate/chloroform extract had higher bacteriostatic potential against *Corynebacterium* and *Proteus* species with a ZOI of 43 mm and 32 mm for *Staphylococcus epidermidis* [129].

Another bioassay investigating chloroform, ethyl acetate, and methanol extracts of *C. anisata* fruits, leaves, stem bark, twigs, and roots reported MIC values of 0.78 to 6.25 mg/mL (for *Pseudomonas aeruginosa*), 1.56 to 6.25 mg/mL (for *Salmonella typhimurium*, *Salmonella kisarawe*, and *Klebsiella oxytoca*), 3.13 to 6.25 mg/mL (for *Proteus mirabilis*), 3.13 to 12.50 mg/mL (for *Escherichia coli* and *Klebsiella pneumoniae*), and 3.13 to 50 mg/mL for *C. albicans* [174]. Qualitative antibacterial activity analysis (bioautography) was performed to assess the bioactivity of the constituents of the acetonetic leaf extracts of *C. anisata* against *Streptococcus pyogenes*, *Escherichia coli*, and *S. aureus*. Three fractions of the extract had remarkable activity against the tested pathogens (with ZOI ranging from  $6 \pm 0.50$  mm to  $9.10 \pm 0.10$  mm), comparable to that of standard drugs (amoxicillin and ciprofloxacin with ZOI =  $12.33 \pm 1.50$  mm and  $12.33 \pm 0.57$  mm against *Streptococcus pyogenes*) [127].

Mutuku indicated that methanol extracts of *C. anisata* (stem bark) had significant antibacterial potential against methicillin-resistant *Staphylococcus aureus* with MIC and minimum bactericidal concentration values of 0.06 mg/mL and 0.78 mg/mL, respectively [175]. Agbor et al. reported that hydroalcoholic extracts of *C. anisata* leaves and stems were active against some isolated periodontal microorganisms (*Streptococcus mutans*, *Lactobacillus* species, *Aggregatibacterium actinomycetemcomitans*, *Fusobacterium nucleatum*, and *Prevotella intermedia*), with MICs between 50 and 100 mg/mL [126]. The lowest MIC was 50 mg/mL for *Aggregatibacterium actinomycetemcomitans* and *Fusobacterium nucleatum* isolates for the leaf extract and *Aggregatibacterium actinomycetemcomitans* for the stem bark extract. A recent investigation by Machumi et al. indicated that *C. anisata* leaf extract and petroleum ether, dichloromethane, ethyl acetate, and alkaloidal fractions exhibited marked antifungal activity against *Cryptococcus neoformans* and *C. albicans* with MIC ranging from 0.39 to 6.25 mg/mL [164]. The study further indicated that the traditional claim of curative potential of this species varies with the harvest season, with the highest activity in the rainy season than in the dry season.

**Isolated compounds.** The carbazole alkaloids clausenol and clausenine, isolated from the ethanolic stem bark of *C. anisata* were evaluated for their antimicrobial activity [144]. Clausenol was reported to be active against gram-positive and gram-negative bacteria and fungi, with MIC values of 7, 14, 12, 14, 13, 5, and 2  $\mu\text{g}/\text{mL}$  against *Escherichia coli*, *B. subtilis*, *Salmonella typhimurium*, *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Candida albicans*, and *Trypophyton rubrum*, respectively. The corresponding MIC for clausenine were 40, > 100, > 100, 16, > 100, 16, and 5  $\mu\text{g}/\text{mL}$ , respectively. However, these MICs were higher than those of the commercial antibiotics (penicillin, streptomycin, griseofulvin, and

nystatin) used in this study [144]. These observations agree with a later work of Amoo et al., wherein fresh and stored leaves and twigs of *C. anisata* elicited antimicrobial activity with MIC and minimum microbicidal concentration values of 1.56–6.25 mg/mL and 6.25 mg/mL or greater against *Candida albicans*, *Pseudomonas aeruginosa*, and *Staphylococcus aureus* [176].

Tatsimo et al. assessed the antibacterial activities of crude methanolic and hexane extracts of *C. anisata* and 21 isolated compounds; they reported that residues of crude leaf and stem bark extracts had broad-spectrum antibacterial activity against *Vibrio cholerae* (*V. cholerae* SG24, *V. cholerae* CO6, *V. cholerae* NB2, *V. cholerae* PC2, *Shigella flexneri* 2a, and *Staphylococcus aureus* ATCC 25,923 [142]. Among the isolated compounds, imperatorin had the highest activity, with an MIC of 34.66 µg/mL. This was followed by phellopterin (MIC = 61.33 µg/mL), murrayamine-A (MIC = 64 µg/mL), 7-methoxy-6 (2'-oxo-3'-methyl butyl) coumarin (MIC = 69.33 µg/mL), chalepin (MIC = 69.33 µg/mL), 7-((E)-7-hydroxy-3,7-dimethylocta-2,5-dienyloxy)-coumarin (MIC = 82.66 µg/mL), 13<sup>2</sup> (R)-hydroxypheophyton *a* (MIC = 128 µg/mL), ekeberginine (MIC = 144 µg/mL), girinimbine (MIC = 144 µg/mL), isooxypeucedanin (MIC = 149.33 µg/mL), xanthoxyletin (MIC = 160 µg/mL), 1-O-methylclausenolide (MIC = 160 µg/mL), clausamine H (MIC = 235.33 µg/mL), bergapten (MIC = 256 µg/mL), gravelliferone (MIC = 277.33 µg/mL), and 3-(1,1-dimethyl allyl) xanthyletin (MIC = 352 µg/mL) [142]. Time-kill kinetic studies of imperatorin against *V. cholerae* CO6 and *V. cholerae* SG24 (as a function of incubation time) indicated that it caused an appreciable reduction (approximately 6-log growth reduction) of bacteria, comparable to that of ampicillin when their minimum bactericidal concentrations were used within 6–10 h [142].

Recently, heptazoline, chalepin, imperatorin, and an incompletely identified compound with a chromene skeleton (63) isolated from the dichloromethane (DCM)/methanol extract of *C. anisata* roots were screened using the agar diffusion method against strains of *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, and *B. subtilis* [60]. Heptazoline and imperatorin had comparable antibacterial activity against *Staphylococcus aureus* and *B. subtilis* (with ZOI of 14 mm and 13 mm, respectively) to that of ciprofloxacin (ZOI = 15 mm) at 20 µg/mL. Interestingly, chalepin has a comparatively higher bacteriostatic potential against *B. subtilis* (ZOI = 16 mm zone of inhibition) than ciprofloxacin (ZOI = 15 mm) [60].

#### Anti-inflammatory activity

Crude ethanolic leaf extracts of *C. anisata* (39–117 mg/kg) induced a significant dose-dependent reduction in induced inflammation and fever. These activities were speculated to have been caused by various compounds reported for this species [6]. Another assay assessed stored and fresh methanolic extracts of *C. anisata* leaves and twigs for their cyclooxygenase (COX) (COX-1 and COX-2) inhibition potency [176]. The extracts caused 26.7% to 65.3% inhibition of the COX enzymes. Interestingly, the inhibition of COX-1 by stored *C. anisata* differed significantly from that of the fresh parts. The anti-inflammatory potential of the extracts on prostaglandin synthesis during the inflammation cascade through inhibition of COX enzymes (especially COX-2) was asserted to be through this pathway, as is the case with nonsteroidal anti-inflammatory allopathic drugs [176].

Another report found that crude acetonic extracts of *C. anisata* leaves exhibited the best inhibitory activity against nitric oxide production (96.9% inhibition) when tested at 6.25 µg/mL, comparable to that of quercetin (91.08% inhibition) [134]. Owing to the release of nitric oxide contributes to inflammation, the extract proved to be a potential scavenger or inhibitor of nitric oxide synthase activity and/or its expression. In vitro, 30 mg/kg of *C. anisata* extracts dwindled nitric oxide production and pro-inflammatory cytokines such as interleukin (IL)-6 and prostaglandin E<sub>2</sub> in lipopolysaccharide-stimulated RAW 264.7 cells [20]. The extract further diminished the expression of pro-inflammatory mediators such as COX-2. While in vivo, *C. anisata* extract significantly reduced inflammatory cell numbers in bronchoalveolar lavage fluid and

suppressed pro-inflammatory cytokine levels, including tumor necrosis factor- $\alpha$ , IL-6, and IL-1 $\beta$ , as well as reactive oxygen species production in bronchoalveolar lavage fluid. Taken together, these findings demonstrate that the extracts inhibited inflammatory responses in a mouse model of lipopolysaccharide-induced acute lung injury and stimulated RAW 264.7 cells. Thus, *C. anisata* extract is a potential candidate for development as an adjunctive treatment for inflammatory disorders such as acute lung injury [20].

In vivo anti-inflammatory and analgesic investigations of ethanolic extracts of *C. anisata* roots revealed that it had a significant anti-inflammatory effect at 1,000 mg/kg in a carrageenan-induced edema model [145]. In acetic acid-induced writhing and hot plate tests, it had a strong but dose-dependent analgesic effect, with maximum analgesic activity of 72.1% at 1,000 mg/kg [145]. The isolated compounds (osthol 37 and anisocoumarin B 52) displayed the highest anti-inflammatory activity at 9 mg/kg, which was better than that of indomethacin. Heptaphylline (17) was the most potent (48.7%), while others (except xanthotoxol 41) showed significant analgesic activity at 6 mg/kg (higher than diclofenac). The analgesic activity of 52 (50.4%) was the highest among the isolates tested and the standard, tramadol, in the hot plate test. The nonselective opioid receptor antagonist naloxone abolished the analgesic effect of the crude extract and the tested isolates (41 and 52) in the hot plate test, suggesting an effect via the central opioidergic system [145].

The potential of selected 21 species used against *Propionibacterium acnes* (*P. acnes*) in South African herbal medicine has been assessed [157]. Interestingly, the ethanolic extracts of *C. anisata* leaves and stems inhibited the growth of *P. acnes* with an MIC of 31.25 µg/mL, but the isolated compound (*trans*-4-hydroxy-1-methyl-L-proline 90) had an MIC value > 500 µg/mL. The anti-inflammatory activity of *C. anisata* was evaluated along with its probable mechanism of action using lipase and hyaluronidase [157]. Cells stimulated with *P. acnes* and treated with 50, 25, 12.50, and 6.25 µg/mL of *C. anisata* extracts had diminished Interleukin-8 production at concentrations of 322.48, 365.98, 383.62, and 409.52 pg/mL, respectively. The extracts at 500 µg/mL resulted in 21.93% inhibition of lipase activity, which could be translated into their demonstrated potential to ameliorate sebum production associated with *Acne vulgaris* progression. At the same concentration, the extract portrayed 49.02% hyaluronidase activity inhibition, interpreted as its potential to deter the proliferation of *P. acnes* cells to adjoining cells [157].

Another probe assay was performed to evaluate the inhibitory activities of acetonic and aqueous extracts of *C. anisata* on secretory phospholipase A<sub>2</sub> (sPLA<sub>2</sub>), lipoxygenase (15-LOX), and COX enzymes, which are South African plants used for skincare and beauty [177]. The extract had a high inhibitory activity against the COX-2 enzyme (IC<sub>50</sub> = 15.18 and 6.89 µg/mL, respectively). The IC<sub>50</sub> values for sPLA<sub>2</sub>, 15-LOX, and COX-1 were 24.82 and 24.61 µg/mL, 67.62 and 41.45 µg/mL, and 4.29 and 3.13 µg/mL, respectively. Usually, free arachidonic acid released by sPLA<sub>2</sub> activity is metabolized by 15-LOX to form hydroxyeicosatetraenoic acid, resulting in atherosclerotic plaque formation. Given the results that indicated low inhibition of sPLA<sub>2</sub> encountered, it was deduced that the extracts exerted their inhibitory activity via other mechanisms [177]. It is recommended that the extracts have the potential to serve as nonselective inhibitors of COX-1 and COX-2 enzymes, implying that incorporation of the extracts selective for COX-2 and 15-LOX into beauty product formulations could enhance skin beauty by reducing scarring, dark spots, and uneven skin usually encountered with chronic inflammation and progression [177].

#### Anticancer activity

Various extracts and compounds isolated from this species elicited marked bioactivity against cancer cells (Table 6) [178–181]. Although other malignancies such as breast and prostate cancer have been treated using *C. anisata* leaves (Table 2), no study has evaluated the potential of the plant extracts against cell lines of these cancers. The mechanisms by which the evaluated extracts (compounds) mediate their antitumor activity have been elucidated in other

**Table 6 Anticancer activities of extracts and isolated compounds from *C. anisata***

Extract/compounds	Cancer cell lines tested	Anticancer activity	References
Ethanol leaf extract	Human epidermal melanocyte cell line (PCS-200-013) and cellular tyrosinase	Nontoxic to melanocytes at 15 µg/mL and but inhibited cellular tyrosinase with IC <sub>50</sub> of 1.56 µg/mL	[178]
Hydroethanolic root extract	Jurkat (human acute-cell leukemia) cells	IC <sub>50</sub> = 293.14 µg/mL	[179]
Leaves and stem bark ethanol extracts	HL-60 monocyte lymphoma (U937) cells	Low cytotoxicity with IC <sub>50</sub> = 74.46 µg/mL	[157]
Leaves ethanol extract	Human rhabdomyosarcoma cancer cell line	CC <sub>50</sub> of 8.83 µg/mL	[79]
Methanolic and DCM aerial part extracts	HL-60 cells	IC <sub>50</sub> = 118.50 µg/mL and 225.40 µg/mL	[180]
Methanolic stem bark extract	Hoechst 33342, alamar blue, calcein-AM, glutathione depletion and oxygen-consumption assays	Minimal toxic dosage of < 7.81 to 250 µg/mL	[181]
Murrayamine-A, 3-(1, 1-dimethyl allyl) xanthyletin (38), gravelliferone (64), excavatin D (66), 7-((E)-7-hydroxy-3, 7-dimethylocta-2, 5-dienyloxy)-coumarin (69), phellopterin (70), 1-O-methylclausenolide (79)	HeLa cells	For 70, LC <sub>50</sub> was > 10 lg/mL. For the other compounds, LC <sub>50</sub> = 1.14 to 3.26 lg/mL and selective to normal Vero cells (LC <sub>50</sub> = 69.15–434.78 lg/mL)	[142]
Roots, leaves and bark extracts	Leukemia CCRF-CEM cells	Inhibited less than 50% of cancer cells at 40 mg/mL	[130]
Methanolic and hexanic leaves and stem bark extracts	HeLa cells	–	[142]
Clausamines B–E (2, 3, 9, 10), Clausine F (7), mukonal (20) and glycosinine (21)	HL-60	Clausamine E had the highest cytotoxicity (cell viability 47.3%); clausamines B and C had lower cytotoxicity (viability ~80%)	[140]
Clausamines A–E (1–3, 9, 10), ekeberginine (4), methyl carbazole-3-carboxylate (5), Clausine F (7) and O-demethylmurrayanine (8)	Epstein-barr virus early antigen activation induced by 12-O-tetradecanoylphorbol-13-acetate in Raji cells	All the compounds inhibited Epstein-Barr virus activation at 1 × 10 <sup>2</sup> mol ratio	[139]

*C. anisata*, *Clausena anisata* (Willd.) Hook.f. ex Benth.; DCM, dichloromethane; IC<sub>50</sub>, inhibitory concentration; CC<sub>50</sub>, 50% cytotoxic concentration; LC<sub>50</sub>, lethal concentration; HeLa, human cervical cancer; HL-60, human leukaemic; –, not mentioned.

*Clausena* species. For example, girinimbine (18) and other alkaloids isolated from *Clausena dunniana* and clausine E (from *Clausena vestita*) exhibit growth inhibitory activity by inducing apoptosis and cell cycle arrest in the S and G2/M phases [182, 183].

#### Antiparasitic (antiplasmodial/antimalarial and anthelmintic) activities

Extracts from *C. anisata* have been investigated using either nonradioactive antiplasmodial (in vitro) or four-day *Plasmodium berghei* ANKA suppressive (in vivo) assays. In an earlier study dichloromethane:methanol (1:1) extracts of *C. anisata* twigs and leaves were investigated using the parasite lactate dehydrogenase assay; the extracts had IC<sub>50</sub> values of 18, 55, and > 100 µg/mL against the *Plasmodium falciparum* strain D10, and were subsequently not labeled as highly active antiplasmodial plants [184]. Another study using petroleum ether, dichloromethane, and methanolic extracts of *C. anisata* roots, stem bark, and leaves used a parasite lactate dehydrogenase assay and reported that the extracts had IC<sub>50</sub> values between 5 and > 499 g/mL against the multidrug-resistant K1 strain of *Plasmodium falciparum* [185]. Similarly, chloroform, aqueous, and its fractions obtained from *C. anisata* leaves were assayed using Peter's 4-day test. Ethanol and aqueous extracts and their fractions showed dose-dependent chemosuppressive and schizonticidal effects on (*P. berghei*) infection in mice (4-day test) with parasitemia [131]. Another in vivo assessment of the suppressive, curative, prophylactic, and acute toxicity properties of hexane and chloroform extracts of *C. anisata* stem bark against murine malaria has been performed [51]. The authors revealed that chloroform extract (500 mg/kg/day) had

66.1% and 73.4% parasite (*P. berghei*) reductions in the prophylactic and suppressive tests, respectively. The same dose of the hexanic extract had prophylactic effects (56.7% and 30.7%, respectively). The mean survival times of 12.3 days and 9.3 days, respectively, were recorded for the extracts in the curative test [51].

The crude ethanol extract of *C. anisata* leaves exhibited anthelmintic activity, whereas the hexanic extract showed no bioactivity when tested at concentrations of 1–2 mg/mL in 2-hour and 7-day anthelmintic assays [186]. Similarly, organic (methanol and dichloromethane) extracts of *C. anisata* aerial parts displayed in vitro antitrypanosomal effects on the parasitic kinetoplastid (*Trypanosoma brucei brucei*) with IC<sub>50</sub> values of 92.20 µg/mL and 49.80 µg/mL [180]. Christensen et al. assessed the antiparasitic potential of the ethanol extract of *C. anisata* radix against the obligate anaerobic parasitic gut protist *Blastocystis* (subtype 4) and reported that the extract was among the best 24 plant extracts assayed, with an IC<sub>50</sub> of 314 mg/mL after 24-hour incubation [187]. However, the authors noted that the antimicrobial activity of the extracts at 800 mg/mL could have influenced the anti-*Blastocystis* activity observed [187]. In another screening involving both African and Caribbean plants, ethanol *C. anisata* root extract exhibited the highest anthelmintic activity against *Ascaris suum* with an EC<sub>50</sub> value of 74 lg/mL [188].

A Ghanaian study screened the antiprotozoal efficacy and potential mechanisms of action of 112 crude extracts of indigenous plants against *Trypanosoma*, *Leishmania*, and *Plasmodium* parasites [179]. Hydroethanolic root extracts of *C. anisata* was found to possess cytotoxic effects to *Trypanosoma brucei brucei* (IC<sub>50</sub> = 29.50 µg/mL), *L. donovani* (IC<sub>50</sub> = 12.10 µg/mL), and *Plasmodium falciparum* (IC<sub>50</sub> =



487.56 µg/mL). These results implied that the extracts elicited strong anti-Leishmania activity [179]. Another study in Cameroon reported that *C. anisata* leaf Eos had a very high IC<sub>50</sub> (> 100 µg/mL) against *Trypanosoma brucei* [189].

#### Antidiabetic activities

The hypoglycemic effect of *C. anisata* root methanol extract was investigated in normal (normoglycemic) and streptozotocin-treated (90 mg/kg) diabetic Wistar rats [87]. Graded doses of *C. anisata* extract were administered to fasted normal and fasted diabetic rats. Relatively moderate to high doses of the extracts (100–800 mg/kg) induced a dose-dependent reduction in blood glucose concentrations in both normal and diabetic rats. At 800 mg/kg, the extract reduced the mean basal blood glucose concentrations of normal and diabetic rats by 57.52% and 51.3%, respectively. The mechanism of this activity was not elucidated, but it was expected to be elicited by the terpenoids and coumarins that are abundant in this species. The study findings agreed with the folkloric utilization of *C. anisata* roots in the treatment of adult-onset type-2 diabetes mellitus in South Africa [87].

The inhibitory effects of *C. anisata* extracts against human urinary α-amylase, α-glucosidase, and glucose 6-phosphatase in vitro, as well as rat α-amylase and α-glucosidase in vivo, were investigated [190]. These enzymes are associated with carbohydrate metabolism, and this study was conducted to establish the mechanism of hypoglycemic action of the leaf extracts. Aqueous and methanolic extracts reportedly had significant inhibition of α-amylase (> 80%) and hepatic glucose-6-phosphatase (60% and 58%). However, they are less potent inhibitors of α-amylase than acarbose [190]. Acetonic and hexanic extracts similarly showed significant inhibition (> 80%) of *Bacillus stearothermophilus* α-glucosidase. Thus, in vivo confirmation of the observed in vitro evaluation of phosphatase inhibition by the extracts and their hypoglycemic action is not mediated through the inhibition of intestinal carbohydrate hydrolyzing enzymes [190, 191].

In another study, aqueous and methanolic extracts of *C. anisata* leaves inhibited human urinary α-amylase with IC<sub>50</sub> values of 1,947 and 2,436 µg/mL [33]. This inhibition has been suggested to be reversible and non-competitive. Acetonic and hexane extracts of *C. anisata* inhibited *Bacillus stearothermophilus* α-glucosidase, with IC<sub>50</sub> values of 1,020 and 2,068 µg/mL. The aqueous and methanolic extracts of *C. anisata* leaves also inhibited hepatic glucose 6-phosphatase with IC<sub>50</sub> values of 493.60 and 1,012 µg/mL, and the effect of the latter was far less than that of the reference inhibitor (sodium vanadate). Although both aqueous and hexane extracts of *C. anisata* leaves inhibited human urinary α-amylase and *B. stearothermophilus* α-glucosidase in vitro, the aqueous extract was more potent in vivo inhibition of rat hepatic glucose 6-phosphatase than the known inhibitor of the catalytic subunit of this multi-component enzyme system [33].

Arsia et al. found that the percentage increase in glucose uptake into yeast cells was correlated with the glucose concentrations used when *C. anisata* leaf extracts were tested [129]. The aqueous and alcoholic leaf extracts of *C. anisata* produced maximum increase in 10 mM glucose concentration, i.e., 95% and 95.3% at 2,000 µg/mL. The alcoholic, aqueous, and acetone extracts showed appreciable (> 20%) enzyme inhibitory activity against human urinary alpha amylase, followed by hexane (3.4%) and ethyl acetate + chloroform extracts (8.1%).

The hypoglycemic effect of AgNPs synthesized using ethanolic leaf extract of *C. anisata* was assessed using glucose uptake by yeast cells, alpha-amylase inhibition assay, adsorption capacity, and glucose diffusion assay [169]. The silver nanoparticles (SNPs) had alpha-amylase inhibitory activity of 80.32% at 500 µg/mL, with IC<sub>50</sub> of 100 µg/mL. Glucose uptake by yeast cells was found to be dose-dependent, with a maximum uptake found to be 68.29% at 10 mM, and the rate of glucose diffusion across the membrane ranged from 30 to 180 min [169]. The same team tested two doses (100 mg/kg body weight (BW) and 200 mg/kg BW) of crude leaf and root extracts and 5 mg/kg BW and 10 mg/kg BW of SNPs synthesized from the leaf and root extracts of *C. anisata* [192]. The extracts were

administered to alloxan-induced diabetic rats for 30 days, and the results indicated that the activities (triglycerides, antioxidants, cholesterol, serum glucose, liver biomarkers, and gluconeogenic enzymes) were reduced. The extracts proved beneficial for the regeneration of pancreatic β-cells in alloxan-induced diabetic rats. SNPs from the root extracts at 10 mg/kg BW were the most potent [192].

#### Anticonvulsant activity

A lead study by Makanju highlighted that aqueous extracts of *C. anisata* root bark depresses the central nervous system with mild-to-moderate anticonvulsant activity at 250–500 mg/kg in mice [96]. A later report using the Gamma aminobutyrric acid-benzodiazepine receptor assay reiterated that the ethanolic extract of *C. anisata* bark and aerial parts had weak and dose-dependent anticonvulsant activity [193]. Later, Mbah and Kenchukwu observed that ethanolic extracts of the leaves and stem bark of this species had no anti-convulsive effect but sedative properties, with intraperitoneal administration of bark extract affording only partial protection at 800 mg/kg [194]. In extension of the foregoing study, the antiepileptic potential of *C. anisata* leaf, root and stem bark ethanolic extracts (labeled CALE, CARE, and CASE, respectively) against induced seizures was explored [132]. Their study found that CARE (800 mg/kg) delayed the onset of convulsions, providing 33.33% protection. The remaining extracts did not provide any appreciable protection. The authors reiterated that CARE-housed therapeutic compounds (primarily tannins and saponins) have potential benefits for petit mal epilepsy, giving a pharmacological reason for its claimed use in epilepsy phytotherapy [132]. Further research is required to confirm the efficacy of the compounds isolated from this species.

#### Antihypertensive activity

To validate the folkloric claim of using *C. anisata* leaves by hypertensive Africans, aqueous and ethanolic extracts were screened for the angiotensin-converting enzyme (ACE) inhibitors. The in vitro results indicated that extracts had 54% and 1% ACE inhibition [91]. Based on this observation, another study was undertaken to investigate the blood pressure-lowering potential of aqueous leaf extracts of this species in a spontaneously hypertensive rat model. The authors investigated whether the antihypertensive effects were mediated by diuresis, inhibition of the renin-angiotensin-aldosterone blood pressure control system, and/or possible negative inotropic or chronotropic cardiac effects. Extracts administered intra-arterially at 400 mg/kg BW reduced aortic blood pressure via the reduction in plasma angiotensin II levels, reaffirming the postulation by Duncan et al. [37, 91]. This bioactivity was predicted to be elicited by coumarins in the extracts because they have been previously shown to possess antihypertensive properties [195].

#### Antinociceptive and anti-arthritis properties

Hydroethanolic extract of *C. anisata* leaves at 400 mg/kg reduced the duration of paw licking/biting during the early (42.12%) and late (75.79%) phases of formalin-induced nociception [75]. Findings from the study illustrated that *C. anisata* elicits antinociceptive activity through interaction with opioidergic, noradrenergic, L-arginine-nitric oxide, and serotonergic pathways, as well as anti-arthritis properties conferred by its potential to inhibit the release of inflammatory mediators and oxidative stress [75].

#### Antifeedant, repellent, fumigant, and contact toxicity (pesticidal activities)

*Eos. C. anisata* extracts and Eos were bioassayed against houseflies (*Musca domestica* L.) and cockroaches (*Periplaneta americana* L.) for various insect control activities [196]. The Eos had knockdown effects on *Musca domestica* (KD<sub>50</sub> = 0.74). The lethal doses (LD<sub>50</sub>) of Eos were 2.21 and 0.86 g against flies and cockroach nymphs, respectively. The LD<sub>50</sub> values of the solvent extractives were between 0.41 and 1,746.20 mg cm<sup>2</sup> for houseflies and 0.69 to 2.60 mg cm<sup>2</sup> for cockroaches. The

crude extracts elicited poor growth regulatory and anti-feedant effects [197].

In the western highlands of Cameroon, the incorporation of dry leaves of *C. anisata* into stored food products was investigated using leaf powder, Eos, and anethole (2) against *Callosobruchus maculatus* (*C. maculatus*) and *Callosobruchus chinensis* (Coleoptera, Bruchidae) in stored beans [198]. The dry leaves and Eos evoked significant reduction in the survival rate and number of progeny production, with up to 91% adult mortality and a reduction in adult emergence in treated grain at a dose of 5 g/50 g grain for powder or with 8 µL/40 g grain for Eos. The oils and anethole were markedly toxic to both pests, but anethole had a higher contact toxicity, particularly upon coating it on the grains (LD<sub>50</sub> of Eos = 5, 97 and 4.30 µL/40 g grain and that of anethole = 0.37 and 0.65 µL/40 g grain). Both oils and anethole exhibit strong repellency against test insects [198].

Similarly, *C. anisata* leaf Eos were tested along with other 32 plant Eos and extracts against *C. maculatus* [199]. The number of eggs laid by the beetles on the treated bean seeds was lower than that of the untreated controls. The Eos caused repellence, with 19.8% percentage mortality. A previous study by the same group had indicated that distilled Eos of this species had no repellence activity against *C. maculatus* but prevented oviposition completely and reduced the longevity of adult beetles drastically at doses of 1.25 and 2.50 mL/kg [200]. The insecticidal efficacy of *C. anisata* Eos and an oil-aromatized clay powder against *Acanthoscelides obtectus* (*A. obtectus*) has been investigated [201]. Contact toxicity studies (evaluated through the coating of bean seeds) revealed significant dose-dependent mortality of *A. obtectus*. The aromatized clay powder exhibited greater toxicity (LD<sub>50</sub> = 0.07 µL/g) than Eos (LD<sub>50</sub> = 0.08 µL/g), indicating that the former could be used to stabilize the latter for greater insecticidal efficacy. The Eos and clay powder exhibited reduced toxicity after 24 and 36 h, respectively. The F1 progeny of *A. obtectus* production were reduced by clay and Eos. The highest recorded repellency of the Eos was 90% at 0.25 µL/cm<sup>2</sup>, while it showed high fumigant toxicity (LC<sub>50</sub> = 0.09 µL/cm<sup>3</sup>) against the adult beetles [201].

Govindarajan found that leaf Eos of *C. anisata* (50, 100, 150, 200, and 250 ppm) elicited significant larvicidal activity with 24-hour LC<sub>50</sub> values of 140.96, 130.19, and 119.59 ppm against *Culex quinquefasciatus* (*C. quinquefasciatus*), *Aedes aegypti* (*A. aegypti*), and *Anopheles stephensi* (*A. stephensi*) late III instar larvae [202]. Five isolated pure compounds from the Eos (β-pinene, sabinene, germacrene D, estragole, and linalool) tested at concentrations of 5–75 ppm had LC<sub>50</sub> values of 23.17, 19.67, 16.95, 11.01, and 35.17 ppm against *A. stephensi*; 27.69, 21.20, 18.76, 12.70, and 38.64 ppm against *A. aegypti* and 32.23, 25.01, 21.28, 14.01, and 42.28 ppm against *C. quinquefasciatus* larvae [202]. An assay using Eos from *C. anisata* grown in Tanzania indicated that the oils had low repellency against *Anopheles gambiae* s.s. of the essential oil was low (1.1 × 10<sup>-5</sup> g/cm<sup>2</sup>), unlike those of Western Kenyan plant species, which had no mosquitocidal activity [108, 159]. Toxicity studies on estragole, a major component of most Eos from *C. anisata*, indicate that the compound is a naturally occurring genotoxic carcinogen unsafe for human use [203].

The variation in the constituents of *C. anisata* and *Plectranthus glandulosus* (*P. glandulosus*) Eos, and their levels in stored food products over time was investigated to establish their safety when used in the preservation of food products [204]. Corn grains and flour were treated with plant Eos and stored for 150 days, within which the composition of the Eos was extracted by hydrodistillation and analyzed by gas chromatography/ flame ionization detector. Results indicated the concentration of the oils on food products reduced with storage time (half-life times recorded were 24.16 and 34.61 days for *C. anisata* and 25 and 38.75 days for *P. glandulosus*). At 150 days, only six constituents of *C. anisata* (linalool, estragole, methyl salicylate, *E*-ocimene, thymol, and *E*-nerolidol) and three constituents of *P. glandulosus* (limonene, fenchone, and δ-3-carene) were found on the grains whereas ten (sabinene, linalool, estragole, methyl salicylate, *E*-ocimene, *Z*-ocimene, δ-elemene, *cis*-p-meth-2-en-1-ol, thymol, and *E*-nerolidol) and seven components (limonene, α-terpinolene,

terpinene-4-ol, fenchone, isopulegone-4-methyl, piperitenone oxide, and germacrene D) were present in the flour. Thus, the persistent constituents were more than 62.5 times lower than the toxic concentration observed from the day of treatment, implying that they can safely be used to control insect pests on the corns [204]. With this efficacy, the authors recommended the use of Eos against storage pests, as it had comparable bioactivity to that of imidacloprid [205].

A knowledge-based Cameroonian investigation was undertaken to establish the larvicidal potential of Eos from *C. anisata* leaves and *Dysphania ambrosioides* aerial parts against *C. quinquefasciatus* (the lymphatic filariasis vector) and house flies (*Musca domestica*) [118]. A duplex mixture of the oils was also assessed for probable synergy with the insects, and the mechanism of insecticidal action (inhibitory action on acetylcholinesterase (AChE) was explored. It was found that *C. anisata* Eos were more toxic to third instar larvae of *C. quinquefasciatus* (LC<sub>50</sub> = 29.30 µL/L) than *Musca domestica* (LC<sub>50</sub> = 90.10 µg/adult), while the *Dysphania ambrosioides* Eos was more toxic to *Musca domestica* adults (LC<sub>50</sub> = 51.70 µg/adult) than mosquito larvae (LC<sub>50</sub> = 62.10 µL/L). The mixture of Eos had a synergistic effect against *C. quinquefasciatus* larvae with (LC<sub>50</sub> of 19.30 µL/L), whereas this was not the case with *Musca domestica* adults (LD<sub>50</sub> = 75.90 µg/adult). *C. anisata* Eos did not show any AChE inhibitory effect, though 77 ± 6.86% and 82.2 ± 3.74% AChE inhibition was previously reported at 1 mg/mL of its leaves and twigs methanol extract [136]. As *C. anisata* Eos are known to primarily house phenylpropanoids, their insecticidal activity could be mediated through the inhibition of the activities of cytochrome P450 detoxicative enzymes [206]. In another trial, estragole-rich Eos of *C. anisata* leaves completely inhibited the viability of larvae and the emergence of adult butterflies (*Sitotroga cerealella*) when tested at a dose of 0.50 µL/mL [207].

In another toxicological test, *C. anisata* volatile oils were investigated for their ovicidal, larvicidal, and adulticidal activities against eggs, larvae, and adult female *Anopheles coluzzii* at concentrations of 2%, 1.5%, 1%, and 0.5% [208]. The oils had significant inhibitory potential on *Anopheles coluzzii* eggs, with IC<sub>50</sub> of 0.147%, while 100% mortality was recorded against the 3<sup>rd</sup> instar 48 h post-exposure to *C. anisata* Eos (LC<sub>50</sub> = 0.44%). The knockdown effect of oil-impregnated nets was enhanced with increment in exposure time, with median knockdown times of 1.93, 18.58, 51.02, and 60.15 minutes, respectively for 2%, 1.5%, 1%, and 0.5% dilutions of the oils. The adulticidal effect of the oils 24 h post-exposure had LC<sub>50</sub> value of 1.215% [208].

Mollong et al. assessed the acaricidal efficacy of *C. anisata* Eos on the larvae of *Amblyomma variegatum* (*A. variegatum*) (Acarina: Ixodidae), *Rhipicephalus (Boophilus) (Rh. (B.)) decoloratus*, and *Rh. (B.) microplus* ticks. A concentration of 0.06 µL/cm<sup>2</sup> resulted in 100% larval mortality in the tested tick species, with *A. variegatum* and *Rh. (B.) decoloratus* recording a similar LD<sub>50</sub> of 0.02, whereas the LD<sub>99</sub> of *Rh. (B.) decoloratus* was 0.08 [161]. Although the commercial acaricide (flumethrin) was more effective, with 100% mortality for *A. variegatum* at the lowest dilution of 0.01, these obtained results indicated that *C. anisata* Eos could be harnessed into a potential bio-acaricide [161]. The toxicity of the Eos was again tested on eggs and larvae of *A. variegatum* when diluted with palm kernel and *Jatropha curcas* oils as the vehicle [209]. At 0.12 µL/cm<sup>2</sup>, the Eos inhibited the hatching of 95% of the eggs and provoked 60% mortality of the larvae after 24-h exposure. Despite its toxicity compared to the reference acaricide (bayticol), Eos can be harnessed for managing *A. variegatum* at the preimaginal stages [209]. A continuation of a previous study assessed the effects of *C. anisata* Eos and palm kernel oils on the biological parameters of engorged females of three major tick species in Togo. The Eos of *C. anisata* resulted in 100% mortality in all engorged females. A one-eighth dilution caused the destruction of 100% of eggs laid by *A. variegatum*. Palm kernel oils resulted in 100% mortality in female species of the genus *Rh. (B.)*. The authors indicated that palm kernel oils had appreciable toxicity, indicating that the two oils could be harnessed as an alternative control for tick species.



Eos from *C. anisata* showed 82% fumigant toxicity (insect mortality) at 160  $\mu\text{L/L}$  concentration after 24 hours against *Sitophilus oryzae* (L.) [210]. The oils had a median ( $\text{LC}_{50}$ ) of 17.84  $\mu\text{L/L}$  in fumigant toxicity, whereas maximum repellency (-0.25) was observed at 50  $\mu\text{L/L}$  after 3 h. Eos (3, 6, and 12  $\mu\text{L/L}$ ) were reported to induce a significant reduction in the protein content and total esterase activity of *Sitophilus oryzae* [210].

Another group investigated Eos from *C. anisata*, *Crithmum maritimum*, *Heracleum sphondylium*, *Litchi chinensis* (*L. chinensis*), *Lippia alba*, *Pimpinella anisum*, and *Syzygium aromaticum* for their contact toxicity against the poultry red mite (*Dermanyssus gallinae*). The mechanism of action and efficacy of the oils were assessed using vapor-phase and residual toxicity tests [211]. It was found that Eos of *Syzygium aromaticum* had the highest contact toxicity ( $\text{LC}_{50}$  = 8.90  $\mu\text{g/mL}$ ), followed by *Crithmum maritimum* ( $\text{LC}_{50}$  = 23.70  $\mu\text{g/mL}$ ) and *L. chinensis* Eos ( $\text{LC}_{50}$  = 24.70  $\mu\text{g/mL}$ ). The Eos of *C. anisata* and *L. chinensis* showed the highest vapor toxicity among the Eos.

**Extracts.** The larvicidal potential of some selected traditional South African plant extracts, including that of *C. anisata* was assessed against *Anopheles arabiensis* to verify their use as mosquito repellents [38]. Dose-response bioassay results indicated that *C. anisata* leaf extracts had 100% and 88% larvicidal activity at 500  $\mu\text{g/mL}$  and 250  $\mu\text{g/mL}$  with  $\text{LC}_{50}$  of 112.70  $\mu\text{g/mL}$ . Furthermore, the antifeedant activities of aqueous and organic (hexane, petroleum ether, chloroform, ethyl acetate, and methanolic extracts of *C. anisata* leaves and roots) extracts were evaluated against *Helicoverpa armigera* (Hübner) [23]. Root extracts had higher antifeedant activities, with petroleum ether and chloroform exhibiting the strongest activities, with concentrations eliciting 50% deterrence of 0.014% and 0.016%, respectively. Further fractionation of the root extracts resulted in one chloroform and one petroleum ether fraction that were active owing to the presence of osthol. The highest osthol concentration was observed in the chloroform root extract. The antifeedant activities of the root extracts, as measured by the 50% deterrence values, were highly correlated with their osthol contents. Approximately 99% of the variation in bioactivity of the root extracts could be accounted for by variation in osthol content; osthol, therefore, appeared to be an antifeedant component of *C. anisata* to *Helicoverpa armigera* [23].

The application of aqueous extracts of *C. anisata* leaves and stem bark along with other plant extracts to nematode-inoculated tomato plants was reported to significantly reduce (up to 86%) tomato root galling [135]. In another study, powdered *C. anisata* leaves admixed with maize grains were tested on Cameroonian and German strains of *Sitophilus zeamais* and *Prostephanus truncatus* to assess their insecticidal potential and their effects on progeny production, grain damage, and population increase [212]. Significant treatment, exposure time, and insect strain affected the adulticidal potential of the powder. The treatment also significantly affected the number of progeny produced, and the percentage reduction in adult emergence relative to the control, while 3-months monitoring results indicated that the powder significantly reduced the rate of increase in the population of the insects [212].

Experiments were performed to evaluate the fungitoxicity of methylene chloride/methanol leaf extracts of seven plants (including *C. anisata* leaves) on *Phytophthora infestans* development [213]. Plant extracts at 3% were assessed for their potential to inhibit/retard sporangial germination, latent and incubation periods, lesion size, and late blight severity in tomato plants. *C. anisata* along with *Entandrophragma angolense*, *Ageratum houstonianum*, and *Tephrosia vogelli* extracts provided excellent inhibition (3, 3.68, 0.18, and 2.18  $\text{cm}^2$ , respectively) of late blight disease progression, with comparable fungitoxic effects to that of the commercial fungicides metalaxyl (0.40  $\text{cm}^2$ ) and maneb (1.33  $\text{cm}^2$ ), utilized as the positive controls in the study. The study concluded that the extracts could be incorporated into an integrated pathogen management program for tomato late blight [213]. Another report assessed the efficacy and toxicity of some acetic extracts of plants used in ethnoveterinary medicine in South Africa on egg hatching and larval development of *Haemonchus contortus* [214]. *C. anisata* had  $\text{EC}_{50}$  values of  $1.80 \pm 0.09$  and  $2.07 \pm$

0.15  $\text{mg/mL}$  in egg hatching and larval development assays, respectively. The extract was the least cytotoxic, with an  $\text{LC}_{50}$  of 0.17  $\text{mg/mL}$  and a selectivity index of 0.10 and 0.08 for the two assays.

The larvicidal activity of 1,2-benzenedicarboxylic acid, mono (2-ethylhexyl) ester (339) isolated from the ethyl acetate extract of leaves of *C. anisata* was evaluated in the fourth instar larvae of *C. quinquefasciatus*, *A. aegypti*, and *A. stephensi*. The compound elicited 100% larval mortality against *A. aegypti* and *A. stephensi* at 40 ppm, with  $\text{LC}_{50}$  values of 8.94 and 9.23 ppm [156]. The compound showed 98% mortality, with an  $\text{LC}_{50}$  value of 12.07 ppm against *C. quinquefasciatus*.

The different bioactivities of *C. anisata* extracts and oils against insects have been attributed to the abundance of compounds such as coumarins, limonoids, and alkaloids [215]. For example, coumarins have been known to elicit insecticidal, feeding deterrence, and repellent activities through bioenergetic disruption of muscles or exerting neurotoxicity by ACE inhibition by blocking the octopamine receptor [216, 217]. Specifically, furanocoumarins may alter the detoxification capacity of organisms through the reversible and/or irreversible inhibition of cytochrome P450 detoxification enzymes [218]. On the other hand, alkaloids deter feeding by inhibiting impulse generation in sugar-sensitive cells in lepidopterans or by competitively blocking sucrose responses [14]. Their insecticidal potential varies for different target insects but is usually mediated by exerting effects on ACE and sodium channels [219, 220]. Limonoids, on the other end of the spectrum, have recognized insect antifeedant activities, usually mediated through suppression of the phagostimulant receptor and stimulation of deterrent cells located in the medial sensillum of insects [221]. The growth-regulating activities of this chemical group are achieved through the direct action on enzymes and other physiological regulatory processes necessary for normal insect growth or insect progenies [14].

#### Molecular docking studies

In silico methods have gained attention in modern drug design because they have the ability to understand drug-receptor interactions and quantum chemical properties. Eswaremoorthy et al. confirmed the mode of antibacterial binding and other properties of heptaphyline (35) and three coumarins (36, 40, and 63) isolated from *C. anisata* [60, 222]. Docking studies showed that compound 40 exhibited better docking scores against both DNA gyrase B and the LasR-binding domain compared to ciprofloxacin. In contrast, toxicological predictions revealed that these compounds are non-cytotoxic, non-hepatotoxic, non-carcinogenic, non-irritant, and immunogenic. Compound 40 had a large electronegativity, mutagenicity value, global softness, and global electrophilicity comparable to that of ciprofloxacin, suggesting that it could elicit better bioactivity and chemical reactivity with considerable intramolecular charge transfer between the electron-donor and electron-acceptor groups [222]. Molecular docking of *trans*-4-hydroxy-1-methyl-L-proline (90) from *C. anisata* leaf and stem extracts indicated a GOLD fitness score of 31.26 with only a single hydrogen bond appreciated with residue Glu208 in the active sites of lipase and hyaluronidase enzymes [157]. Analysis of the active site of the hyaluronidase enzyme indicated that its binding residues were Ala221, Leu222, Gly223, Thr224, Leu225, Lys226, Ile227, Thr271, Gly273, Lys274, and Leu275. Active site docking of compound 90 gave a GOLD fitness score of 17.20, which was lower than that of lipase [157].

#### Toxicity profile of *C. anisata* extracts and isolated compounds

Toxicological evaluation of medicinal plants, isolated pure compounds, and corresponding herbal products is pivotal for enhancing their approval and licensing as pharmaceutical products. As summarized in Table 7, it can be deduced that extracts from *C. anisata* are relatively non-toxic, which could also explain their widespread use in traditional medicine [223, 224]. Irungu et al. focused on the safety of extracts from *C. anisata* for specific solvents [51].

Table 7 Toxicity of *C. anisata* as reported in open literature

Toxicity <sup>a</sup>	Assay	Inferences	Reference (s)
LD <sub>50</sub> = 948.68 mg/kg and 1,265 mg/kg (ethanol extract of leaves, root bark and stem bark)	Acute oral toxicity test	Moderate toxicity	[132]
LD <sub>50</sub> = 27,172.5 mg/L (aqueous leaf extracts)	Brine shrimp lethality assay	Low toxicity	[37]
LC <sub>50</sub> > 100 µg/mL (ethanol and ethyl acetate extracts of fruits, leaves, root and stem bark), LC <sub>50</sub> = < 20 µg/mL (corresponding chloroform extracts)	Brine shrimp lethality assay	Nontoxic	[174]
Leaf ethanolic extracts (up to 5,000 mg/kg)	Acute oral toxicity test	Moderately toxic. No mortality up to 14 <sup>th</sup> day but toxicity behaviors noted	[75]
LD <sub>50</sub> = 393.7 mg/kg (leaf methanolic extract)	Acute oral toxicity test	Toxic as mortality was recorded	[6]
LD <sub>50</sub> = 17,036.85 mg/kg bw (leaf ethanolic extract)	Acute oral toxicity test	Nontoxic to white mouse	[223]
LC <sub>50</sub> = 71.9 and 60.5 µg/mL (roots DCM and ethanol extracts)	Brine shrimp lethality assay	Potentially toxic	[224]
LC <sub>50</sub> = 533.0 µg/mL (leaves methanolic extract)	Brine shrimp lethality assay	Moderately toxic	[82]

LC<sub>50</sub>, median lethal concentration; LD<sub>50</sub>, median lethal dose; DCM, dichloromethane; *C. anisata*, *Clausena anisata* (Willd.) Hook.f. ex Benth.

The authors recorded no mortality in mice that received hexane extract of *C. anisata* stem bark, whereas the LD<sub>50</sub> of animals administered chloroform extract was 4,166.70 mg/kg. This mild oral acute toxicity was speculated to be due to the abundance of cytotoxic carbazole alkaloids or coumarins (such as chalepin), which are known to be mutagenic and toxic, with an LD<sub>50</sub> of 100 mg/kg in rats [88, 139, 140, 225]. This is substantiated by another report wherein the micromolar addition of chalepin led to the inhibition of rat liver mitochondrial respiration [149]. The authors inferred that chalepin (i) inhibits pyruvate/malate oxidation, (ii) does not interrupt succinate oxidation, (iii) inhibits site I phosphorylation, and (iv) is approximately one-tenth that of rotenone, a natural coumarin insecticide [149]. These results imply that further studies are required to provide sufficient evidence of the toxicity of the isolated compounds.

#### Clinical studies

Clinical validation of pharmaceutical and herbal products is mandatory and important as it promotes their acceptance and use. However, throughout this exhaustive review, only one clinical trial utilizing an herbal product from *C. anisata* was performed. This single-blind, randomized controlled trial of a Ghanaian herbal formulation (*C. anisata* and *Cassia sieberiana* D.C. roots) assessed the safety and effectiveness of the product in managing patients with osteoarthritis [226]. After 8 weeks of using the product by 44 subjects, the intensity of pain was reduced, and the disability score and days similarly declined. These observations support the use of herbal medicines. Toxicological assessments further indicated that the product was safe for human use. More clinical studies on this species are required to unravel its potential in the management of diseases across Africa.

#### Conclusion

*C. anisata* has a long history of use as an herbal medicine in Africa. Significant strides have been made over the last five decades to isolate, elucidate, and evaluate its phytoconstituents and the pharmacological activities of those constituents. Both crude extracts and isolated compounds from *C. anisata* possess anti-inflammatory, antimicrobial, antioxidant, antiplasmodial, antiproliferative, anti-human immunodeficiency virus 1, antimycobacterial, antihypertensive, and antidiabetic activities. The relevant bioactive compounds include alkaloids, coumarins, limonoids, and phenylpropanoids. Although some studies have indicated moderate

toxicity of the extracts, some of the dominant compounds in this species, such as estragole and carbazole alkaloids, are mutagenic or cytotoxic. Accumulating evidence supports the traditional use of this species in herbal medicine. However, these studies do not explain the relationships between traditional uses, pharmacological activities, and underlying molecular mechanisms. Thus, future research studies should focus on the structure-bioactivity relationships and clinical potential of crude extracts and isolated compounds from *C. anisata*.

#### References

- Suksaeree J, Chankana N, Luprasong C, Monton C. Optimization of dynamic maceration of *Clausena anisata* (Willd.) Hook. f. ex Benth. leaves to maximize trans-anethole content. *SN Appl Sci*. 2021;3(4):514. <https://doi.org/10.1007/s42452-021-04509-4>
- Quattrocchi U. CRC World Dictionary of Plant Names: Common Names, Scientific Names, Eponyms, Synonyms, and Etymology. 2012;5:3960. [https://www.studmed.ru/quattrocchi-u-crc-world-dictionary-of-plant-names-common-names-scientific-names-eponyms-synonyms-and-etymology-vol-1-a-c\\_0cbc692ce1.html](https://www.studmed.ru/quattrocchi-u-crc-world-dictionary-of-plant-names-common-names-scientific-names-eponyms-synonyms-and-etymology-vol-1-a-c_0cbc692ce1.html)
- Tchinda AT. *Clausena anisata* (Willd.) Hook. f. Ex Benth. Schmelzer GH, Gurib-Fakim A (eds.) Prota 11(2): Medicinal plants/PlantesMédicinales 2. PROTA; 2011. [https://www.prota4u.org/database/protav8.asp?p=Clausena+anisata+\(Willd.\)+Hook.f.+ex+Benth](https://www.prota4u.org/database/protav8.asp?p=Clausena+anisata+(Willd.)+Hook.f.+ex+Benth)
- Zhang D, Hartley TG. Flora of China. 2008;11:83–85. <http://flora.huh.harvard.edu/china/PDF/PDF11/Clausena.pdf>.
- The Medicinal Plant Names Services. *Clausena anisata* (Willd.) Hook.f. ex Benth. 2021. <https://mpns.science.kew.org/mpns-portal/plantDetail?plantId=724765&query=clausena+anisata&filter=&fuzzy=false&nameType=all&db=wcsCmp>
- Okokon JE, Udoh AE, Andrew UE, Amazu LU. Antiinflammatory and antipyretic activities of *Clausena anisata*. *Mol Clin Pharmacol*. 2012;3(1):47–54. [https://www.researchgate.net/profile/Anwanabasi-Udoh/publication/345243821\\_Anti-inflammatory\\_and\\_antipyretic\\_activities\\_of\\_Clausena\\_anisata/links/5fa15f0092851c14bcff6ae9/Anti-inflammatory-and-antipyretic-activities-of-Clausena-anisata.pdf](https://www.researchgate.net/profile/Anwanabasi-Udoh/publication/345243821_Anti-inflammatory_and_antipyretic_activities_of_Clausena_anisata/links/5fa15f0092851c14bcff6ae9/Anti-inflammatory-and-antipyretic-activities-of-Clausena-anisata.pdf)
- BioTik. *Clausena anisata* (Willd.) J.Hk. ex Benth-RUTACEAE. 2021. [http://www.biotik.org/india/species/c/clauanis/clauanis\\_en.html](http://www.biotik.org/india/species/c/clauanis/clauanis_en.html)

8. Wikimedia Commons. File: *Clausena anisata* 1.jpg. 2021. [https://commons.wikimedia.org/wiki/File:Clausena\\_anisata\\_1.jpg](https://commons.wikimedia.org/wiki/File:Clausena_anisata_1.jpg)
9. Hyde MA, Wursten BT, Ballings P, Coates Palgeave M. *Flora of Zimbabwe: Species Information: records of: Clausena anisata*; 2022. [https://www.zimbabweflora.co.zw/speciesdata/species-display.php?species\\_id=133210](https://www.zimbabweflora.co.zw/speciesdata/species-display.php?species_id=133210)
10. Agyepong N, Agyare C, Adarkwa-Yiadom M, Gbedema SY. Phytochemical investigation and anti-microbial activity of *Clausena anisata* (Willd.) Hook. *Afr J Tradit Complement Altern Med*. 2014;11(3):200–209. <https://doi.org/10.4314/ajtcam.v11i3.28>
11. Lunyera J, Wang D, Maro V, et al. Traditional medicine practices among community members with diabetes mellitus in Northern Tanzania: an ethnomedical survey. *BMC Complement Altern Med*. 2016;16(1):282. <https://doi.org/10.1186/s12906-016-1262-2>
12. Schmelzer GH, Aguilar NO. Record from Proseabase. PROSEA (Plant Resources of South-East Asia) Foundation, Bogor, Indonesia. 2001. <https://www.cabi.org/isc/abstract/20013181428>
13. Molino JF. Révision du genre *Clausena* Burm. f. (Rutaceae). *Bull Mus Nat Hist Nat, Paris Adansonia*. 1994;1:105–153. [https://horizon.documentation.ird.fr/exl-doc/pleins\\_textes/pleins\\_textes\\_7/b\\_fdi\\_57-58/010024458.pdf](https://horizon.documentation.ird.fr/exl-doc/pleins_textes/pleins_textes_7/b_fdi_57-58/010024458.pdf)
14. Mukandiwa L, Naidoo V, Katerere DR. The use of *Clausena anisata* in insect pest control in Africa: a review. *J Ethnopharmacol*. 2016;194:1103–1111. <https://doi.org/10.1016/j.jep.2016.11.002>
15. Coates Palgrave K, Coates Palgrave P, Coates Palgrave M. *Everyone's Guide to Trees of South Africa* (fourth ed.). Cape Town, South Africa: Struik, Publishers; 2000. ISBN:1868724891
16. Bussmann RW, Paniagua-Zambrana NY, Njoroge GN. *Clausena anisata* (Willd.) Hook. f. ex Benth. Rutaceae. Bussmann RW (eds) *Ethnobotany of the Mountain Regions of Africa*. Springer, Cham. 2021;289–291. [https://doi.org/10.1007/978-3-030-38386-2\\_40](https://doi.org/10.1007/978-3-030-38386-2_40)
17. Meijer TM. The essential oil from the leaves of *Clausena anisata* Hook. f. *Recueil des Travaux Chimiques des Pays-Bas*. 1947;66(6):395–400. <https://doi.org/10.1002/recl.19470660608>
18. Lawal IO, Uzokwe VNE, Rafiu B, Afolayan AJ. Foliar anatomy of *Clausena anisata* (Willd.) Hook: a South African medicinal plant. *Niger J Nat Prod Med*. 2018;22:149–154. <https://www.ajol.info/index.php/njnpm/article/view/178105>
19. Kadiri M, Ojewumi AW, Onatade T. Indigenous uses and phytochemical contents of plants used in the treatment of menstrual disorders and after-child birth problems in abeokuta south local government of Ogun state, Nigeria. *J Drug Del Therapeut*. 2015;5(3):33–42. <https://doi.org/10.22270/jddt.v5i3.1144>
20. Jeon CM, Shin IS, Shin NR, et al. *Clausena anisata*-mediated protection against lipopolysaccharide-induced acute lung injury in mice. *Int J Mol Med*. 2016;37(4):1091–1098. <https://doi.org/10.3892/ijmm.2016.2515>
21. KewScience. *Plants of the World Online. Clausena anisata* (Willd.) Hook.f. ex Benth. 2021. <http://www.plantsoftheworldonline.org/taxon/urn:lsid:ipni.org:names:772111-1>
22. Ajene I, Khamis FM, Pietersen G, Asch BV. Mitochondrial genetic variation reveals phylogeographic structure and cryptic diversity in *Triozia erytraea*. *Sci Rep*. 2020;10(1):8893. <https://doi.org/10.1038/s41598-020-65880-7>
23. Pitan OOR, Ayelaagbe OO, Wang HL, Wang CZ. Identification, isolation and characterization of the antifeedant constituent of *Clausena anisata* against *Helicoverpa armigera* (Lepidoptera: noctuidae). *Insect Sci*. 2009;16(3):247–253. <https://doi.org/10.1111/j.1744-7917.2009.01255.x>
24. Kémeuzé VA, Solefack MCM, Nkongmeneck BA, Decocq G, Jiofack T, Madeleine J. Variation altitudinale de la distribution des plantes à activité insecticide dans la forêt communautaire de Kilum-Ijim: cas de *Clausena anisata* (Willd.) Hook. f. ex-Benth. *Bois et Forêts des Tropiques*. 2009;299:71–77. <https://doi.org/10.19182/bft2009.299.a20425>
25. Monton C, Suksaeree J, Luprasong C. Validated reversed-phase ion-interaction high-performance liquid chromatography for quantitation of nitrate content of *Clausena anisata* (Willd.) Hook. f. ex Benth. Leaves. *Scientifica (Cairo)*. 2020;2020:6424682. <https://doi.org/10.1155/2020/6424682>
26. Avlessi F, Dangou J, Wotto VD, Alitonou GA, Sohounhloue DK, Menut C. Propriétés antioxydantes de l'huile essentielle des feuilles de *Clausena anisata* (Willd.) Hook. *Comptes Rendus Chimie*. 2004;7(10–11):1057–1061. <https://doi.org/10.1016/j.crci.2003.11.015>
27. Ayedoun MA, Houenon JG, Sossou PV, Menut C, Lamaty G, Bessiere JM. Aromatic plants of tropical West Africa. v. major components of the leaf and fruit Eos of *Clausena anisata* (willd.) j.d. hook from Benin. *J Essent Oil Res*. 1997;9(2):247–248. <https://doi.org/10.1080/10412905.1997.9699473>
28. Axtell BL, Fairman RM. Minor oil crops (Part III. Eos). *FAO Agricultural Services Bulletin*. 1992:94. <https://www.fao.org/3/X5043E/x5043E00.htm>
29. Maina FM. *Investigation of Volatile Oil Of Clausena Anisata Growing In Kenya*. BSc Research Report, University of Nairobi, Nairobi, Kenya;1983. <http://erepository.uonbi.ac.ke/bitstream/handle/11295/96874/Maina-Investigation%20Of%20Volatile%20Oil%20Of%20Clausena%20Anisata%20Growing%20In%20Kenya.pdf?sequence=1>
30. Mavundza EJ, Maharaj R, Kabera G, Finnie JF, Van Staden J. An ethnobotanical survey of mosquito repellent plants in uMkhanyakude district, KwaZulu-Natal Province, South Africa. *J Ethnopharmacol*. 2011;137(3):1516–1520. <https://doi.org/10.1016/j.jep.2011.08.040>
31. Lawal IO, Grierson DS, Afolayan AJ. Phytochemical and antioxidant investigations of a *Clausena anisata* hook, a South African medicinal plant. *Afr J Tradit Complement Altern Med*. 2015;12(1):28–37. <https://doi.org/10.4314/ajtcam.v12i1.5>
32. van Vuuren S, Frank L, Review: Southern African medicinal plants used as blood purifiers. *J Ethnopharmacol*. 2020;249:112434. <https://doi.org/10.1016/j.jep.2019.112434>
33. Mkhombo MH. The effects of *Clausena anisata* (Willd) Hook (Rutaceae) leaf extracts on selected diabetic related carbohydrate metabolizing enzymes. MSc Thesis, University of Limpopo, South Africa; 2010. <http://ulspace.ul.ac.za/bitstream/handle/10386/272/M%20H%20MKHOMBO%20%20THESIS%20MSC%20IN%20BIOCHEMISTRY%202010.pdf?sequence=1>
34. Lawal IO, Grierson DS, Afolayan AJ. Phytotherapeutic Information on plants used for the treatment of tuberculosis in Eastern Cape Province, South Africa. *Evid-Based Complement Alternat Med*. 2014;2014:735423. <https://doi.org/10.1155/2014/735423>
35. York T, van Vuuren SF, de Wet H. Plants used for treating respiratory infections in rural Maputaland, KwaZulu-Natal, South Africa. *J Ethnopharmacol*. 2011;135(30):696–710. <https://doi.org/10.1016/j.jep.2011.03.072>
36. Asowata-Ayodele AM, Afolayan AJ, Otunola GA. Ethnobotanical survey of culinary herbs and spices used in the traditional medicine system of Nkonkobe Municipality, Eastern Cape, South Africa. *South Afr J Bot*. 2016;104:69–75. <https://doi.org/10.1016/j.sajb.2016.01.001>
37. Lechaba NMT, Schutte PJ, Hay L, Böhmer L, Govender MM. The effects of an aqueous leaf extract of *Clausena anisata* (Willd.) Hook.f.ex Benth. on blood pressure, urine output, angiotensin II



- levels and cardiac parameters in spontaneously hypertensive rats. *J Med Plants Res.* 2016;10(28):425–434. <https://doi.org/10.5897/JMPR2016.6135>
38. Mavundza EJ, Maharaj R, Chukwujekwu JC, Finnie JF, Van Staden J. Larvicidal activity against *Anopheles arabiensis* of 10 South African plants that are traditionally used as mosquito repellents. *South Afr J Bot.* 2013;88:86–89. <https://doi.org/10.1016/j.sajb.2013.05.007>
  39. Mukungu N, Abuga K, Okalebo F, Ingwela R, Mwangi J. Medicinal plants used for management of malaria among the Luhya community of Kakamega East sub-County, Kenya. *J Ethnopharmacol.* 2016;194:98–107. <https://doi.org/10.1016/j.jep.2016.08.050>
  40. Okello SV, Nyunja RO, Netondo GW, Onyango JC. Ethnobotanical study of medicinal plants used by Sabaots of Mt. Elgon Kenya. *Afr J Tradit Complement Altern Med.* 2009;7(1):1–10. <https://doi.org/10.4314/ajtcam.v7i1.57223>
  41. Muthaura CN, Rukunga GM, Chhabra SC, Mungai GM, Njagi ENM. Traditional phytotherapy of some remedies used in treatment of malaria in Meru district of Kenya. *South Afr J Bot.* 2007;73(3):402–411. <https://doi.org/10.1016/j.sajb.2007.03.004>
  42. Nguta JM, Mbaria JM, Gakuya DW, Gathumbi PK, Kiama SG. Traditional antimalarial phytotherapy remedies used by the South Coast community, Kenya. *J Ethnopharmacol.* 2010;131(2):256–267. <https://doi.org/10.1016/j.jep.2010.06.031>
  43. Kigen G, Kipkore W, Wanjohi B, Haruki B, Kemboi J. Medicinal plants used by traditional healers in Sangurur, Elgeyo Marakwet County, Kenya. *Pharmacognosy Res.* 2017;9(4):333–347. [https://doi.org/10.4103/pr.pr.42\\_17](https://doi.org/10.4103/pr.pr.42_17)
  44. Kokwaro JO. *Medicinal Plants of East Africa* (2nd Eds). Nairobi, Kenya: East African Literature Bureau;1993. <http://41.204.161.209/handle/11295/34820>
  45. Rufford. Ethnomedicine of Tugen Community, Baringo County-Kenya;2020. [https://ruffordorg.s3.amazonaws.com/media/project\\_reports/19802-1%20Medicinal%20Plants%20of%20Baringo%2C%20Kenya.pdf](https://ruffordorg.s3.amazonaws.com/media/project_reports/19802-1%20Medicinal%20Plants%20of%20Baringo%2C%20Kenya.pdf)
  46. World Agroforestry Center, Useful trees and shrubs for Kenya. *Clausena anisata.* 2014. [http://www.worldagroforestry.org/usefultrees/pdflib/Clausena\\_anisata\\_KEN.pdf](http://www.worldagroforestry.org/usefultrees/pdflib/Clausena_anisata_KEN.pdf)
  47. Muthee JK, Gakuya DW, Mbaria JM, Kareru PG, Mulei CM, Njonge FK. Ethnobotanical study of anthelmintic and other medicinal plants traditionally used in Loitokitok district of Kenya. *J Ethnopharmacol.* 2011;135(1):15–21. <https://doi.org/10.1016/j.jep.2011.02.005>
  48. Mbuni YM, Wang S, Mwangi BN, et al. Medicinal plants and their traditional uses in local communities around cherangani hills, Western Kenya. *Plants (Basel).* 2020;9(3):331. <https://doi.org/10.3390/plants9030331>
  49. Ichikawa M. A preliminary report on the ethnobotany of the Suiei Dorobi in Northern Kenya. *Afr Stud Monographs.* 1987;7:1–52. <https://doi.org/10.14989/68347>
  50. Pakia M. Plant ecology and ethnobotany of two sacred forests (kayas) at the Kenya coast. MSc Thesis, University of Natal, Durban; 2000. <https://ukzn-dspace.ukzn.ac.za/handle/10413/3829>
  51. Irungu BN, Mbabu MJ, Kiboi DM, Moindi E, Kinyua J, Romano K M. In vivo antimalarial and acute toxicity properties of hexane and chloroform extracts from *Clausena anisata* (Willd.) Benth. *Afr J Pharmacol Therapeut.* 2012;1:24–29. <http://repository.embuni.ac.ke/handle/123456789/309>
  52. Muthaura CN, Keriko JM, Mutai C, et al. Antiplasmodial potential of traditional phytotherapy of some remedies used in treatment of malaria in Meru-Tharaka Nithi County of Kenya. *J Ethnopharmacol.* 2015;175:315–323. <https://doi.org/10.1016/j.jep.2015.09.017>
  53. Kemoli AM, van Amerongen WE, de Soet JJ. Antimicrobial and buffer capacity of crude extracts of chewing sticks (Miswaki) from Kenya. *ASDC J Dent Child.* 2001;68(3):183–188,152. <https://pubmed.ncbi.nlm.nih.gov/11693010/>
  54. Gumisiriza H, Birungi G, Olet EA, Sesaazi CD. Medicinal plant species used by local communities around Queen Elizabeth National Park, Maramagambo Central Forest Reserve and Ihimbo Central Forest Reserve, South western Uganda. *J Ethnopharmacol.* 2019;239:111926. <https://doi.org/10.1016/j.jep.2019.111926>
  55. Gradé JT, Tabuti JRS, Van Damme P. Ethnoveterinary knowledge in pastoral Karamoja, Uganda. *J Ethnopharmacol.* 2009;122(20):273–293. <https://doi.org/10.1016/j.jep.2009.01.005>
  56. Tabuti JRS, Lye KA, Dhillion SS. Traditional herbal drugs of Bulamogi, Uganda: plants, use and administration. *J Ethnopharmacol.* 2003;88(1):19–44. [https://doi.org/10.1016/s0378-8741\(03\)00161-2](https://doi.org/10.1016/s0378-8741(03)00161-2)
  57. Garneau FX, Pichette A, Oteng-Yeboah A, et al. (E)- and (Z)-Foeniculin, constituents of the leaf oil of a new chemovariety of *Clausena anisata*. *J Essent Oil Res.* 2000;12(6):757–762. <https://doi.org/10.1080/10412905.2000.9712209>
  58. Yetein MH, Houessou LG, Loubégnon TO, Tekka O, Tente B. Ethnobotanical study of medicinal plants used for the treatment of malaria in plateau of Allada, Benin (West Africa). *J Ethnopharmacol.* 2013;146(1):154–163. <https://doi.org/10.1016/j.jep.2012.12.022>
  59. Allabi AC, Busia K, Ekanmian V, Bakiono F. The use of medicinal plants in self-care in the Agonlin region of Benin. *J Ethnopharmacol.* 2011;133(1):234–243. <https://doi.org/10.1016/j.jep.2010.09.028>
  60. Tamene D, Endale M. Antibacterial activity of coumarins and carbazole alkaloid from roots of *Clausena anisata*. *Adv Pharmacol Sci.* 2019;2019:5419854. <https://doi.org/10.1155/2019/5419854>
  61. Tesfaye S, Belete A, Engidawork E, Gedif T, Asres K. Ethnobotanical study of medicinal plants used by traditional healers to treat cancer-like symptoms in eleven districts, Ethiopia. *Evid Based Complement Alternat Med.* 2020;2020:7683450. <https://doi.org/10.1155/2020/7683450>
  62. Teklehaymanot T, Giday M. Ethnobotanical study of medicinal plants used by people in Zegie Peninsula, Northwestern Ethiopia. *J Ethnobiol Ethnomed.* 2007;3:12. <https://doi.org/10.1186/1746-4269-3-12>
  63. Kefalew A, Asfaw Z, Kelbessa E. Ethnobotany of medicinal plants in Ada'a District, East Shewa Zone of Oromia Regional State, Ethiopia. *J Ethnobiol Ethnomed.* 2015;11:25. <https://doi.org/10.1186/s13002-015-0014-6>
  64. Abera B. Medicinal plants used in traditional medicine by Oromo people, Ghimbi District, Southwest Ethiopia. *J Ethnobiol Ethnomed.* 2014;10:40. <https://doi.org/10.1186/1746-4269-10-40>
  65. Mekuanent T, Zebene A, Solomon Z. Ethnobotanical study of medicinal plants in Chilga District, Northwestern Ethiopia. *J Nat Remed.* 2015;15(2):88–112. <https://doi.org/10.18311/jnr%2F2015%2F476>
  66. Teklehaymanot T. Ethnobotanical study of knowledge and medicinal plants use by the people in Dek Island in Ethiopia. *J Ethnopharmacol.* 2009;124(1):69–78. <https://doi.org/10.1016/j.jep.2009.04.005>
  67. Giday M, Asfaw Z, Woldu Z. Ethnomedicinal study of plants used by Sheko ethnic group of Ethiopia. *J Ethnopharmacol.* 2010;132(1):75–85. <https://doi.org/10.1016/j.jep.2010.07.046>
  68. Suleman S, Alemu T. A survey on utilization of ethnomedicinal

- plants in Nekemte town, East Wellega (Oromia), Ethiopia. *J Herbs Spices Med Plants*. 2012;18(1):34–57. <https://doi.org/10.1080/10496475.2011.645188>
69. Alemneh D. Ethnobotanical study of plants used for human ailments in Yilmana Densa and Quarit Districts of West Gojjam Zone, Amhara Region, Ethiopia. *BioMed Res Int*. 2021;2021:6615666. <https://doi.org/10.1155/2021/6615666>
  70. Kidane B, van der Maesen LJG, van Andel T, Asfaw Z. Ethnoveterinary medicinal plants used by the Maale and Ari ethnic communities in Southern Ethiopia. *J Ethnopharmacol*. 2014;153(1):274–282. <https://doi.org/10.1016/j.jep.2014.02.031>
  71. Van Vuuren SF, Viljoen AM. The in vitro antimicrobial activity of toothbrush sticks used in Ethiopia. *South Afr J Bot*. 2006;72(4):646–648. <https://doi.org/10.1016/j.sajb.2006.03.009>
  72. Hamza OJM, van den Bout-van den Beukel CJP, Matee MIN, et al. Antifungal activity of some Tanzanian plants used traditionally for the treatment of fungal infections. *J Ethnopharmacol*. 2006;108(1):124–132. <https://doi.org/10.1016/j.jep.2006.04.026>
  73. Moshi MJ, Kagashe GAB, Mbwambo ZH. Plants used to treat epilepsy by Tanzanian traditional healers. *J Ethnopharmacol*. 2005;97(2):327–336. <https://doi.org/10.1016/j.jep.2004.11.015>
  74. Ajibesin KK, Ekpo BA, Bala DN, Essien EE, Adesanya SA. Ethnobotanical survey of Akwa Ibom State of Nigeria. *J Ethnopharmacol*. 2008;115(3):387–408. <https://doi.org/10.1016/j.jep.2007.10.021>
  75. Ishola IO, Olayemi SO, Oreagba IA, Ifeanyi CI, Popoola TO. Antinociceptive and anti-arthritic properties of hydroethanolic leaf extract of *Clausena anisata* (Willd.) Hook. f. ex Benth (Rutaceae) in Rodents: possible mechanism of actions. *Niger J Physiol Sci*. 2015;30(1–2):39–49. <https://pubmed.ncbi.nlm.nih.gov/27506169/>
  76. Iwu MM. *Handbook of African Medicinal Plants*. New York, CRC Press; 1994:164–166. ISBN:9780429101397
  77. Segun PA, Ogbale OO, Ajaiyeoba EO. Medicinal plants used in the management of cancer among the Ijebus of Southwestern Nigeria. *J Herbal Med*. 2018;14:68–75. <https://doi.org/10.1016/j.hermed.2018.04.002>
  78. Ndukwe KC, Okeke IN, Lamikanra A, Adesina SK, Aboderin O. Antibacterial activity of aqueous extracts of selected chewing sticks. *J Contemp Dent Pract*. 2005;6(3):86–94. <https://pubmed.ncbi.nlm.nih.gov/16127476/>
  79. Ogbale OO, Segun PA, Adeniji AJ. In vitro cytotoxic activity of medicinal plants from Nigeria ethnomedicine on Rhabdomyosarcoma cancer cell line and HPLC analysis of active extracts. *BMC Complement Altern Med*. 2017;17(1):494. <https://doi.org/10.1186/s12906-017-2005-8>
  80. Kpodar MS, Karou SD, Katawa G, et al. An ethnobotanical study of plants used to treat liver diseases in the Maritime region of Togo. *J Ethnopharmacol*. 2016;181:263–273. <https://doi.org/10.1016/j.jep.2015.12.051>
  81. Koudouvo K, Karou DS, Kokou K, et al. An ethnobotanical study of antimalarial plants in Togo Maritime Region. *J Ethnopharmacol*. 2011;134(1):183–190. <https://doi.org/10.1016/j.jep.2010.12.011>
  82. Tafadzwa M. Screening of some traditional medicinal plants from Zimbabwe for biological and anti-microbial activity. MPhil Thesis, University of Zimbabwe; 2005. <http://41.175.146.201/handle/10646/964>
  83. Tomani JCD, Gankam LOT, Nshutiyayesu S, et al. An ethnobotanical survey and inhibitory effects on NLRP3 inflammasomes/Caspase-1 of herbal recipes' extracts traditionally used in Rwanda for asthma treatment. *J Ethnopharmacol*. 2018;227:29–40. <https://doi.org/10.1016/j.jep.2018.08.016>
  84. Senthilkumar A, Venkatesalu V. Phytochemical analysis and antibacterial activity of essential oil of *Clausena anisata* (Willd.) Hook. f. ex Benth. *Int J Integr Biol*. 2009;5(2):116–120. <https://www.semanticscholar.org/paper/Phytochemical-analysis-and-antibacterial-activity-Senthilkumar-Venkatesalu/2ff9640a2dd7793e2d45ab6f58b87f352fa818f>
  85. MMS. *Clausena anisata*; 2021. <https://www.yumpu.com/en/document/view/11003169/constituents-mmh-mms>
  86. Arbab IA, Abdul AB, Aspollah M, Abdullah R, Abdelwahab SI, Ibrahim MY, Ali LZ. A review of traditional uses, phytochemical and pharmacological aspects of selected members of *Clausena* genus (Rutaceae). *J Med Plants Res*. 2012;6(38):5107–5118. <https://doi.org/10.5897/JMPR12.317>
  87. Ojewole JA. Hypoglycaemic effect of *Clausena anisata* (Willd.) Hook methanolic root extract in rats. *J Ethnopharmacol*. 2002;81(2):231–237. [https://doi.org/10.1016/s0378-8741\(02\)00085-5](https://doi.org/10.1016/s0378-8741(02)00085-5)
  88. Uwaifo AO. The mutagenicities of seven coumarin derivatives and a furan derivative (nimbolide) isolated from three medicinal plants. *J Toxicol Environ Health*. 1984;13(4–6):521–530. <https://doi.org/10.1080/15287398409530517>
  89. Nguta JM, Mbaria JM, Gathumbi PK, Gakuya D, Kabasa JD, Kiama SG. Ethnodiagnostic skills of the Digo community for malaria: a lead to traditional bioprospecting. *Front Pharmacol*. 2011;2:30. <https://doi.org/10.3389/fphar.2011.00030>
  90. Beentje HJ, Joy A, Dhan B, et al. *Kenya Trees, Shrubs, and Lianas*. Nairobi, Kenya: National Museums of Kenya; 1994. ISBN:9966986103
  91. Duncan AC, Jäger AK, van Staden J. Screening of Zulu medicinal plants for angiotensin converting enzyme (ACE) inhibitors. *J Ethnopharmacol*. 1999;68(1–3):63–70. [https://doi.org/10.1016/s0378-8741\(99\)00097-5](https://doi.org/10.1016/s0378-8741(99)00097-5)
  92. Ragunathan M, Weldegerima B. Medico ethno botany; a study on the Amhara ethnic group of Gondar District of North Gondar Zone Ethiopia. *J Nat Remed*. 2007;7(2):200–206. <http://scholar.info/index.php/jnr/article/view/28496>
  93. Iyamah PC, Idu M. Ethnomedicinal survey of plants used in the treatment of malaria in Southern Nigeria. *J Ethnopharmacol*. 2015;173:287–302. <https://doi.org/10.1016/j.jep.2015.07.008>
  94. Lindsay RS. *Medicinal Plants of Marakwet, Kenya*. Royal Botanic Gardens (Kew): Herbarium; 1978. <https://wellcomecollection.org/works/fu3evjnp>
  95. Adesina SK, Ette EI. The isolation and identification of anticonvulsant agents from *Clausena anisata* and *Afraegle paniculata*. *Fitoterapia*. 1982;53(3):63. [https://scholar.google.com/scholar\\_lookup?title=The%20isolation%20and%20identification%20of%20anticonvulsant%20agents%20from%20Clausena%20anisata%20and%20Afraegle%20paniculata&publication\\_year=1982&author=S.K%20Adesina&author=E.I%20Ette](https://scholar.google.com/scholar_lookup?title=The%20isolation%20and%20identification%20of%20anticonvulsant%20agents%20from%20Clausena%20anisata%20and%20Afraegle%20paniculata&publication_year=1982&author=S.K%20Adesina&author=E.I%20Ette)
  96. Makanju OOA. Behavioral and anticonvulsant effects of an aqueous extract from the roots of *Clausena anisata* (Rutaceae). *Int J Crude Drug Res*. 1983;21(1):29–32. <https://doi.org/10.3109/13880208309070606>
  97. Adesina SK. Studies on some plants used as anticonvulsants in Amerindian and African traditional medicine. *Fitoterapia*. 1982;53:147–162. <https://cir.nii.ac.jp/crid/1573668925071598208?lang=en>
  98. Malweyi SL. Evaluation of prioritized medicinal plants for their bioactivity in Kaimosi area of Nandi and Vihiga Counties of Kenya. MSc Thesis, University Of Eldoret, Kenya; 2013. <http://41.89.164.27/handle/123456789/817>
  99. Gachathi FN. Kikuyu botanical dictionary of plant names and uses. *Tropical Botany*; 1989. <https://agris.fao.org/agris-search/search.do?recordID=US2013>

00290007

100. Kokwaro JO. *Medicinal Plants of East Africa (3rd ed)*. Nairobi, Kenya:University of Nairobi Press;2009. <https://xs2.dailyheadlines.cc/books?id=msyHLY0dhPwC&printsec=frontcover&hl=zh-CN>
101. Moshi MJ, van den Beukel CJP, Hamza OJM, et al. Brine shrimp toxicity evaluation of some tanzanian plants used traditionally for the treatment of fungal infections. *Afr J Complement Altern Med*. 2006;4(2):219–225. <https://doi.org/10.4314/ajtcam.v4i2.31211>
102. Lewis WH. *Elvin-Lewis MP (1977) Medical Botany*. New York;John Wiley and Sons:1977:1–515. ISBN:0471861340
103. Hutchings A, Scoth AH, Lewis G, Cunningham A. *Clausena anisata (Wild)*. Hook. F. ex Benth, *Zulu Medicinal Plants: An Inventory*. Pietermaritzburg, South Africa:University of Natal Press;1996:153–154. ISBN:0869809237
104. York T, van Vuuren SF, deWet H. An antimicrobial evaluation of plants used for the treatment of respiratory infections in rural Maputaland, KwaZulu-Natal, South Africa. *J Ethnopharmacol*. 2012;144(1):118–127. <https://doi.org/10.1016/j.jep.2012.08.038>
105. Watt JM, Breyer-Brandwijk MG. *The Medicinal and Poisonous Plants of Southern and Eastern Africa (2nd ed)*. Livingstone, London;1962. <https://agris.fao.org/agris-search/search.do?recordID=XF2016015447>
106. Kokwaro JO. *Medicinal Plants of East Africa*. Nairobi, Kenya:East African Literature Bureau;1976. <http://41.204.161.209/handle/11295/34820>
107. Palgrave KC. *Trees of Southern Africa (3rd ed)*. Struik, Cape Town, South Africa: Pooley E.; 1998. *A Field Guide to Wild Flowers-KwaZulu-Natal and the Eastern Region*. Durban, South Africa: Natal Flora Publications Trust;2002. ISBN:9781920572754
108. Innocent E, Hassanali A. Constituents of Eos from three plant species used in traditional medicine and insect control in tanzania. *J Herbs Spices Med Plants*. 2015;21(3):219–229. <https://doi.org/10.1080/10496475.2014.949997>
109. Yineger H, Yewhalaw D, Teketay D. Ethnomedicinal plant knowledge and practice of the Oromo ethnic group in southwestern Ethiopia. *J Ethnobiol Ethnomed*. 2008;4:11. <https://doi.org/10.1186/1746-4269-4-11>
110. Hamill FA, Apio S, Mubiru NK, et al. Traditional herbal drugs of Southern Uganda, II: literature analysis and antimicrobial assays. *J Ethnopharmacol*. 2003;84(1):57–78. [https://doi.org/10.1016/s0378-8741\(02\)00289-1](https://doi.org/10.1016/s0378-8741(02)00289-1)
111. Gakuya DW, Itonga SM, Mbaria JM, Muthee JK, Musau JK. Ethnobotanical survey of biopesticides and other medicinal plants traditionally used in Meru central district of Kenya. *J Ethnopharmacol*. 2013;145(2):547–553. <https://doi.org/10.1016/j.jep.2012.11.028>
112. Kigen G, Kamuren Z, Njiru E, Wanjohi B, Kipkore W. Ethnomedicinal survey of the plants used by traditional healers in Narok county, Kenya. *Evid-Based Complement Alternat Med*. 2019;2019:8976937. <https://doi.org/10.1155/2019/8976937>
113. Bryant AT (Alfred T). *Zulu Medicine and Medicine-Men*. C. Struik, Cape Town;1966. [https://ehrafworldcultures.yale.edu/ehrafe/citation.do?method=citation&forward=browseCulturesFullContext&col=collection\(%27/eHRAF/ethnography/Africa/FX20%27\)&docId=fx20-009&tocOffsetId=tocPubInfoP](https://ehrafworldcultures.yale.edu/ehrafe/citation.do?method=citation&forward=browseCulturesFullContext&col=collection(%27/eHRAF/ethnography/Africa/FX20%27)&docId=fx20-009&tocOffsetId=tocPubInfoP)
114. Mhlongo LS, Van Wyk BE. Zulu medicinal ethnobotany: new records from the Amandawe area of KwaZulu-Natal, South Africa. *South Afr J Bot*. 2019;122:266–290. <https://doi.org/10.1016/j.sajb.2019.02.012>
115. Sikolia SF. Medicinal plants of Kakamega forests and their consistency applications: opportunities and challenges to-date. *Inst Rep*. 2018;13:48–54. <http://41.204.161.159/handle/123456789/2991>
116. Kaingu CK, Oduma JA, Kanui TI. Practices of traditional birth attendants in Machakos District, Kenya. *J Ethnopharmacol*. 2011;137(1):495–502. <https://doi.org/10.1016/j.jep.2011.05.044>
117. Ole-Miaron JO, Mapenay M. Maasai ethnoveterinary medicine: an avenue of bioprospecting for anthelmintics. *Kenya Veterinarian*. 2004;27:3–4. <https://doi.org/10.4314/kenvet.v27i1.39549>
118. Pavela R, Maggi F, Lupidi G, et al. *Clausena anisata* and *Dysphania ambrosioides* Eos: from ethno-medicine to modern uses as effective insecticides. *Environ Sci Pollut Res Int*. 2017;25(11):10493–10503. <https://doi.org/10.1007/s11356-017-0267-9>
119. Chavunduka DM. Plants regarded by Africans as being of medicinal value to animals. *Rhod Vet J*. 1976;7:6–12. <https://www.semanticscholar.org/paper/Plants-regarded-by-Africans-as-being-of-medicinal-Chavunduka/24a808ca475bb95bf84cec60ff17c70ca453ae6c>
120. Mukandiwa L, Eloff JN, Sibanda DR, Naidoo V. An acetone extract of *Clausena anisata* may be a potential control agent for flies encountered in cutaneous myiasis. *Onderstepoort J Vet Res*. 2016;83(1):a1045. <https://doi.org/10.4102/ojvr.v83i1.1045>
121. Ole-Miaron J. The Maasai ethnodagnostic skill of livestock diseases: a lead to traditional bioprospecting. *J Ethnopharmacol*. 2003;84(1):79–83. [https://doi.org/10.1016/s0378-8741\(02\)00283-0](https://doi.org/10.1016/s0378-8741(02)00283-0)
122. Karunamoorthi K, Husen E. Knowledge and self-reported practice of the local inhabitants on traditional insect repellent plants in Western Hararghe zone, Ethiopia. *J Ethnopharmacol*. 2012;141(1):212–219. <https://doi.org/10.1016/j.jep.2012.02.022>
123. Tapondjou AL, Bouda H, Fontem DA, Zapfack L, Lontsi D, Sondengam BL (2000) Local plants used for traditional stored product protection in the Menoua division of the Western highlands of Cameroon. IOBC/WPRS Working Group Integrated Protection in Stored Products. Proceedings of the meeting at Berlin, Germany, 22–24. 1999;23:73–77. <https://www.cabdirect.org/cabdirect/abstract/20023077012>
124. Chhabra SC, Mahunnah RL, Mshiu EN. Plants used in traditional medicine in Eastern Tanzania. II. Angiosperms (capparidaceae to ebenaceae). *J Ethnopharmacol*. 1989;25(3):339–359. [https://doi.org/10.1016/0378-8741\(89\)90038-x](https://doi.org/10.1016/0378-8741(89)90038-x)
125. Omara T, Kiprop AK, Kosgei VJ. *Albizia coriaria* Welw ex Oliver: a review of its ethnobotany, phytochemistry and ethnopharmacology. *Adv Tradit Med*. 2021. <https://doi.org/10.1007/s13596-021-00600-8>
126. Agbor AM, Fotsing KP, Dountio SH, Fokunang NC, Naidoo S. Phytochemical investigation and anti-microbial activity of *Clausena anisata* on cariogenic and periodontopathic bacteria. *Eur J Med Plants*. 2020;31(9):11–25. <https://doi.org/10.9734/ejmp/2020/v31i1930345>
127. Lawal IO, Galadima M, Ogunbamowo PO. Isolation of bioactive compounds of *Clausena anisata* (Willd.) Hook. growing in South Africa by liquid chromatography–mass spectroscopy profiling, and their antibacterial activities. *J Med Plants Econ Dev*. 2018;2(1):a22. <https://hdl.handle.net/10520/EJC-1053352c82>
128. Kumatia EK, Ayertey F, Appiah AA, Ocloo A. Analgesic, anti-inflammatory and anti-oxidant activities of *Clausena anisata* leaf and stem extracts and quantitative and qualitative phytochemical analysis of their contents. *Der Pharmazie*. 2018;11(4):16–23. [https://www.researchgate.net/publication/333520620\\_Analgesic\\_Anti-Inflammatory\\_and\\_Anti-Oxidant\\_Activities\\_of\\_Clausena\\_anisata\\_Leaf\\_and\\_Stem\\_Extracts\\_and\\_Quantitative\\_and\\_Qualitative](https://www.researchgate.net/publication/333520620_Analgesic_Anti-Inflammatory_and_Anti-Oxidant_Activities_of_Clausena_anisata_Leaf_and_Stem_Extracts_and_Quantitative_and_Qualitative)



## Phytochemical Analysis of their Contents

129. Arsia Y, Begum TN, Lamya N. Study of invitro antioxidant, antidiabetic and antibacterial activity of *Clausena anisata* (Willd.) Hook (Rutaceae) leaf extracts. *Int J Res Pharmacol Pharmacother*. 2014;3(2):112–125. <https://vdocuments.net/study-of-invitr-antioxidant-antidiabetic-and-antibacterial-activity-of-clausena-anisata-willd-hook-rutaceae-leaf-extracts.html>
130. Kuete V, Tankeo SB, Saeed MEM, Wiench B, Tane P, Efferth T. Cytotoxicity and modes of action of five Cameroonian medicinal plants against multi-factorial drug resistance of tumor cells. *J Ethnopharmacol*. 2014;153(1):207–219. <https://doi.org/10.1016/j.jep.2014.02.025>
131. Okokon JE, Etebong EO, Udobang JA, Essien GE. Antiplasmodial and analgesic activities of *Clausena anisata*. *Asian Pac J Trop Med*. 2012;5(3):214–219. [https://doi.org/10.1016/S1995-7645\(12\)60027-3](https://doi.org/10.1016/S1995-7645(12)60027-3)
132. Kenechukwu FC, Mbah CJ, Momoh MA, Chime SA, Umeyor CE, Ogbonna JDN. Pharmacological justification for the ethnomedical use of *Clausena anisata* root-bark extract in the management of epilepsy. *J Appl Pharmaceut Sci*. 2012;2(9):036–040. <http://dx.doi.org/10.7324/JAPS.2012.2907>
133. Asante KI, Amuzu N, Donkor A, Annan K. Screening of some Ghanaian medicinal plants for phenolic compounds and radical scavenging activities. *Pharmacog J*. 2009;1(3):201–206. <http://ugspace.ug.edu.gh/handle/123456789/29385?show=full>
134. Adebayo SA, Dzoyem JP, Shai LJ, Eloff JN. The anti-inflammatory and antioxidant activity of 25 plant species used traditionally to treat pain in southern African. *BMC Compl Alternat Med*. 2015;15:159. <https://doi.org/10.1186/s12906-015-0669-5>
135. Nyoni M, Muzemu S, Chinheya C, Mushayabasa T, Ncube B. Screening indigenous nematocidal plants in Zimbabwe against *Meloidogyne javanica*. 1st International research conference on driving socio-economic development through value addition and sustainable use of resources, 14th-16th July 2015, Elephant Hills Resort, Victoria Falls, Zimbabwe. <http://hdl.handle.net/11408/1319>
136. Amoo SO, Aremu AO, Moyo M, Van Staden J. Antioxidant and acetylcholinesterase-inhibitory properties of long-term stored medicinal plants. *BMC Compl Alternat Med*. 2012;12:87. <https://doi.org/10.1186/1472-6882-12-87>
137. Otang WM, Grierson DS, Afolayan AJ. A survey of plants responsible for causing irritant contact dermatitis in the Amathole district, eastern cape, South Africa. *J Ethnopharmacol*. 2014;157:274–284. <https://doi.org/10.1016/j.jep.2014.10.002>
138. Ito C, Katsuno S, Itoigawa M, Ruangrunsi N, Furukawa H. Structures of Clausamine-A, -B, -C, three novel carbazole alkaloids from *Clausena anisata*. *Chem Pharmaceut Bulletin*. 1998;46(2):344–346. <https://doi.org/10.1248/cpb.46.344>
139. Ito C, Katsuno S, Itoigawa M, et al. New carbazole alkaloids from *Clausena anisata* with antitumor promoting activity. *J Nat Prod*. 2000;63:125–128. <https://doi.org/10.1021/np990285x>
140. Ito C, Itoigawa M, Aizawa K, Yoshida K, Ruangrunsi N, Hiroshi F.  $\gamma$ -Lactone Carbazoles from *Clausena anisata*. *J Nat Prod*. 2009;72(6):1202–1204. <https://doi.org/10.1021/np800735m>
141. Songue JL, Kouam DE, Mpondo TN, White RL. Chemical constituents from stem bark and roots of *Clausena anisata*. *Molecules*. 2012;17(11):13673–13686. <https://doi.org/10.3390/molecules171113673>
142. Tatsimo SJN, Tamokou JDD, Lamsho'ft M, et al. LC-MS guided isolation of antibacterial and cytotoxic constituents from *Clausena anisata*. *Med Chem Res*. 2015;24:1468–1479. <https://doi.org/10.1007/s00044-014-1233-4>
143. Ngadjui BT, Ayafor JF, Sondengam BL, Connolly JD. Quinolone and carbazole alkaloids from *Clausena anisata*. *Phytochemistry*. 1989;28(5):1517–1519. [https://doi.org/10.1016/S0031-9422\(00\)97778-1](https://doi.org/10.1016/S0031-9422(00)97778-1)
144. Chakraborty A, Chowdhury BK, Bhattacharyya P. Clausenol and clausenine-two carbazole alkaloids from *Clausena anisata*. *Phytochemistry*. 1995;40(1):295–298. [https://doi.org/10.1016/0031-9422\(95\)00047-b](https://doi.org/10.1016/0031-9422(95)00047-b)
145. Kumatia EK, Annan K, Dickson RA, et al. Anti-inflammatory and analgesic effects in rodent models of ethanol extract of *Clausena anisata* roots and their chemical constituents. *Nat Prod Comm*. 2017;12(1):67–72. <https://pubmed.ncbi.nlm.nih.gov/30549828/>
146. Okorie DA. A new carbazole alkaloid and coumarins from roots of *Clausena anisata*. *Phytochemistry*. 1975;14(12):2720–2721. <https://doi.org/10.1016/0031-9422%2875%2985267-8>
147. Mester I, Bergenthal D, Reisch J. Inhaltsstoffe aus *Clausena anisata* (Willd.) Oliv. (Rutaceae), III 13C-NMR-Spektren des Mupamins, des Carbazols und einiger Carbazol-Derivate/Constituents of *Clausena anisata* (Willd.) Oliv. (Rutaceae), III 13C NMR Spectra of Mupamine, Carbazole and Some Carbazole Derivatives. *Zeitschrift Für Naturforschung B*. 1979;34(4):650–652. <https://doi.org/10.1515/znb-1979-0427>
148. Mester I, Szendrei K, Reisch J. Constituents of *Clausena anisata* (Willd.) Oliv. (Rutaceae), I. coumarins from the root bark. *Planta Med*. 1977;32(1):81–85. <https://doi.org/10.1055/s-0028-1097563>
149. Olorunsogo OO, Emerole GO, Malomo SO, Thabrew MI. Inhibition of mitochondrial respiration by chalepin, a naturally occurring furocoumarin. *Toxicol Lett*. 1983;15(4):323–327. [https://doi.org/10.1016/0378-4274\(83\)90151-0](https://doi.org/10.1016/0378-4274(83)90151-0)
150. Ayafor JF, Ngadjui B, Sondengam BL. PA12 novel bioactive coumarins from *Clausena anisata*. *Int Symp Chem Nat Prod*. 1988. [https://doi.org/10.24496/intnaturalprod.1988.0\\_90](https://doi.org/10.24496/intnaturalprod.1988.0_90)
151. Lakshmi V, Prakash D, Raj K, Kapil RS, Popli SP. Monoterpenoid furanocoumarin lactones from *Clausena anisata*. *Phytochemistry*. 1984;23(11):2629–2631. [https://doi.org/10.1016/S0031-9422\(00\)84114-X](https://doi.org/10.1016/S0031-9422(00)84114-X)
152. Ngadjui BT, Mouncherou SM, Ayafor JF, Sondengam BL, Tillequint F. Geranyl coumarins from *Clausena anisata*. *Phytochemistry*. 1991;30(8):2809–2811. [https://doi.org/10.1016/0031-9422\(91\)85155-S](https://doi.org/10.1016/0031-9422(91)85155-S)
153. Ngadjui BT, Ayafor JF, Sondengam BL, Connolly JD. Coumarins from *Clausena anisata*. *Phytochemistry*. 1989;28(2):585–589. [https://doi.org/10.1016/0031-9422\(89\)80056-1](https://doi.org/10.1016/0031-9422(89)80056-1)
154. Mukandiwa L, Ahmed A, Eloff JN, Naidoo V. Isolation of seselin from *Clausena anisata* (Rutaceae) leaves and its effects on the feeding and development of *Lucilia cuprina* larvae may explain its use in ethnoveterinary medicine. *J Ethnopharmacol*. 2013;150(3):886–891. <https://doi.org/10.1016/j.jep.2013.09.037>
155. Ngadjui BT, Ayafor JF, Sondengam BL, Connolly JD. Limonoids from *Clausena anisata*. *J Nat Prod*. 1989;52(4):832–836. <https://doi.org/https://doi.org/10.1021/np50064a027>
156. Jayaraman M, Senthilkumar A, Adaikala Raj G, Venkatesalu V. Isolation of mosquito larvicidal molecule from the leaves of *Clausena anisata*. *J Exp Sci*. 2015;6:12–16. <https://doi.org/10.19071/JES.2015.V6.2927>
157. De Canha MN, Kishore N, Kumar V, et al. The potential of *Clausena anisata* (Willd.) Hook.f. ex Benth against *Propionibacterium acnes*. *South Afr J Bot*. 2018;119:410–419. <https://doi.org/10.1016/j.sajb.2018.09.019>
158. Guenther E. *The EOs, vol I*. New York, van Nostrand Company Inc;1960:372. <https://doi.org/10.1002/jps.3030370518>
159. Simiyu S. Bioevaluation and chemical identification of insecticidal/repellent plants and derivatives from central region

- of Kenya. Masters Thesis, Kenyatta University, Kenya;2003.  
<https://ir-library.ku.ac.ke/handle/123456789/2475?show=full>
160. Ekundayo O, Oguntimehin BO, Hammerschmidt FJ. Constituents of the essential oil of *Clausena anisata* leaves. *Planta Med.* 1986;52(6):505–506.  
<https://doi.org/10.1055/s-2007-969274>
  161. Mollong E, Nuto Y, Bawa R, Kassene DB, Gomina M. Acaricidal effect of essential oil of *Clausena anisata* (Rutaceae) on larvae of three tick species: *amblyomma variegatum*, *Rhipicephalus* (*Boophilus*) *decoloratus* and *Rhipicephalus* (*Boophilus*) *microplus*. *J Entomol Nematol.* 2018;10(4):23–32.  
<https://doi.org/10.5897/JEN2018.0203>
  162. Gundidza M, Chinyangaya F, Chagonda L, Depooter HL, Mavi S. Phytoconstituents and antimicrobial activity of the leaf essential oil of *Clausena anisata* (Willd.). J.D. Handbook ex. Benth. *Flavour Fragr J.* 1994;9(6):299–303.  
<https://doi.org/10.1002/ffj.2730090604>
  163. Essilfie MK, Oteng-Yeboah AA. Morphometric studies of *Clausena anisata* (Willd.) Hook. f. ex. Benth. in Coastal Savanna Zone of Ghana. *West Afr J Appl Ecol.* 2013;21(1):1–13.  
<https://www.ajol.info/index.php/wajae/article/view/94726>
  164. Machumi F, Innocent E, Yand P. Seasonal impacts on antifungal activity and chemical composition of extracts from medicinal plants *Turraea holstii* and *Clausena anisata*. *J Med Plants Res.* 2021;15(4):133–139.  
<https://doi.org/10.5897/JMPR2020.7056>
  165. Lawal IO, Ajay H, Dairo P, Ogunbamowo PO. Comparative studies of Eos of *Clausena anisata* (Hook) using solvent-free microwave extraction and hydrodistillation methods. *J Med Plants Econ Dev.* 2019;3(1):a59.  
<https://hdl.handle.net/10520/EJC-1719ea88b6>
  166. Lawal IO. Chemical composition, cytotoxicity and anti-mycobacterium activities of *Clausena anisata* essential oils. *J Med Plants Econ Dev.* 2020;4:a90.  
<https://doi.org/10.4102/jomped.v4.i1.90>
  167. Goudoum A, Tinkeu LSN, Ngamo TL, Ngassoum MB, Mbofung CM. Antioxidant activities of Eos of *Clausena anisata* (Rutaceae) and *Plectranthus glandulosus* (Labiatae), plants used against stored grain insects in North Cameroon. *Int J Biol Chem Sci.* 2009;3(3):567–577.  
<https://doi.org/10.4314/ijbcs.v3i3.45338>
  168. Yaouba A, Tatsadjieu LN, Dongmo PMJ, et al. Evaluation of *Clausena anisata* Eos from Cameroon for controlling food spoilage fungi and its potential use as an antiradical agent. *Nat Prod Comm.* 2011;6(9):1367–1371.  
<https://doi.org/10.1177%2F1934578X1100600937>
  169. Arsia TY, Nargis BT, Muhammad IMH, Shilu M, Archunan G, Ishtia Q. Green synthesis, Antioxidant potential and hypoglycemic effect of silver nanoparticles using ethanolic leaf extract of *Clausena anisata* (Willd.) Hook. F. Ex Benth. of Rutaceae. *Pharmacog J.* 2016;8(6):565–575.  
<http://dx.doi.org/10.5530/pj.2016.6.8>
  170. Senatore F, Oliviero F, Scandolera E, et al. Chemical composition, antimicrobial and antioxidant activities of anethole-rich oil from leaves of selected varieties of fennel [*Foeniculum vulgare* Mill. ssp. *vulgare* var. *azoricum* (Mill.) Thell]. *Fitoterapia.* 2013;90:214–219.  
<https://doi.org/10.1016/j.fitote.2013.07.021>
  171. Osei-Safo D, Addae-Mensah I, Garneau F, Koumaglo H. A comparative study of the antimicrobial activity of the leaf Eos of chemo-varieties of *Clausena anisata* (Willd.) Hook. f. ex Benth. *Indust Crops Prod.* 2010;32(3):634–638.  
<https://doi.org/10.1016/j.indcrop.2010.07.016>
  172. Sessou P, Yaovi BA, Yovo M, et al. Phytochemistry and antibacterial activity of plants extracts compared with two commercial antibiotics against *E. coli* responsible for avian colibacillosis in Benin. *Int J Phytomed.* 2018;10(3):168–174.  
<https://doi.org/10.5138/09750185.2259>
  173. Adamu M, Naidoo V, Eloff JN. Some southern African plant species used to treat helminth infections in ethnoveterinary medicine have excellent antifungal activities. *BMC Compl Alternat Med.* 2012;12:213.  
<https://doi.org/10.1186/1472-6882-12-213>
  174. Makirita WE, Chauka LJ, Chacha M. Antimicrobial and cytotoxicity activity of *Clausena anisata*, *acokanthera shemperii* and *olea europaea* growing in Tanzania. *Eur J Med Plants.* 2016;14:1–9.  
<https://dSPACE.nm-aist.ac.tz/handle/20.500.12479/1124>
  175. Mutuku AM. Antimicrobial activity of *warburgia ugandensis*, *euclea divinorum*, *rhus vulgaris*, *ocimum gratissimum* and *Clausena anisata* against oral pathogenic microorganisms. MSc Thesis, University of Eldoret, Eldoret, Kenya;2018.  
<http://41.89.164.27/handle/123456789/369>
  176. Amoo SO, Aremu AO, Moyo M, Van Staden J. Assessment of long-term storage on antimicrobial and cyclooxygenase inhibitory properties of South African medicinal Plants. *Phyther Res.* 2013;27(7):1029–1035.  
<https://doi.org/10.1002/ptr.4830>
  177. Thibane VS, Ndhlala AR, Finnie JF, Van Staden J. Modulation of the enzyme activity of secretory phospholipase A<sub>2</sub>, lipoxygenase and cyclooxygenase involved in inflammation and disease by extracts from some medicinal plants used for skincare and beauty. *South Afri J Bot.* 2019;120:198–203.  
<https://doi.org/10.1016/j.sajb.2018.06.001>
  178. Thibane VS, Ndhlala AR, Finnie JF, Van Staden J. Cosmeceutical efficiency by some plant extracts used traditionally for skin care in inhibiting tyrosinase activity in a human epidermal melanocyte (HEM) cell line. *South Afri J Bot.* 2019;126:256–260.  
<https://doi.org/10.1016/j.sajb.2019.06.031>
  179. Ohashi M, Amoa-Bosompem M, Kwofie KD, et al. In vitro antiprotozoan activity and mechanisms of action of selected Ghanaian medicinal plants against *Trypanosoma*, *Leishmania* and *Plasmodium* parasites. *Phyther Res.* 2018;32(8):1617–1630.  
<https://doi.org/10.1002/ptr.6093>
  180. Nibret E, Wink M. Trypanocidal and cytotoxic effects of 30 Ethiopian medicinal plants. *Zeitschrift Fur Naturforschung C.* 2011;66(11–12):541–546.  
<https://doi.org/10.1515/znc-2011-11-1202>
  181. Van den Bout-van den Beukel CJP, Hamza OJM, Moshi MJ, et al. Evaluation of cytotoxic, genotoxic and CYP450 enzymatic competition effects of tanzanian plant extracts traditionally used for treatment of fungal infections. *Basic Clin Pharmacol Toxicol.* 2008;102(6):515–526.  
<https://doi.org/10.1111/j.1742-7843.2008.00225.x>
  182. Lin W, Wang Y, Lin S, et al. Induction of cell cycle arrest by the carbazole alkaloid Clausoline-I from *Clausena vestita* DD. Tao via inhibition of the PKC $\delta$  phosphorylation. *Eur J Med Chem.* 2012;47:214–220.  
<https://doi.org/10.1016/j.ejmech.2011.10.047>
  183. Cui CB, Yan SY, Cai B, Yao XS. Carbazole alkaloids as new cell cycle inhibitor and apoptosis inducers from *Clausena dunniana* levl. *J Asian Nat Prod Res.* 2002;4(4):233–241.  
<https://doi.org/10.1080/1028602021000049041>
  184. Clarkson C, Maharaj VJ, Crouch NR, et al. In vitro antiplasmodial activity of medicinal plants native to or naturalised in South Africa. *J Ethnopharmacol.* 2004;92(2–3):177–191.  
<https://doi.org/10.1016/j.jep.2004.02.011>
  185. Weenen H, Nkunya MH, Bray DH, Mwasumbi LB, Kinabo LS, Kilimali VA. Antimalarial activity of Tanzanian medicinal plants. *Planta Med.* 1990;56(4):368–370.  
<https://doi.org/10.1055/s-2006-960984>
  186. McGaw LJ, Ja`ger AK, van Staden J. Antibacterial, anthelmintic and anti-amoebic activity in South African medicinal plants. *J Ethnopharmacol.* 2000;72(1–2):247–263.  
[https://doi.org/10.1016/s0378-8741\(00\)00269-5](https://doi.org/10.1016/s0378-8741(00)00269-5)

187. Christensen CB, Soelberg J, Stensvold CR, Jäger AK. Activity of medicinal plants from Ghana against the parasitic gut protist *Blastocystis*. *J Ethnopharmacol*. 2015;174:569–575. <https://doi.org/10.1016/j.jep.2015.03.006>
188. Williams AR, Soelberg J, Jäger AK. Anthelmintic properties of traditional African and Caribbean medicinal plants: identification of extracts with potent activity against *Ascaris suum* in vitro. *Parasite*. 2016;23:24. <https://doi.org/10.1051/parasite/2016024>
189. Kamte SLN, Ranjbarian F, Campagnaro GD, et al. Trypanosoma brucei Inhibition by Eos from medicinal and aromatic plants traditionally used in cameroon (*Azadirachta indica*, *Aframomum melegueta*, *Aframomum daniellii*, *Clausena anisata*, *Dichrostachys cinerea* and *Echinops giganteus*). *Int J Environ Res Public Health*. 2017;14(7):737. <https://doi.org/10.3390/ijerph14070737>
190. Mogale MA, Mkhombo MH, Lebelo SL, et al. The effects of *Clausena anisata* (Willd) Hook leaf extracts on selected diabetic related metabolizing enzymes. *J Med Plants Res*. 2012;6(25):4200–4207. <https://doi.org/10.5897/JMPR12.208>
191. Peter EL, Nagendrappa PB, Hilonga S, et al. Pharmacological reflection of plants traditionally used to manage diabetes mellitus in Tanzania. *J Ethnopharmacol*. 2021;269:113715. <https://doi.org/10.1016/j.jep.2020.113715>
192. Arsia TY, Nargis BT, Syed JH, Shilu M, Muhammad IMH. In vivo analysis of *Clausena anisata* (Willd.) Hook. f. ex Benth crude and SNP extracts against antioxidant and hypoglycemic activity. *Int J Pharmaceut Clin Res*. 2018;10:28–39. <http://hdl.handle.net/10576/11314>
193. Risa J, Risa A, Adersen A, et al. Screening of plants used in southern Africa for epilepsy and convulsions in the GABA<sub>A</sub>-benzodiazepine receptor assay. *J Ethnopharmacol*. 2004;93(2–3):177–182. <https://doi.org/10.1016/j.jep.2004.01.021>
194. Mbah C, Kenechukwu F. Preliminary screening of ethanolic extracts of *Clausena anisata* for anticonvulsant activity. *Bio-Res*. 2009;7(2):548–550. <https://doi.org/10.4314/br.v7i2.56601>
195. Gilani AH, Shaheen E, Saeed SA, et al. Hypotensive action of coumarin glycosides from *Daucus carota*. *Phytomedicine*. 2000;7(5):423–426. [https://doi.org/10.1016/s0944-7113\(00\)80064-1](https://doi.org/10.1016/s0944-7113(00)80064-1)
196. Oussou KR, Kanko C, Philomène KA, Guessennnd KN, Boti JB. Chemical composition and inhibitory effects of Eos of two Rutaceae of Côte d'Ivoire: *Clausena anisata* and *Citrus aurantifolia*. *Int J Eng Appl Sci*. 2017;4(12):8–12. <https://www.neliti.com/publications/257319/chemical-composition-and-inhibitory-effects-of-essential-oils-of-two-rutaceae-of-cote>
197. Dongdem FA. Effects of extracts and hydrodistillates of *Clausena anisata* (Willd.) ex. Benth. and *Hyptis spicigera* Lam. on *Musca domestica* L. and *Periplaneta americana* (L.). MPhil Thesis, University of Ghana, Ghana;1997. <http://ugspace.ug.edu.gh/handle/123456789/6216>
198. Tapondjou AL, Adler C, Bouda H, Reichmuth C. Ability of products derived from the leaves of *Clausena anisata* to protect stored legumes from attack by *Callosobruchus maculatus* and *C. chinensis* (Coleoptera, Bruchidae). *Integr Protect Stored Prod IOBC Bullet*. 2002;25(3):153–164. [http://www.iobc-wprs.org/pub/bulletins/iobc-wprs\\_bulletin\\_20\\_02\\_25\\_03.pdf#page=177](http://www.iobc-wprs.org/pub/bulletins/iobc-wprs_bulletin_20_02_25_03.pdf#page=177)
199. Boeke SJ, Baumgart IR, van Loon JJA, van Huis A, Dicke M, Kossou DK. Toxicity and repellence of African plants traditionally used for the protection of stored cowpea against *Callosobruchus maculatus*. *J Stored Prod Res*. 2004;40(4):423–438. [https://doi.org/10.1016/S0022-474X\(03\)00046-8](https://doi.org/10.1016/S0022-474X(03)00046-8)
200. Boeke SJ, Barnaud C, van Loon JJA, Kossou DK, van Huis A, Dicke M. Efficacy of plant extracts against the cowpea beetle, *Callosobruchus maculatus*. *Int J Pest Manage*. 2004;50(4):251–258. <https://doi.org/10.1080/09670870410001702206>
201. Ndomo AF, Ngamo LT, Tapondjou LA, Tchouanguép FM, Hance T. Insecticidal effects of the powdery formulation based on clay and EO from the leaves of *Clausena anisata* (Willd.) J. D. Hook ex. Benth. (Rutaceae) against *Acanthoscelides obtectus* (Say) (Coleoptera: Bruchidae). *J Pest Sci*. 2008;81(227):227–234. <https://doi.org/10.1007/s10340-008-0211-3>
202. Govindarajan M. Chemical composition and larvicidal activity of leaf essential oil from *Clausena anisata* (Willd.) Hook. f. ex. Benth (Rutaceae) against three mosquito species. *Asian Pac J Trop Med*. 2010;3(11):874–877. [https://doi.org/10.1016/S1995-7645\(10\)60210-1](https://doi.org/10.1016/S1995-7645(10)60210-1)
203. European Medicines Agency, Evaluation of medicines for human use: Publicstatement on the use of herbal medicinal products containing estragol. Committee on herbal medicinal products. Doc Ref: EMEA/HMPC/137212/2005, London;2005. [https://www.ema.europa.eu/en/documents/regulatory-procedural-guideline/compassionate-use-centralised-medicinal-products-according-article-83-regulation-ec-no-726/2004\\_en.pdf](https://www.ema.europa.eu/en/documents/regulatory-procedural-guideline/compassionate-use-centralised-medicinal-products-according-article-83-regulation-ec-no-726/2004_en.pdf)
204. Goudoum A, Ngamo Tinkeu LS, Ngamo TL, Ngassoum MB, Mbofung CM. Biodegradation of insecticidal compounds of *Clausena anisata* and *Plectranthus glandulosus* Eos applied as protectant on stored grains. *Agric J*. 2012;7(2):165–171. <https://doi.org/10.3923/AJ.2012.165.171>
205. Goudoum A, Ngamo TL, Ngassoum MB, Mbofung CM. Variation in technological and nutritional parameters of maize treated with Eos during storage. *Afr J Food Sci*. 2012;6(2):34–40. <https://doi.org/10.5897/AJFS11.198>
206. Afshar FH, Maggi F, Iannarelli R, Cianfaglione K, Isman MB. Comparative toxicity of *Helosciadium nodiflorum* EOs and combinations of their main constituents against the cabbage looper, *Trichoplusia ni* (Lepidoptera). *Indust Crops Prod*. 2017;98:46–52. <https://doi.org/10.1016/j.indcrop.2017.01.004>
207. Adjalian E, Sessou P, Alitonou GA, et al. Ovicide and larvicide activities of the Eos extracted from *polyalthia longifolia* and *Clausena anisata* on *Sitotroga cerealella*. *Online J Biol Sci*. 2017;17(3):193–200. <https://doi.org/10.3844/ojbsci.2017.193.200>
208. Aurelie FDG, Pierre BNM, Ascension NM, Lebel TJ. Chemical composition and biocide properties of *Clausena anisata* (Rutaceae) essential oil against developmental stages of the malaria vector *Anopheles coluzzii*. *Amer J Essent Oil Nat Prod*. 2018;6(1):9–15. <https://www.essencejournal.com/pdf/2018/vol6issue1/PartA/5-4-7-762.pdf>
209. Mollong E, Bawa RA, Boma S, Nuto Y, Amevoin K. Sensitivity of eggs and larvae of *Amblyomma variegatum* (Acarina: Ixodidae) to *Clausena anisata* essential oil mixed in two vegetable oils as vehicle. *Sci J Vet Adv*. 2018;7:237–246. <https://iranjournals.nlai.ir/handle/123456789/773255>
210. Moses JP, Nattudurai G, Baskar K, Arokiyaraj S, Jayakumar M. Efficacy of essential oil from *Clausena anisata* and its impact on biochemical changes of *Sitophilus oryzae*. *Environ Sci Pollut Res Int*. 2020;27:23215–23221. <https://doi.org/10.1007/s11356-020-08928-5>
211. Tabari MA, Rostami A, Khodashenas A, et al. Acaricidal activity, mode of action, and persistent efficacy of selected Eos on the poultry red mite (*Dermanyssus gallinae*). *Food Chem Toxicol*. 2020;138:111207. <https://doi.org/10.1016/j.fct.2020.111207>
212. Nukenine EN, Adler C, Reichmuth C. Efficacy of *Clausena anisata* and *Plectranthus glandulosus* leaf powder against *Prostephanus truncatus* (Coleoptera: Bostrichidae) and two strains of *Sitophilus zeamais* (Coleoptera: Curculionidae) on maize. *J Pest Sci*. 2010;83:181–190.

- <https://doi.org/10.1007/s10340-009-0285-6>
213. Goufo P, Fontem DA, Ngnokam D. Evaluation of plant extracts for tomato late blight control in Cameroon. *New Zealand J Crop Hort Sci.* 2010;38(3):171–176. <https://doi.org/10.1080/01140671.2010.495374>
214. Adamu M, Naidoo V, Eloff JN. Efficacy and toxicity of thirteen plant leaf acetone extracts used in ethnoveterinary medicine in South Africa on egg hatching and larval development of *Haemonchus contortus*. *BMC Vet Res.* 2013;9:38. <https://doi.org/10.1186/1746-6148-9-38>
215. Gebreyesus T, Chapya A. Antifeedants from *Clausena anisata* (Willd.) Hook F. Ex Benth (Rutaceae). *Natural Products for Innovative Pest Management* (Whitehead, DL and Bower, WS eds). Oxford: Pergamon Press; 1983; 237–241. ISBN:0080288936
216. Zheng J, Leong D, Lees G, Nicholson RA. Studies on the interaction of surangin B with insect mitochondria, insect synaptosomes, and rat cortical neurones in primary culture. *Pestic Biochem Physiol.* 1998;61(1):1–13. <https://doi.org/10.1006/pest.1998.2349>
217. Khanikor B, Parida P, Yadav RNS, Bora D. Comparative mode of action of some terpene compounds against octopamine receptor and acetyl cholinesterase of mosquito and human system by the help of homology modeling and docking studies. *J Appl Pharm Sci.* 2013;3(2):6–12. <https://doi.org/10.7324/JAPS.2013.30202>
218. Neal JJ, Wu D. Inhibition of insect cytochromes P450 by furanocoumarins. *Pest Biochem Physiol.* 1994;50(1):43–50. <https://doi.org/10.1006/pest.1994.1056>
219. Flattum RF, Shankland DL. Acetylcholine receptors and the diphasic action of nicotine in the American cockroach, *Periplaneta americana* (L.). *Comp Gen Pharmacol.* 1971;2(6):159–167. [https://doi.org/10.1016/0010-4035\(71\)90006-1](https://doi.org/10.1016/0010-4035(71)90006-1)
220. Amar M, Pichon Y, Inoue I. Micromolar concentrations of veratridine activate sodium channels in embryonic cockroach neurones in culture. *Pflügers Arch.* 1991;417:500–508. <https://doi.org/10.1007/BF00370946>
221. Koul O. Phytochemicals and insect control: an antifeedant approach. *Crit Rev Plant Sci.* 2008;27(1):1–24. <https://doi.org/10.1080/07352680802053908>
222. Eswaremoorthy R, Hailekiros H, Kedir F, Endale M. In silico molecular docking, DFT analysis and ADMET studies of carbazole alkaloid and coumarins from roots of *Clausena anisata*: a potent inhibitor for Quorum Sensing. *Adv Appl Bioinform Chem.* 2021;14:1–12. <https://doi.org/10.214/AABC.S290912>
223. Yusuf H. Acute toxicity test of *Clausena* leaf ethanol extract (*Clausena anisata* Hook.f.). *Syiah Kuala Med J.* 2011;11(1):1–7. <http://www.jurnal.unsyiah.ac.id/JKS/article/view/3439>
224. Moshi MJ, Innocent E, Magadula JJ, et al. Brine shrimp toxicity of some plants used as traditional medicines in Kagera Region, north western Tanzania. *Tanzania J Health Res.* 2010;12(1):63–67. <https://doi.org/10.4314/thrb.v12i1.56287>
225. Emerole G, Thabrew MI, Anosa V, Okorie DA. Structure-activity relationship in the toxicity of some naturally occurring coumarins –chalepin, imperatorin and oxypeucedanine. *Toxicology.* 1981;20(1):71–80. [https://doi.org/10.1016/0300-483X\(81\)90107-4](https://doi.org/10.1016/0300-483X(81)90107-4)
226. Thomford KP, Yorke J, Thomford AK, Amponsah IK. A formulation of *Clausena anisata* (Willd.) Hook. f. Ex Benth and *Cassia sieberiana* DC. alleviates the symptoms associated with osteoarthritis: a single-blind, randomised controlled trial of a traditional Ghanaian remedy. *Clin Phytosci.* 2021;7:76. <https://doi.org/10.1186/s40816-021-00313-0>